

# Apoptosis by dietary agents for prevention and treatment of prostate cancer

Naghma Khan, Vaqar Mustafa Adhami and Hasan Mukhtar

Department of Dermatology, Medical Sciences Center, University of Wisconsin-Madison, 4385, 1300 University Avenue, Madison, Wisconsin 53706, USA

(Correspondence should be addressed to H Mukhtar; Email: hmukhtar@wisc.edu)

## Abstract

Accumulating data clearly indicate that induction of apoptosis is an important event for chemoprevention of cancer by naturally occurring dietary agents. In mammalian cells, apoptosis has been divided into two major pathways: the extrinsic pathway, activated by pro-apoptotic receptor signals at the cellular surface; and the intrinsic pathway, which involves the disruption of mitochondrial membrane integrity. This process is strictly controlled in response to integrity of pro-death signaling and plays critical roles in development, maintenance of homeostasis, and host defense in multicellular organisms. For chemoprevention studies, prostate cancer (PCa) represents an ideal disease due to its long latency, its high incidence, tumor marker availability, and identifiable preneoplastic lesions and risk groups. In this article, we highlight the studies of various apoptosis-inducing dietary compounds for prevention of PCa *in vitro* in cell culture, in preclinical studies in animals, and in human clinical trials.

*Endocrine-Related Cancer* (2010) 17 R39–R52

## Introduction

Prostate cancer (PCa) is one of the most common cancers in men in the United States and is the second leading cause of male cancer death worldwide after lung cancer. The number of new PCa cases expected to be diagnosed in the United States alone in 2009 are 192 280 with an estimated 27 360 disease-related deaths (Jemal *et al.* 2009). PCa is an ideal disease for chemopreventive intervention as it grows slowly before the onset of symptoms and the establishment of diagnosis and it is usually diagnosed in men more than 50 years of age. Therefore, pharmacological or nutritional intervention could considerably impact the quality of life of patients by delaying the progression of cancer (Syed *et al.* 2007).

Multicellular organisms normally eliminate damaged cells effectively through apoptosis, a controlled cellular mechanism resulting in cell death. The concept of physiological cell death was developed by Kerr *et al.* (1972) with the publication of a seminal paper on apoptosis. The term apoptosis is derived from the Greek word describing the falling off of petals from a flower or leaves from a tree. It has become clear that apoptosis is a highly conserved mechanism that

has evolved to maintain cell numbers and cellular positioning within tissues comprised of different cell compartments (Fadeel & Orrenius 2005, Khan *et al.* 2007). It involves the concerted action of a number of intracellular signaling pathways, including members of the caspase family of cysteine proteases, stored in most cells as zymogens or procaspases. Characteristic apoptotic features include cell shrinkage, membrane blebbing, chromatin condensation, and formation of a DNA ladder with multiple fragments caused by internucleosomal DNA cleavage finally ending with the engulfment by macrophages or neighboring cells, thereby avoiding an inflammatory response in surrounding tissues (Savill & Fadok 2000). Cells can undergo apoptosis via two different pathways: the intrinsic or mitochondrial-mediated pathway and the extrinsic or death receptor-mediated pathway.

The intrinsic pathway is usually activated by the loss of growth factor signals or in response to many different damaging influences, for example DNA damage, oxidative stress, hypoxia, or chemotherapeutic drugs. Intrinsically triggered apoptosis is mainly regulated by proteins of the Bcl-2 family, which control the release of pro-apoptotic factors from the mitochondrial intermembrane space. Caspase-9 is

activated when cytochrome *c* is released into the cytoplasm from the mitochondrial intermembranous space. Activated caspase-8 and -9 activate executioner caspases, including caspase-3, which in turn cleave a number of cellular proteins that include structural proteins, nuclear proteins, cytoskeletal proteins, and signaling molecules. The Bcl-2 family of proteins plays a central role in controlling the mitochondrial pathway. More than 20 members of this family have been identified to date in humans, including suppressors (Bcl-2, Bcl-xL, Mcl-1, Bfl-1/A1, Bcl-W, and Bcl-G) and promoters (Bax, Bak, Bok, Bad, Bid, Bik, Bim, Bcl-Xs, Krk, Mtd, Nip3, Nix, Nora, and Bcl-B) of apoptosis (Iannolo *et al.* 2008). In addition to cytochrome *c*, mitochondria release a large number of other polypeptides, including AIF, Endo G, second mitochondrial activator of caspases (Smac/Diablo), and HtrA2/Omi from the intermembrane space. Smac/Diablo and Omi/HtrA2 promote caspase activation through neutralizing the inhibitory effects of inhibitor of apoptosis proteins (IAPs). In addition to cytochrome *c*, mitochondria release a large number of other polypeptides, including AIF, Endo G, Smac/Diablo, and HtrA2/Omi from the intermembrane space. Smac/Diablo and Omi/HtrA2 promote caspase activation through neutralizing the inhibitory effects of IAPs.

The extrinsic pathway is initiated by binding of the transmembrane death receptors such as Fas, tumor necrosis factor (TNF) receptor, DR3, DR4, or DR5 with their specific ligands. These cell surface receptors are activated when cross-linked by their ligands. Activation of death receptors by cross-linking with their natural ligands induces receptor clustering and formation of a death-inducing signaling complex. The complex recruits procaspase-8 via the adaptor molecule Fas-associated death domain protein (FADD), resulting in the activation of caspase-8. Next, procaspase-8 is proteolytically activated and serves as the ‘initiator’ caspase, further activating downstream effectors’ proteins such as caspases-3 and -7 to initiate cell degradation, causing inevitable apoptosis.

For the maintenance of prostate growth, the complex equilibrium between cell growth, proliferation factors, and apoptosis-inducing factors is essential. Fluctuations in this balance cause over-expression of factors causing cell survival and proliferation and loss of apoptosis leading to tumorigenesis and cancer. The deregulation of prostate growth in PCa cells is notable by apoptotic evasion, loss of differentiation, and uncontrolled proliferation. For the treatment of advanced metastatic PCa and the appearance of therapeutic resistance of prostate

tumors, the challenges in the implementation of effective therapeutic strategies involve functional significance of anti-apoptotic pathways (Reynolds & Kyprianou 2006). In PCa progression, loss/suppression of apoptosis has been heavily implicated and apoptosis induction has been considered as an effective means of therapeutic approach for the treatment of prostate tumors. For the effective establishment of the efficacy of the markers of apoptosis in PCa, their evaluation should be performed in combination with numerous clinical parameters and other biochemical markers.


Various research studies have demonstrated that dietary agents may be used alone or in combination with conventional chemotherapeutic agents to prevent the occurrence and spread of cancer. Being rich sources of abundant bioactive compounds, consumption of fruits, vegetables, and whole grains may reduce the risk of cancer in some individuals. In this review, we discuss the use of dietary components present in the mostly consumed fruits, vegetables, and beverages as chemopreventive and/or chemotherapeutic agents against cancer. The rationale for selecting these compounds is that they are present in large amounts in the dietary substances and have been shown to exhibit chemopreventive and/or chemotherapeutic effects against PCa. We summarized studies on the effects of selected dietary components like (–)-epigallocatechin-3-gallate (EGCG), genistein, curcumin, resveratrol, lycopene, pomegranate, fisetin, and lupeol (Fig. 1) on PCa within the context of apoptosis (Fig. 2) *in vitro*, *in vivo*, and where available in human clinical trials.

### (–)-Epigallocatechin-3-gallate

Tea, the most consumed beverage in the world next to water, is derived from the plant *Camellia sinensis* and is processed in different ways in different parts of the world to give green, black, or oolong tea. Green tea contains characteristic polyphenolic compounds, namely, EGCG, (–)-epigallocatechin (EGC), (–)-epicatechin-3-gallate (ECG), and (–)-epicatechin (EC). There is vast amount of scientific literature, which suggests that EGCG is responsible for the majority of the potential health benefits attributed to green tea consumption (Khan *et al.* 2008a).

### *In vitro* studies


We reported for the first time that EGCG (80 μM) induced apoptosis in human PCa DU145 cells (Ahmad *et al.* 1997), and recently reported that treatment of human PCa cells such as LNCaP, PC-3, and



**Figure 1** Chemical structures of dietary agents.

CWR22Rv1 with combination of EGCG (10–40  $\mu\text{M}$ ) and cyclooxygenase-2 (COX-2) inhibitor resulted in enhanced cell growth inhibition, apoptosis induction, and inhibition of NF- $\kappa\text{B}$  (Adhami *et al.* 2007). Study from our laboratory has also shown that EGCG (20 and 40  $\mu\text{M}$ ) sensitizes TRAIL-resistant LNCaP cells to TRAIL-mediated apoptosis through modulation of intrinsic and extrinsic apoptotic pathways (Siddiqui *et al.* 2008a). In our earlier studies, using isogenic cell lines, we have demonstrated that EGCG (20–80  $\mu\text{M}$ )

activates growth arrest and apoptosis in prostate carcinoma cells primarily via p53-dependent pathway that involves the function of both p21 and Bax such that downregulation of either molecule confers a growth advantage to the cells (Hastak *et al.* 2005). It has been shown that EGCG, the major green tea catechin, inhibited the growth of SV40-immortalized human prostate epithelial cells (PNT1A) as well as tumorigenic, poorly differentiated PC-3 cells, but had no effects on normal human prostate epithelial cells



**Figure 2** Effects of dietary agents on the extrinsic and intrinsic pathways of apoptosis in prostate cancer. The extrinsic pathway is triggered by death receptors and involves activation of the initiator caspase-8, which directly activates caspase-3 causing apoptosis. The intrinsic pathway is activated by different apoptotic stimuli that lead to release of cytochrome *c* from mitochondria and activation of caspase-9. This step can be inhibited by the anti-apoptotic members of the Bcl-2 family of apoptosis regulators. Cytochrome *c* interacts with APAF-1 and caspase-9 to promote the activation of caspase-3. Most dietary agents interfere with the key regulators of the apoptotic pathway.

(Caporali *et al.* 2004). It has been shown that EGCG (20–80  $\mu\text{M}$ ) induced apoptosis in human PCa LNCaP cells via stabilization of p53 by phosphorylation on critical serine residues and p14ARF-mediated downregulation of murine double minute 2 (MDM2) protein, negative regulation of NF- $\kappa\text{B}$  activity, activation of caspases, causing a change in the ratio of Bax/Bcl-2 in a manner that favors apoptosis (Hastak *et al.* 2003). The high levels of fatty acid synthase (FAS) activity in LNCaP cells were shown to be dose dependently inhibited by EGCG (40–150  $\mu\text{M}$ ) and it was paralleled by decreased endogenous lipid synthesis, inhibition of cell growth, and induction of apoptosis. Treatment of nonmalignant cells exhibiting low levels of FAS activity with EGCG led to a decrease in growth rate but not to induction of apoptosis. These studies showed that EGCG inhibited FAS activity and selectively caused apoptosis in LNCaP cells but not in nontumoral fibroblasts (Brusselmans *et al.* 2003). EGCG and (+)-gallicocatechin gallate at dose level of 1–100  $\mu\text{M}$  potently and specifically inhibit the chymotrypsin-like activity of purified 20S proteasome and the 26S proteasome in tumor cell lysates. EGCG treatment (10  $\mu\text{M}$ ) also accumulated the pro-apoptotic Bax protein and induced apoptosis in LNCaP cells expressing high basal levels of Bax (Smith *et al.* 2002). Green tea catechins (100  $\mu\text{M}$ ) suppressed the growth and induced apoptosis through an increase in reactive oxygen species formation and mitochondrial depolarization in human PCa DU145 cells (Chung *et al.* 2001). We have shown earlier that EGCG (10–80  $\mu\text{M}$ ) negatively modulates PCa cell growth by the induction of apoptosis, which may be mediated by WAF1/p21-caused  $G_0/G_1$  phase cell cycle arrest, irrespective of the androgen association or p53 status of the cells (Gupta *et al.* 2000).

### **In vivo studies**

We have earlier reported that green tea polyphenols (GTP) containing EGCG consumption significantly inhibited PCa development and metastasis in transgenic adenocarcinoma of the mouse prostate (TRAMP) model. This was achieved by an oral infusion of GTP equivalent to six cups of green tea a day, i.e. at doses that are achievable in humans. GTP consumption was found to cause significant apoptosis of PCa cells, which possibly resulted in reduced dissemination of cancer cells, thereby causing inhibition of PCa development, progression, and metastasis to distant organ sites (Gupta *et al.* 2001). Recently, we have reported that continuous GTP infusion for 32 weeks resulted in substantial reduction in expression of NF- $\kappa\text{B}$ , IKK $\alpha$ ,

IKK $\beta$ , RANK, NIK, STAT-3, and osteopontin in dorsolateral prostate of TRAMP mice. There was also induction of Bax and downregulation of Bcl-2 proteins (Siddiqui *et al.* 2008b). In athymic nude mice, implanted with CWR22Rv1 cells, combination treatment with GTP and celecoxib resulted in enhanced tumor growth inhibition, lowering of prostate-specific antigen (PSA), insulin-like growth factor 1 (IGF1) levels, and increase in IGF-binding protein (IGFBP)-3 levels (Adhami *et al.* 2007). EGCG inhibited early stage PCa, but not late stage PCa, in the TRAMP mice. In the ventrolateral prostate, EGCG significantly reduced cell proliferation, induced apoptosis, and decreased androgen receptor (AR), IGF1, IGF1R, phospho-extracellular signal-regulated kinases 1/2, COX-2, and inducible nitric oxide synthase (Harper *et al.* 2007). It has also been reported previously that treatment of athymic nude mice implanted with CWR22Rv1 cells with GTP, water extract of black tea, and their major constituents, EGCG and theaflavins respectively, resulted in significant inhibition in growth of implanted prostate tumors. This was accompanied with induction of apoptosis as evidenced by upregulation in Bax and decrease in Bcl-2 proteins and decrease in the levels of vascular endothelial growth factor (VEGF) protein (Siddiqui *et al.* 2006). In a study, only 20% of mice developed the neoplasms who were receiving 0.3% GTC in drinking water, while 100% of TRAMP mice developed CaP in the control group. In TRAMP mice, the clusterin (CLU) gene was dramatically downregulated during onset and progression of CaP. In GTC-treated TRAMP mice in which tumor progression was chemoprevented, CLU mRNA and protein progressively accumulated in the prostate gland. CLU dropped again to undetectable levels in animals in which GTC chemoprevention failed and PCa developed (Caporali *et al.* 2004).

### **Clinical trials**

The effects of green tea consumption to various stages of PCa patients have been reported in several studies, but apoptotic effects have not been reported in any of these studies. Recently, the results of a phase II clinical trial have been reported. The PCa patients were given daily doses of Polyphenon E, which contained 800 mg (–)-EGCG and lesser amounts of (–)-EC, (–)-EGC, and (–)-ECG, until time of radical prostatectomy. There was a significant reduction in serum levels of PSA, hepatocyte growth factor, and VEGF in men with PCa after brief treatment with EGCG (Polyphenon E) without the elevation in liver enzymes (McLarty *et al.* 2009). In a clinical trial, men with high-grade prostatic

intraepithelial neoplasia were given either extracts of green tea or a placebo for 1 year. There were no significant side effects and lower urinary tract symptoms with 3% tumor incidence in GTC-treated men as compared to 30% in placebo-treated men (Bettuzzi *et al.* 2006). In patients of hormone-refractory PCa, green tea was found to have minimal clinical activity against the disease with nine patients reported to have progressive disease within 2 months of starting therapy and six patients developed progressive disease after additional 1–4 months of therapy (Choan *et al.* 2005). A case–control study was conducted in Hangzhou, Southeast China, in patients with histologically confirmed adenocarcinoma of the prostate. The risk of PCa declined with increasing frequency, duration, and quantity of green tea consumption (Jian *et al.* 2004). In a phase II clinical trial, decrease in PSA was observed in 2% of cohort, but there was also green tea toxicity in 69% of patients with androgen-independent PCa (Jatoi *et al.* 2003).

## Genistein

Phytoestrogens are naturally occurring phenolic compounds classified as flavones, isoflavones, coumestans, and lignans. The beneficial effects of a soy diet have been attributed to isoflavones. Genistein (5,7,4'-trihydroxyisoflavone), the predominant isoflavone in human diet, is derived mainly from soybeans but is also found in other legumes, including peas, lentils, or beans.

### *In vitro* studies

It has been recently reported that genistein (15–120  $\mu\text{M}$ ) inhibited proliferation and induced apoptosis of DU145 and HeLa cells and had minimal effects on normal L-O2 cells (Yuan-Jing *et al.* 2009). Genistein-combined polysaccharide (GCP) at doses of 1–200  $\mu\text{M}$  has been reported to mediate growth inhibition and promote apoptosis through molecular mimicry of androgen ablation via AR downregulation and by providing an AR-independent, pro-apoptotic signal through mammalian target of rapamycin (mTOR) inhibition (Tepper *et al.* 2007). Genistein (15  $\mu\text{M}$ ) combined with radiation was found to cause greater inhibition in PC-3 colony formation compared to genistein or radiation alone. Treatment with genistein caused dose- and time-dependent  $G_2/M$  phase cell cycle arrest with increased WAF1/p21 and decreased cyclin B1 expression. Radiation-induced activation of NF- $\kappa\text{B}$  activity was strongly inhibited by genistein pretreatment. A significant increase in apoptosis was measured by an increase in cleavage

of poly (ADP-ribose) polymerase (PARP) protein (Raffoul *et al.* 2006). Genistein (15–50  $\mu\text{M}$ ) reduced MDM2 protein and mRNA levels in human cancer cell lines of breast, colon and prostate, primary fibroblasts, and breast epithelial cells. At the post-translational level, genistein induced ubiquitination of MDM2, which led to its degradation. Treatment with genistein also led to induction of apoptosis,  $G_2$  arrest, and inhibition of cell proliferation (Li *et al.* 2005). Both genistein (10–70  $\mu\text{M}$ ) and  $\beta$ -lapachone (1.2  $\mu\text{M}$ ) caused dose-dependent growth inhibition and treatment-induced apoptosis in PC-3 cells. Treatment with caspase-3 inhibitor, DEVD-FMK before exposure to genistein, significantly inhibited caspase-3 expression and treatment-induced apoptosis (Kumi-Diaka *et al.* 2004). Genistein in combination with polysaccharide (10  $\mu\text{M}$ ) significantly suppressed cell growth in LNCaP and PC-3 cells, which was associated with apoptosis in LNCaP cells, but not in PC-3 cells (Bemis *et al.* 2004). Inhibition of the proteasome by genistein (50–200  $\mu\text{M}$ ) was associated with accumulation of ubiquitinated proteins and three known proteasome target proteins, Kip1/p27, I $\kappa\text{B}\alpha$  and the pro-apoptotic protein Bax in PCa LNCaP and breast cancer MCF-7 cells. Genistein-mediated proteasome inhibition was accompanied by induction of apoptosis in these tumor cells (Kazi *et al.* 2003). Polyphenols from tomatoes and soy such as genistein, quercetin, kaempferol, biochanin A, daidzein, and rutin (5–50  $\mu\text{M}$ ) modulated IGF1-induced *in vitro* proliferation and apoptotic resistance in the AT6.3 rat PCa cell line via inhibition of multiple intracellular signaling pathways involving tyrosine kinase activity (Wang *et al.* 2003). A cDNA microarray gene expression profile of genistein (100  $\mu\text{M}$ ) -treated LNCaP cells revealed that the expression of many genes, including survivin, DNA topoisomerase II, cdc-6, and MAPK-6, was downregulated. The glutathione peroxidase (GPx)-1 gene expression level was the most upregulated (Suzuki *et al.* 2002). It was reported that genistein (75–150  $\mu\text{M}$ ) inhibited the growth of LNCaP cells, which was accompanied by the suppression of DNA synthesis and the induction of apoptosis (Onozawa *et al.* 1998).

### *In vivo* studies

In an orthotopic prostate carcinoma model of PC-3 cells in nude mice, genistein combined with prostate tumor irradiation led to a greater control of the growth of the primary tumor and metastasis to lymph nodes than genistein or radiation alone, resulting in greater survival. There was an increase in giant cells,

apoptosis, inflammatory cells, and fibrosis with decreased tumor cell proliferation consistent with increased tumor cell destruction after radiation and genistein treatment (Hillman *et al.* 2004). It was also shown that MDM2 overexpression abrogated genistein-induced apoptosis *in vitro* and that genistein inhibited MDM2 expression and tumor growth in PC-3 xenografts (Li *et al.* 2005). The 2% GCP-supplemented diet significantly slowed LNCaP tumor growth, increased apoptosis, and decreased proliferation over 4 weeks in xenograft model (Bemis *et al.* 2004). Genistein regulated the expression of multiple genes involved in the control of cell growth, apoptosis, and metastasis in PC-3 cells and in experimental PC-3 bone tumors created by injecting PC-3 cells into human bone fragments previously implanted in severe combined immunodeficient mice (Li *et al.* 2004).

### Clinical trials

A nonrandomized, nonblinded trial with historically matched controls from archival tissue was designed to determine the effects of acute exposure to a dietary supplement of isoflavones in men with clinically significant PCa before radical prostatectomy. Prior to surgery, 20 men consumed 160 mg/day of red clover-derived dietary isoflavones, containing a mixture of genistein, daidzein, formononetin, and biochanin A. There were no significant differences in serum PSA, Gleason score, serum testosterone, or biochemical factors between pre- and post-treatment in the subjects. Apoptosis in radical prostatectomy specimens from treated patients was significantly higher than in control subjects, specifically in regions of low to moderate grade (Jarred *et al.* 2002).

### Curcumin

Curcumin (diferuloylmethane) is a major chemical component of turmeric (*Curcuma longa* Linn.) and is used as a spice to give a specific flavor and yellow color to food in the Indian subcontinent. It has been used for centuries in indigenous medicine for the treatment of a variety of inflammatory conditions and other diseases. There are diverse mechanisms that are implicated in the inhibition of tumorigenesis by curcumin and include a combination of anti-inflammatory, anti-oxidant, immunomodulatory, pro-apoptotic, and anti-angiogenic properties via pleiotropic effects on genes and cell-signaling pathways at multiple levels (Khan *et al.* 2008a).

### In vitro studies

Curcumin (30  $\mu$ M) sensitized PC-3 cells to TRAIL by inhibiting Akt-regulated NF- $\kappa$ B and NF- $\kappa$ B-dependent anti-apoptotic Bcl-2, Bcl-xL, and X-linked inhibitor of apoptosis protein (XIAP) (Deeb *et al.* 2007). Curcumin has been shown to enhance the apoptosis-inducing potential of TRAIL in PC-3 cells and LNCaP cells. On treatment with curcumin (20–40  $\mu$ M), the expressions of Bcl-2, Bcl-xL, survivin, and XIAP were inhibited, and there was induction in the expressions of Bax, Bak, PUMA, Bim, Noxa, and death receptors DR4 and DR5 in both cell lines. Treatment of cells with curcumin resulted in activation of caspases-3 and -9 and drop in mitochondrial membrane potential. Combination with TRAIL further led to the enhancement of these events (Shankar *et al.* 2007a). It has also been reported that combination of phenethyl isothiocyanate (PEITC; 10  $\mu$ M) and curcumin (25  $\mu$ M) significantly increased the activity of PARP and cleavage of caspase-3 in correlation with apoptotic cell death. The phosphorylations of I $\kappa$ B $\alpha$  and Akt were significantly attenuated by the combination of PEITC and curcumin. Treatment with PEITC and curcumin caused suppression of epidermal growth factor (EGF) receptor phosphorylation and inhibition of EGF-induced phosphorylation of Akt and induction of phosphatidylinositol 3-kinase (PI3K) in PC-3 cells (Kim *et al.* 2006). Combined curcumin and TRAIL treatment led to induction of apoptosis as observed by accumulation of hypodiploid cells in sub-G<sub>1</sub> phase, enhanced annexin V binding, DNA fragmentation, cleavage of procaspases-3, -8, and -9, truncation of pro-apoptotic Bid, and release of cytochrome *c* from mitochondria (Deeb *et al.* 2005). Curcumin (2–5  $\mu$ M) in combination with radiation showed significant enhancement of radiation-induced clonogenic inhibition and apoptosis in PC-3 cells. However, curcumin in combination with radiation showed inhibition of TNF- $\alpha$ -mediated NF- $\kappa$ B activity resulting in Bcl-2 protein downregulation. Significant activation of cytochrome *c* and caspases-9 and -3 was also observed in cells treated with a combination of curcumin and radiation (Chendil *et al.* 2004). Treatment of PCa cells with curcumin (1–100  $\mu$ M) suppressed both constitutive (DU145) and inducible (LNCaP) NF- $\kappa$ B activation and potentiated TNF-induced apoptosis. Curcumin treatment (50–100  $\mu$ M) induced apoptosis in both cell types, which correlated with the downregulation of the expression of Bcl-2 and Bcl-xL and the activation of procaspase-3 and -8 (Mukhopadhyay *et al.* 2001).

### **In vivo studies**

Recently, curcumin has been shown to inhibit the growth of LNCaP xenografts in nude mice by inducing apoptosis, inhibiting proliferation, and also sensitized tumors to undergo TRAIL-induced apoptosis. Curcumin upregulated the expression of TRAIL-R1/DR4, TRAIL-R2/DR5, Bax, Bak, p21/WAF1, and p27/KIP1, and inhibited the activation of NF- $\kappa$ B and its gene products in xenografted tumors (Shankar *et al.* 2008). Combination of PEITC and curcumin showed stronger tumor growth-inhibitory effects in NCr immunodeficient (nu/nu) mice bearing subcutaneous xenografts of PC-3 cells. There was inhibition of Akt- and NF- $\kappa$ B-signaling pathways contributing to the inhibition of cell proliferation and induction of apoptosis (Khor *et al.* 2006). Curcumin caused a marked decrease in the extent of cell proliferation and a significant increase in the extent of apoptosis as measured by an *in situ* cell death assay in excised tumors of athymic nude mice. A significant decrease in the microvessel density as observed by the CD31 antigen staining was also observed (Dorai *et al.* 2001).

### **Resveratrol**

Resveratrol (3,5,4'-trihydroxystilbene), a naturally occurring phytoalexin found in red wine, grapes, peanuts, mulberries, and in a variety of other plants, has anti-oxidant and anti-inflammatory properties. It was first detected in the dried roots of *Polygonum cuspidatum*, traditionally used in Chinese and Japanese medicines as an anti-inflammatory agent. It exists in two isoforms; *trans*-resveratrol and *cis*-resveratrol, where the *trans*-isomer is the more stable form (Khan *et al.* 2008a,b, Athar *et al.* 2009). It has gained considerable attention because of its potential cancer chemopreventive properties. Resveratrol has attracted considerable attention since 1997, when Jang *et al.* (1997) reported that it inhibits the carcinogenic process at the initiation, promotion, and progression stages. These studies were performed using a panel of cells originating from human lymphocytic leukemia and murine mammary and skin tumors.

### **In vitro studies**

It has been shown that resveratrol (100  $\mu$ M) sensitized docetaxel-resistant tumor cells to TRAIL by blocking CLU expression. It was further indicated that resveratrol acts as an effective tyrosine kinase inhibitor and could inhibit Src and Jak kinases, thus resulting in the loss of Stat1 activation (Sallman *et al.* 2007). Methoxy- and hydroxy-substituted resveratrol derivatives exerted

cytotoxic effects in PC-3, LNCaP, and DU145 human PCa cell lines. The most potent compounds, 3,3',4,4',5,5'-hexahydroxy-stilbene and 3,4,4',5-tetramethoxystilbene, induced apoptosis at concentrations lower than resveratrol and caused cell cycle arrest. It was concluded in this study that introducing additional hydroxy- and methoxy moieties to the stilbene ring of resveratrol was capable of enhancing its cytotoxic and pro-apoptotic effects in hormone-responsive and nonresponsive PCa cells (Horvath *et al.* 2007). Resveratrol pretreatment (50  $\mu$ M) sensitized PC-3 and DU145 cells to agents that specifically target death receptors but not to agents that initiate apoptosis through other mechanisms. Resveratrol also altered the expression of IAPs and Bax, and decreased Akt phosphorylation in PC-3 cells leading to increased caspase activation and apoptosis (Gill *et al.* 2007). Resveratrol (0–30  $\mu$ M) was found to inhibit growth and induced apoptosis in LNCaP cells, but had no effect on normal human prostate epithelial cells. The expression of Bax, Bak, PUMA, Noxa, Bim, TRAIL-R1/DR4, and TRAIL-R2/DR5 was upregulated, and the expression of Bcl-2, Bcl-xL, survivin, and XIAP was down-regulated on treatment with resveratrol. There was also generation of reactive oxygen species, translocation of Bax and p53 to mitochondria, subsequent drop in mitochondrial membrane potential, release of mitochondrial proteins, activation of caspase-3 and -9, and induction of apoptosis on treatment of cells with resveratrol. The dominant negative FADD, caspase-8 siRNA, or *N*-acetyl cysteine inhibited the ability of resveratrol to sensitize TRAIL-resistant LNCaP cells. (Shankar *et al.* 2007b). Resveratrol (1–150  $\mu$ M) induced a decrease in proliferation rate and an increase in apoptosis in LNCaP and PC-3 cells in a dose- and time-dependent manner. The resveratrol-induced apoptosis was mediated by activation of caspases-9 and -3 and a change in the Bax/Bcl-2 ratio (Benitez *et al.* 2007). Treatment of LNCaP cells with resveratrol (1–50  $\mu$ M) was found to result in a significant loss of mitochondrial membrane potential, inhibition in the protein level of anti-apoptotic Bcl-2, and increase in pro-apoptotic members of the Bcl-2 family, i.e. Bax, Bak, Bid, and Bad. Resveratrol caused apoptosis of LNCaP cells via inhibition of PI3K/Akt activation and modulations in Bcl-2 family proteins (Aziz *et al.* 2006). Resveratrol (100  $\mu$ M) is a potent sensitizer of tumor cells for TRAIL-induced apoptosis through p53-independent induction of p21 and p21-mediated cell cycle arrest associated with survivin depletion. Resveratrol-induced G<sub>1</sub> arrest was associated with downregulation of survivin expression and sensitization for TRAIL-induced apoptosis.

Resveratrol-mediated cell cycle arrest followed by survivin depletion and sensitization for TRAIL was impaired in p21-deficient cells. The downregulation of survivin using survivin anti-sense oligonucleotides sensitized cells for TRAIL-induced apoptosis (Fulda & Debatin 2004).

### **In vivo studies**

Resveratrol delayed LNCaP tumor growth in athymic nude mice and inhibited expression of a marker for steroid hormone responses. Exposure to resveratrol also led to increased angiogenesis and inhibition of apoptosis in the xenograft (Wang et al. 2008). An *in vivo* experiment was performed to explore the effect of resveratrol in the TRAMP model, featuring the rat probasin promoter/SV40 T antigen. Resveratrol suppressed PCa growth and induced apoptosis through AR downregulation and suppression of the androgen-responsive glandular kallikrein 11, known to be an ortholog of the human PSA, at the mRNA level without any signs of toxicity (Seeni et al. 2008).

### **Lycopene**

Lycopene is a nonprovitamin A carotenoid that gives red color to tomatoes. Humans and animals depend on dietary sources and do not synthesize lycopene. The dietary sources of lycopene include tomatoes and tomato products, apricots, pink guava, papaya, watermelon, and pink grapefruit. It has been suggested that carotenoids may modulate processes related to mutagenesis, carcinogenesis, cell differentiation, and proliferation (Khan et al. 2008a,b).

### **In vitro studies**

It has been reported recently that treatment of PCa LNCaP and PC-3 cells with lycopene-based agents (100 nM) resulted in mitotic arrest with cells accumulating in G<sub>0</sub>/G<sub>1</sub> phase. There was block in G<sub>1</sub>/S transition mediated by decreased levels of cyclins D1, E, cdk-4, and suppression of retinoblastoma phosphorylation. These responses correlated with decreased IGF1R expression and activation, increased IGFBP-2 expression, and decreased AKT activation. Exposure to lycopene also induced a profound apoptotic response in LNCaP cells (Ivanov et al. 2007). It has been demonstrated that treatment of LNCaP cells with physiologically attainable concentrations of lycopene (0.3–3.0 μM) significantly reduced mitochondrial transmembrane potential, induced the release of mitochondrial cytochrome *c*, and increased annexin V binding, confirming induction of apoptosis

(Hantz et al. 2005). Treatment of PCa DU145 cells with lycopene (8–32 μM) resulted in G<sub>0</sub>/G<sub>1</sub> phase cell cycle arrest and induction of apoptosis in a dose-dependent manner. The rate of apoptosis was 42.4% lower in DU145 cells treated with lycopene as compared with the untreated control cells (Tang et al. 2005).

### **In vivo studies**

In a study, the timing of initiation of micronutrients, and the effect of micronutrient combinations, on PCa development in Lady transgenic model was examined. Transgenic males were administered either i) a control diet, ii) control diet supplemented with human equivalent doses of vitamin E, selenium, and lycopene, or iii) control diet supplemented with vitamin E and selenium. In separate experiments, the combination of vitamin E, selenium, and lycopene was initiated at 4, 8, 20, and 36 weeks of age. There was significant reduction in PCa and liver metastasis when intervention was commenced within 8 weeks of age by a combination diet of vitamin E, selenium, and lycopene. There was a strong correlation between disease-free state with upregulation of the prognostic marker p27/Kip1 and decreased expression of proliferating cell nuclear antigen (PCNA) and significantly increased apoptotic index. A combination of vitamin E and selenium was not effective in preventing PCa, with 84.6% of animals developing PCa and 11.5% developing high-grade prostatic intraepithelial neoplasia. Early commencement of micronutrients combination of vitamin E, selenium, and lycopene was found to be beneficial in reducing PCa. Lycopene was found to be an essential component of the combination and effective for PCa prevention (Venkateswaran et al. 2009). The inhibitory effect of lycopene on the growth rate of DU145 tumor xenografts was studied in BALB/c male nude mice. There was 55.6 and 75.8% inhibition of tumor growth rate in mice treated with 100 and 300 mg/kg lycopene respectively as compared to the control group. Treatment with lycopene caused accumulation of DU145 cells in the G<sub>0</sub>/G<sub>1</sub> phase and apoptosis in a dose-dependent manner. The rate of apoptosis was 42.4% greater in cells treated with 32 μmol/l lycopene than in control group (Tang et al. 2005).

### **Clinical trials**

In a clinical study, thirty-two patients diagnosed by biopsy with PCa were given tomato sauce pasta entrees (30 mg lycopene/day) for 3 weeks before prostatectomy. Thirty-four patients with PCa that served as controls did not consume tomato sauce and underwent

prostatectomy. When tumor areas with the most apoptotic cells were compared in the biopsy and resected prostate tissue, tomato sauce consumption increased apoptotic cells in benign prostate hyperplasia (BPH) and in carcinomas. The apoptotic cell death in carcinomas increased significantly with treatment, and apoptotic cell death in BPH showed a tendency towards an increase when comparable morphological areas were counted. The differences in values were not significant in BPH and carcinomas when the values of apoptotic cells in BPH and carcinomas of patients who consumed tomato sauce were compared with corresponding control lesions of the patients who did not consume tomato sauce in resected prostate tissue. This study provides the evidence that tomato sauce consumption may suppress the progression of the disease in a subset of patients with PCa by increasing apoptotic cell death (Kim *et al.* 2003).

## Pomegranate

The pomegranate (*Punica granatum* L.) fruit has been used for centuries in ancient cultures for medicinal purposes. The fruits have also been long and widely used in folk and traditional medicine for the treatment of a number of pathologies (Aviram *et al.* 2000). Employing Matrix-assisted laser desorption/ionization time of flight mass spectrometry, pomegranate juice was found to contain six anthocyanins (pelargonidin 3-glucoside, cyanidin 3-glucoside, delphinidin 3-glucoside, pelargonidin 3,5-diglucoside, cyanidin 3,5-diglucoside, and delphinidin 3,5-diglucoside), ellagitannins, and hydrolyzable tannins. The other flavonoids identified included quercetin, kaempferol, and luteolin glycosides (Gil *et al.* 2000).

### *In vitro* studies

We have shown that pomegranate fruit extract (PFE) treatment (10–100  $\mu$ M) of human PCa PC-3 cells resulted in a dose-dependent inhibition of cell growth/cell viability and induction of apoptosis. PFE treatment of PC-3 cells resulted in induction of Bax and Bak (pro-apoptotic) proteins and downregulation of Bcl-xL and Bcl-2 (anti-apoptotic) proteins (Malik *et al.* 2005). It has been reported that pomegranate extract inhibited NF- $\kappa$ B and cell viability of PCa cell lines in a dose-dependent fashion *in vitro*. Pomegranate extract-induced apoptosis was dependent on NF- $\kappa$ B blockade (Rettig *et al.* 2008). Pomegranate oil (35  $\mu$ M), fermented juice polyphenols, and pericarp polyphenols each acutely inhibited *in vitro* proliferation of LNCaP, PC-3, and DU145 human PCa cell

lines. These effects were mediated by changes in both cell cycle distribution and induction of apoptosis (Albrecht *et al.* 2004). Oral administration of PFE to athymic nude mice implanted with androgen-sensitive CWR22Rv1 cells resulted in a significant inhibition of tumor growth concomitant with a significant decrease in serum PSA levels (Malik *et al.* 2005).

### *In vivo* studies

In the LAPC4 xenograft model, pomegranate extract delayed the emergence of LAPC4 androgen-independent xenografts in castrated mice through an inhibition of proliferation and induction of apoptosis. On treatment with pomegranate extract, the observed increase in NF- $\kappa$ B activity during the transition from androgen dependence to androgen independence in the LAPC4 xenograft model was abrogated (Rettig *et al.* 2008).

### Clinical trials

To determine the effects of pomegranate juice consumption, a phase II clinical trial was conducted in men with rising PSA after surgery or radiotherapy. Mean PSA doubling time significantly increased with treatment from a mean of 15 months at baseline to 54 months post-treatment. There was a decrease in cell proliferation and an increase in apoptosis in *in vitro* assays comparing pretreatment and post-treatment patient serum on the growth of LNCaP cells (Pantuck *et al.* 2006).

## Fisetin

Fisetin (3,3',4',7-tetrahydroxyflavone) is found in fruits and vegetables, such as strawberry, apple, persimmon, grape, onion, and cucumber (Arai *et al.* 2000). Fisetin exhibits a wide variety of activities including neurotrophic (Maher 2006), anti-oxidant (Hanneken *et al.* 2006), anti-inflammatory (Higa *et al.* 2003), and anti-angiogenic (Fotsis *et al.* 1998) effects.

### *In vitro* studies

We have reported that fisetin treatment (10–60  $\mu$ M) was found to result in a decrease in the viability of LNCaP, CWR22Rv1, and PC-3 cells but had only minimal effects on normal prostate epithelial cells. Treatment of LNCaP cells with fisetin resulted in induction of apoptosis, PARP cleavage, modulation in the expressions of Bcl-2-family proteins, inhibition of PI3K, and phosphorylation of Akt at Ser<sup>473</sup> and

Thr<sup>308</sup>. There was also induction of mitochondrial release of cytochrome *c* into cytosol, downregulation of XIAP, and upregulation of Smac/Diablo upon treatment of cells with fisetin. Treatment of cells with fisetin also resulted in significant activation of caspases-3, -8, and -9. Pretreatment of cells with caspase inhibitor (Z-VAD-FMK) blocked fisetin-induced activation of caspases (Khan et al. 2008b).

### Lupeol

Lup-20(29)-en-3h-ol (Lupeol), a triterpene found in fruits such as olive, mango, strawberry, grapes and figs, in many vegetables, and in several medicinal plants, is used in the treatment of various diseases. It possesses strong anti-inflammatory, anti-arthritis, anti-mutagenic, and anti-malarial activity when examined in *in vitro* and *in vivo* systems (Geetha & Varalakshmi 2001).

#### *In vitro* studies

Study from our laboratory has shown that lupeol (1–30 μM) induced the cleavage of PARP protein and degradation of acinus protein with a significant increase in the expression of FADD protein and Fas receptor in androgen-sensitive human PCa cells. The small interfering RNA-mediated silencing of the *Fas* gene and inhibition of caspases-6, -8, and -9 by their specific inhibitors confirmed that Lupeol specifically activates the Fas receptor-mediated apoptotic pathway in androgen-sensitive PCa cells. The treatment of cells with a combination of anti-Fas monoclonal antibody and lupeol resulted in higher cell death compared with the additive effect of the two compounds alone, suggesting a synergistic effect. Treatment with lupeol also resulted in a significant inhibition in growth of tumors with concomitant reduction in PSA secretion in athymic nude mice implanted with CWR22Rv1 cells (Saleem et al. 2005). It has been shown recently that lupeol (400–600 μM) caused anti-proliferative effect associated with an increase in G<sub>2</sub>/M phase arrest in PC-3 cells. There was also induction of apoptosis with upregulation of Bax, caspases-3 and -9, and *APAF-1* genes, and downregulation of anti-apoptotic *Bcl-2* gene in PC-3 cells. The role of caspase-induced apoptosis was confirmed by increase in reactive oxygen species, loss of mitochondrial membrane potential followed by DNA fragmentation (Prasad et al. 2008a; Table 1).

**Table 1** Concentrations of dietary agents reported to cause anti-proliferative and apoptotic effects *in vitro* in prostate cancer cells

Dietary agents	<i>In vitro</i> concentrations (μM)	References
EGCG	80	Ahmad et al. (1997)
	10–80	Gupta et al. (2000)
	1–100	Smith et al. (2002)
	40–150	Brusselmans et al. (2003)
	20–80	Hastak et al. (2003, 2005)
	10–40	Adhami et al. (2007)
	20 and 40	Siddiqui et al. (2008a)
Genistein	75–150	Onozawa et al. (1998)
	100	Suzuki et al. (2002)
	5–50	Wang et al. (2003)
	50–200	Kazi et al. (2003)
	10	Bemis et al. (2004)
	10–70	Kumi-Diaka et al. (2004)
	15–50	Li et al. (2005)
Curcumin	15	Raffoul et al. (2006)
	15–20	Yuan-Jing et al. (2009)
	1–100	Mukhopadhyay et al. (2001)
	2–5	Chendil et al. (2004)
	25	Kim et al. (2006)
	20–40	Shankar et al. (2007a)
	30	Deeb et al. (2007)
Resveratrol	100	Fulda & Debatin (2004)
	1–50	Aziz et al. (2006)
	1–150	Benitez et al. (2007)
	0–30	Shankar et al. (2007b)
	50	Gill et al. (2007)
	100	Sallman et al. (2007)
	Lycopene	0.3–3.0
8–32		Tang et al. (2005)
0.1		Ivanov et al. (2007)
Pomegranate	35	Albrecht et al. (2004)
	10–100	Malik et al. (2005)
Fisetin	10–60	Khan et al. (2008b)
Lupeol	1–30	Saleem et al. (2005)
	400–600	Prasad et al. (2008a)

#### *In vivo* studies

In a recent study, lupeol and mango pulp extract supplementation has resulted in a significantly high percentage of apoptotic cells in the hypodiploid region in male Swiss albino mice. The induction of apoptosis in mouse prostate was preceded by the loss of mitochondrial transmembrane potential and DNA laddering. In LNCaP cells, lupeol caused early increase in reactive oxygen species followed by induction of mitochondrial pathway leading to cell death (Prasad et al. 2008b).

### Conclusions and perspectives

Apoptosis is strictly controlled in response to integrity of pro-death signaling and plays critical roles in the development, maintenance of homeostasis, and host defense in multicellular organisms.

In preclinical and clinical settings, various therapeutic approaches to target diseases by regulating apoptosis are being developed. Approaches include the traditional use of small molecules to target specific players in the apoptosis cascade. With the increase in the understanding of apoptosis, additional opportunities will become available for tailor-made therapies that will result in improved therapies. It is of critical importance that the balance between cell death and proliferation is strongly regulated. Consequently, an enormous interest has emerged for devising therapeutic strategies to modulate key molecules involved in apoptosis. Several dietary compounds have been shown to affect the process of apoptosis *in vitro* and *in vivo*. Identifying the key proteins involved in apoptosis represents an attractive way to prevent the development of many diseases including PCa. Understanding how these proteins affect the apoptotic pathways may lead to more effective cancer treatments.

To translate the *in vitro* efficacy of dietary agents in the prevention of cancer to clinical use, attention should be given to physiologically relevant concentrations and chronic exposures to imitate *in vivo* conditions. To optimize the desired physiological response, further consideration should be given to the significant intake of dietary components, their duration, and their validation in suitable animal models.

### Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

### Funding

The original work from the author's (H Mukhtar) laboratory outlined in this review was supported by United States Public Health Service Grants RO1 CA 78809, RO1 CA 101039, RO1 CA 120451, and P50 DK065303.

### References

Adhami VM, Malik A, Zaman N, Sarfaraz S, Siddiqui IA, Syed DN, Afaq F, Pasha FS, Saleem M & Mukhtar H 2007 Combined inhibitory effects of green tea polyphenols and selective cyclooxygenase-2 inhibitors on the growth of human prostate cancer cells both *in vitro* and *in vivo*. *Clinical Cancer Research* **13** 1611–1619.

Ahmad N, Feyes DK, Nieminen AL, Agarwal R & Mukhtar H 1997 Green tea constituent epigallocatechin-3-gallate and induction of apoptosis and cell cycle arrest in human carcinoma cells. *Journal of the National Cancer Institute* **89** 1881–1886.

Albrecht M, Jiang W, Kumi-Diaka J, Lansky EP, Gommersall LM, Patel A, Mansel RE, Neeman I, Geldof AA & Campbell MJ 2004 Pomegranate extracts potently suppress proliferation, xenograft growth, and invasion of human prostate cancer cells. *Journal of Medicinal Food* **7** 274–283.

Arai Y, Watanabe S, Kimira M, Shimoi K, Mochizuki R & Kinai N 2000 Dietary intakes of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration. *Journal of Nutrition* **130** 2243–2250.

Athar M, Back JH, Kopelovich L, Bickers DR & Kim AL 2009 Multiple molecular targets of resveratrol: anti-carcinogenic mechanisms. *Archives of Biochemistry and Biophysics* **486** 95–102.

Aviram M, Dornfeld L, Rosenblat M, Volkova N, Kaplan M, Coleman R, Hayek T, Presser D & Fuhrman B 2000 Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL, and platelet aggregation: studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *American Journal of Clinical Nutrition* **71** 1062–1076.

Aziz MH, Nihal M, Fu VX, Jarrard DF & Ahmad N 2006 Resveratrol-caused apoptosis of human prostate carcinoma LNCaP cells is mediated via modulation of phosphatidylinositol 3'-kinase/Akt pathway and Bcl-2 family proteins. *Molecular Cancer Therapeutics* **5** 1335–1341.

Bemis DL, Capodice JL, Desai M, Buttyan R & Katz AE 2004 A concentrated aglycone isoflavone preparation (GCP) that demonstrates potent anti-prostate cancer activity *in vitro* and *in vivo*. *Clinical Cancer Research* **10** 5282–5292.

Benitez DA, Pozo-Guisado E, Alvarez-Barrientos A, Fernandez-Salguero PM & Castellon EA 2007 Mechanisms involved in resveratrol-induced apoptosis and cell cycle arrest in prostate cancer-derived cell lines. *Journal of Andrology* **28** 282–293.

Bettuzzi S, Brausi M, Rizzi F, Castagnetti G, Peracchia G & Corti A 2006 Chemoprevention of human prostate cancer by oral administration of green tea catechins in volunteers with high-grade prostate intraepithelial neoplasia: a preliminary report from a one-year proof-of-principle study. *Cancer Research* **66** 1234–1240.

Brusselmans K, De Schrijver E, Heyns W, Verhoeven G & Swinnen JV 2003 Epigallocatechin-3-gallate is a potent natural inhibitor of fatty acid synthase in intact cells and selectively induces apoptosis in prostate cancer cells. *International Journal of Cancer* **106** 856–862.

Caporali A, Davalli P, Astancolle S, D'Arca D, Brausi M, Bettuzzi S & Corti A 2004 The chemopreventive action of catechins in the TRAMP mouse model of prostate carcinogenesis is accompanied by clusterin over-expression. *Carcinogenesis* **25** 2217–2224.

Chendil D, Ranga RS, Meigooni D, Sathishkumar S & Ahmed MM 2004 Curcumin confers radiosensitizing effect in prostate cancer cell line PC-3. *Oncogene* **23** 1599–1607.

- Choan E, Segal R, Jonker D, Malone S, Reaume N, Eapen L & Gallant V 2005 A prospective clinical trial of green tea for hormone refractory prostate cancer: an evaluation of the complementary/alternative therapy approach. *Urologic Oncology* **23** 108–113.
- Chung LY, Cheung TC, Kong SK, Fung KP, Choy YM, Chan ZY & Kwok TT 2001 Induction of apoptosis by green tea catechins in human prostate cancer DU145 cells. *Life Sciences* **68** 1207–1214.
- Deeb DD, Jiang H, Gao X, Divine G, Dulchavsky SA & Gautam SC 2005 Chemosensitization of hormone-refractory prostate cancer cells by curcumin to TRAIL-induced apoptosis. *Journal of Experimental Therapeutics and Oncology* **5** 81–91.
- Deeb D, Jiang H, Gao X, Al-Holou S, Danyluk AL, Dulchavsky SA & Gautam SC 2007 Curcumin [1,7-bis(4-hydroxy-3-methoxyphenyl)-1-6-heptadine-3,5-dione; C21H20O6] sensitizes human prostate cancer cells to tumor necrosis factor-related apoptosis-inducing ligand/Apo2L-induced apoptosis by suppressing nuclear factor-kappaB via inhibition of the prosurvival Akt signaling pathway. *Journal of Pharmacology and Experimental Therapeutics* **321** 616–625.
- Dorai T, Cao YC, Dorai B, Buttyan R & Katz AE 2001 Therapeutic potential of curcumin in human prostate cancer. III. Curcumin inhibits proliferation, induces apoptosis, and inhibits angiogenesis of LNCaP prostate cancer cells *in vivo*. *Prostate* **47** 293–303.
- Fadeel B & Orrenius S 2005 Apoptosis: a basic biological phenomenon with wide-ranging implications in human disease. *Journal of Internal Medicine* **258** 479–517.
- Fotsis T, Pepper MS, Montesano R, Aktas E, Breit S, Schweigerer L, Rasku S, Wahala K & Adlercreutz H 1998 Phytoestrogens and inhibition of angiogenesis. *Bailliere's Clinical Endocrinology and Metabolism* **12** 649–666.
- Fulda S & Debatin KM 2004 Sensitization for tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis by the chemopreventive agent resveratrol. *Cancer Research* **64** 337–346.
- Geetha T & Varalakshmi P 2001 Anti-inflammatory activity of lupeol and lupeol linoleate in rats. *Journal of Ethnopharmacology* **76** 77–80.
- Gil MI, Tomas-Barberan FA, Hess-Pierce B, Holcroft DM & Kader AA 2000 Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *Journal of Agricultural and Food Chemistry* **48** 4581–4589.
- Gill C, Walsh SE, Morrissey C, Fitzpatrick JM & Watson RW 2007 Resveratrol sensitizes androgen independent prostate cancer cells to death-receptor mediated apoptosis through multiple mechanisms. *Prostate* **67** 1641–1653.
- Gupta S, Ahmad N, Nieminen AL & Mukhtar H 2000 Growth inhibition, cell-cycle dysregulation, and induction of apoptosis by green tea constituent (–)-epigallocatechin-3-gallate in androgen-sensitive and androgen-insensitive human prostate carcinoma cells. *Toxicology and Applied Pharmacology* **164** 82–90.
- Gupta S, Hastak K, Ahmad N, Lewin JS & Mukhtar H 2001 Inhibition of prostate carcinogenesis in TRAMP mice by oral infusion of green tea polyphenols. *PNAS* **98** 10350–10355.
- Hanneken A, Lin FF, Johnson J & Maher P 2006 Flavonoids protect human retinal pigment epithelial cells from oxidative-stress-induced death. *Investigative Ophthalmology & Visual Science* **47** 3164–3177.
- Hantz HL, Young LF & Martin KR 2005 Physiologically attainable concentrations of lycopene induce mitochondrial apoptosis in LNCaP human prostate cancer cells. *Experimental Biology and Medicine* **230** 171–179.
- Harper CE, Patel BB, Wang J, Eltoum IA & Lamartiniere CA 2007 Epigallocatechin-3-Gallate suppresses early stage, but not late stage prostate cancer in TRAMP mice: mechanisms of action. *Prostate* **67** 1576–1589.
- Hastak K, Gupta S, Ahmad N, Agarwal MK, Agarwal ML & Mukhtar H 2003 Role of p53 and NF-kappaB in epigallocatechin-3-gallate-induced apoptosis of LNCaP cells. *Oncogene* **22** 4851–4859.
- Hastak K, Agarwal MK, Mukhtar H & Agarwal ML 2005 Ablation of either p21 or Bax prevents p53-dependent apoptosis induced by green tea polyphenol epigallocatechin-3-gallate. *FASEB Journal* **19** 789–791.
- Higa S, Hirano T, Kotani M, Matsumoto M, Fujita A, Suemura M, Kawase I & Tanaka T 2003 Fisetin, a flavonol, inhibits TH2-type cytokine production by activated human basophils. *Journal of Allergy and Clinical Immunology* **111** 1299–1306.
- Hillman GG, Wang Y, Kucuk O, Che M, Doerge DR, Yudelev M, Joiner MC, Marples B, Forman JD & Sarkar FH 2004 Genistein potentiates inhibition of tumor growth by radiation in a prostate cancer orthotopic model. *Molecular Cancer Therapeutics* **3** 1271–1279.
- Horvath Z, Marihart-Fazekas S, Saiko P, Grusch M, Ozsuy M, Harik M, Handler N, Erker T, Jaeger W, Fritzer-Szekeres M *et al.* 2007 Novel resveratrol derivatives induce apoptosis and cause cell cycle arrest in prostate cancer cell lines. *Anticancer Research* **27** 3459–3464.
- Iannolo G, Conticello C, Memeo L & De Maria R 2008 Apoptosis in normal and cancer stem cells. *Critical Review in Oncology and Hematology* **66** 42–51.
- Ivanov NI, Cowell SP, Brown P, Rennie PS, Guns ES & Cox ME 2007 Lycopene differentially induces quiescence and apoptosis in androgen-responsive and -independent prostate cancer cell lines. *Clinical Nutrition* **26** 252–263.
- Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CW, Fong HH, Farnsworth NR, Kinghorn AD, Mehta RG *et al.* 1997 Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* **275** 218–220.
- Jarred RA, Keikha M, Dowling C, McPherson SJ, Clare AM, Husband AJ, Pedersen JS, Frydenberg M & Risbridger GP 2002 Induction of apoptosis in low to moderate-grade human prostate carcinoma by red clover-derived dietary isoflavones. *Cancer Epidemiology, Biomarkers and Prevention* **11** 1689–1696.

- Jatoi A, Ellison N, Burch PA, Sloan JA, Dakhil SR, Novotny P, Tan W, Fitch TR, Rowland KM, Young CY *et al.* 2003 A phase II trial of green tea in the treatment of patients with androgen independent metastatic prostate carcinoma. *Cancer* **97** 1442–1446.
- Jemal A, Siegel R, Ward E, Hao Y, Xu J & Thun MJ 2009 Cancer statistics, 2009. *CA: A Cancer Journal for Clinicians* **59** 225–249.
- Jian L, Xie LP, Lee AH & Binns CW 2004 Protective effect of green tea against prostate cancer: a case–control study in Southeast China. *International Journal of Cancer* **108** 130–135.
- Kazi A, Daniel KG, Smith DM, Kumar NB & Dou QP 2003 Inhibition of the proteasome activity, a novel mechanism associated with the tumor cell apoptosis-inducing ability of genistein. *Biochemical Pharmacology* **66** 965–976.
- Kerr JF, Wyllie AH & Currie AR 1972 Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *British Journal of Cancer* **26** 239–257.
- Khan N, Afaq F & Mukhtar H 2007 Apoptosis by dietary factors: the suicide solution for delaying cancer growth. *Carcinogenesis* **28** 233–239.
- Khan N, Afaq F & Mukhtar H 2008a Cancer chemoprevention through dietary antioxidants: progress and promise. *Antioxidants & Redox Signaling* **10** 475–510.
- Khan N, Afaq F, Syed DN & Mukhtar H 2008b Fisetin, a novel dietary flavonoid, causes apoptosis and cell cycle arrest in human prostate cancer LNCaP cells. *Carcinogenesis* **29** 1049–1056.
- Khor TO, Keum YS, Lin W, Kim JH, Hu R, Shen G, Xu C, Gopalakrishnan A, Reddy B, Zheng X *et al.* 2006 Combined inhibitory effects of curcumin and phenethyl isothiocyanate on the growth of human PC-3 prostate xenografts in immunodeficient mice. *Cancer Research* **66** 613–621.
- Kim HS, Bowen P, Chen L, Duncan C, Ghosh L, Sharifi R & Christov K 2003 Effects of tomato sauce consumption on apoptotic cell death in prostate benign hyperplasia and carcinoma. *Nutrition and Cancer* **47** 40–47.
- Kim JH, Xu C, Keum YS, Reddy B, Conney A & Kong AN 2006 Inhibition of EGFR signaling in human prostate cancer PC-3 cells by combination treatment with beta-phenylethyl isothiocyanate and curcumin. *Carcinogenesis* **27** 475–482.
- Kumi-Diaka J, Saddler-Shawnette S, Aller A & Brown J 2004 Potential mechanism of phytochemical-induced apoptosis in human prostate adenocarcinoma cells: therapeutic synergy in genistein and beta-lapachone combination treatment. *Cancer Cell International* **4** 5.
- Li Y, Che M, Bhagat S, Ellis KL, Kucuk O, Doerge DR, Abrams J, Cher ML & Sarkar FH 2004 Regulation of gene expression and inhibition of experimental prostate cancer bone metastasis by dietary genistein. *Neoplasia* **6** 354–363.
- Li M, Zhang Z, Hill DL, Chen X, Wang H & Zhang R 2005 Genistein, a dietary isoflavone, down-regulates the MDM2 oncogene at both transcriptional and posttranslational levels. *Cancer Research* **65** 8200–8208.
- Maher P 2006 A comparison of the neurotrophic activities of the flavonoid fisetin and some of its derivatives. *Free Radical Research* **40** 1105–1111.
- Malik A, Afaq F, Sarfaraz S, Adhami VM, Syed DN & Mukhtar H 2005 Pomegranate fruit juice for chemoprevention and chemotherapy of prostate cancer. *PNAS* **102** 14813–14818.
- McLarty J, Bigelow RL, Smith M, Elmajian D, Ankem M & Cardelli JA 2009 Tea polyphenols decrease serum levels of prostate-specific antigen, hepatocyte growth factor, and vascular endothelial growth factor in prostate cancer patients and inhibit production of hepatocyte growth factor and vascular endothelial growth factor *in vitro*. *Cancer Prevention Research* **2** 673–682.
- Mukhopadhyay A, Bueso-Ramos C, Chatterjee D, Pantazis P & Aggarwal BB 2001 Curcumin downregulates cell survival mechanisms in human prostate cancer cell lines. *Oncogene* **20** 7597–7609.
- Onozawa M, Fukuda K, Ohtani M, Akaza H, Sugimura T & Wakabayashi K 1998 Effects of soybean isoflavones on cell growth and apoptosis of the human prostatic cancer cell line LNCaP. *Japanese Journal of Clinical Oncology* **28** 360–363.
- Pantuck AJ, Leppert JT, Zomorodian N, Aronson W, Hong J, Barnard RJ, Seeram N, Liker H, Wang H, Elashoff R *et al.* 2006 Phase II study of pomegranate juice for men with rising prostate-specific antigen following surgery or radiation for prostate cancer. *Clinical Cancer Research* **12** 4018–4026.
- Prasad S, Nigam N, Kalra N & Shukla Y 2008a Regulation of signaling pathways involved in lupeol induced inhibition of proliferation and induction of apoptosis in human prostate cancer cells. *Molecular Carcinogenesis* **47** 916–924.
- Prasad S, Kalra N & Shukla Y 2008b Induction of apoptosis by lupeol and mango extract in mouse prostate and LNCaP cells. *Nutrition and Cancer* **60** 120–130.
- Raffoul JJ, Wang Y, Kucuk O, Forman JD, Sarkar FH & Hillman GG 2006 Genistein inhibits radiation-induced activation of NF-kappaB in prostate cancer cells promoting apoptosis and G<sub>2</sub>/M cell cycle arrest. *BMC Cancer* **6** 107.
- Rettig MB, Heber D, An J, Seeram NP, Rao JY, Liu H, Klatter T, Belldegrun A, Moro A, Henning SM *et al.* 2008 Pomegranate extract inhibits androgen-independent prostate cancer growth through a nuclear factor-kappaB-dependent mechanism. *Molecular Cancer Therapeutics* **7** 2662–2671.
- Reynolds AR & Kyprianou N 2006 Growth factor signalling in prostatic growth: significance in tumour development and therapeutic targeting. *British Journal of Pharmacology* **147** (Suppl 2) S144–S152.
- Saleem M, Kweon MH, Yun JM, Adhami VM, Khan N, Syed DN & Mukhtar H 2005 A novel dietary triterpene Lupeol induces fas-mediated apoptotic death of androgen-sensitive prostate cancer cells and inhibits tumor growth in a xenograft model. *Cancer Research* **65** 11203–11213.

- Sallman DA, Chen X, Zhong B, Gilvary DL, Zhou J, Wei S & Djeu JY 2007 Clusterin mediates TRAIL resistance in prostate tumor cells. *Molecular Cancer Therapeutics* **6** 2938–2947.
- Savill J & Fadok V 2000 Corpse clearance defines the meaning of cell death. *Nature* **407** 784–788.
- Seeni A, Takahashi S, Takeshita K, Tang M, Sugiura S, Sato SY & Shirai T 2008 Suppression of prostate cancer growth by resveratrol in the transgenic rat for adenocarcinoma of prostate (TRAP) model. *Asian Pacific Journal of Cancer Prevention* **9** 7–14.
- Shankar S, Chen Q, Sarva K, Siddiqui I & Srivastava RK 2007a Curcumin enhances the apoptosis-inducing potential of TRAIL in prostate cancer cells: molecular mechanisms of apoptosis, migration and angiogenesis. *Journal of Molecular Signaling* **2** 10.
- Shankar S, Chen Q, Siddiqui I, Sarva K & Srivastava RK 2007b Sensitization of TRAIL-resistant LNCaP cells by resveratrol (3,4',5 tri-hydroxystilbene): molecular mechanisms and therapeutic potential. *Journal of Molecular Signaling* **2** 7.
- Shankar S, Ganapathy S, Chen Q & Srivastava RK 2008 Curcumin sensitizes TRAIL-resistant xenografts: molecular mechanisms of apoptosis, metastasis and angiogenesis. *Molecular Cancer* **7** 16.
- Siddiqui IA, Zaman N, Aziz MH, Reagan-Shaw SR, Sarfaraz S, Adhami VM, Ahmad N, Raisuddin S & Mukhtar H 2006 Inhibition of CWR22Rnu1 tumor growth and PSA secretion in athymic nude mice by green and black teas. *Carcinogenesis* **27** 833–839.
- Siddiqui IA, Malik A, Adhami VM, Asim M, Hafeez BB, Sarfaraz S & Mukhtar H 2008a Green tea polyphenol EGCG sensitizes human prostate carcinoma LNCaP cells to TRAIL-mediated apoptosis and synergistically inhibits biomarkers associated with angiogenesis and metastasis. *Oncogene* **27** 2055–2063.
- Siddiqui IA, Shukla Y, Adhami VM, Sarfaraz S, Asim M, Hafeez BB & Mukhtar H 2008b Suppression of NFkappaB and its regulated gene products by oral administration of green tea polyphenols in an autochthonous mouse prostate cancer model. *Pharmaceutical Research* **25** 2135–2142.
- Smith DM, Wang Z, Kazi A, Li LH, Chan TH & Dou QP 2002 Synthetic analogs of green tea polyphenols as proteasome inhibitors. *Molecular Medicine* **8** 382–392.
- Suzuki K, Koike H, Matsui H, Ono Y, Hasumi M, Nakazato H, Okugi H, Sekine Y, Oki K, Ito K *et al.* 2002 Genistein, a soy isoflavone, induces glutathione peroxidase in the human prostate cancer cell lines LNCaP and PC-3. *International Journal of Cancer* **99** 846–852.
- Syed DN, Khan N, Afaq F & Mukhtar H 2007 Chemoprevention of prostate cancer through dietary agents: progress and promise. *Cancer Epidemiology, Biomarkers and Prevention* **16** 2193–2203.
- Tang L, Jin T, Zeng X & Wang JS 2005 Lycopene inhibits the growth of human androgen-independent prostate cancer cells *in vitro* and in BALB/c nude mice. *Journal of Nutrition* **135** 287–290.
- Tepper CG, Vinal RL, Wee CB, Xue L, Shi XB, Burich R, Mack PC & de Vere White RW 2007 GCP-mediated growth inhibition and apoptosis of prostate cancer cells via androgen receptor-dependent and -independent mechanisms. *Prostate* **67** 521–535.
- Venkateswaran V, Klotz LH, Ramani M, Sugar LM, Jacob LE, Nam RK & Fleshner NE 2009 A combination of micronutrients is beneficial in reducing the incidence of prostate cancer and increasing survival in the Lady transgenic model. *Cancer Prevention Research* **2** 473–483.
- Wang S, DeGross VL & Clinton SK 2003 Tomato and soy polyphenols reduce insulin-like growth factor-I-stimulated rat prostate cancer cell proliferation and apoptotic resistance *in vitro* via inhibition of intracellular signaling pathways involving tyrosine kinase. *Journal of Nutrition* **133** 2367–2376.
- Wang TT, Hudson TS, Wang TC, Remsberg CM, Davies NM, Takahashi Y, Kim YS, Seifried H, Vinyard BT, Perkins SN *et al.* 2008 Differential effects of resveratrol on androgen-responsive LNCaP human prostate cancer cells *in vitro* and *in vivo*. *Carcinogenesis* **29** 2001–2010.
- Yuan-Jing F, Nan-Shan H & Lian X 2009 Genistein synergizes with RNA interference inhibiting survivin for inducing DU-145 of prostate cancer cells to apoptosis. *Cancer Letters* **284** 189–197.