SOCS-3 antagonises the proliferative and migratory effects of fibroblast growth factor-2 in prostate cancer by inhibition of p44/p42 MAPK signalling

Martin Puhr, Frédéric R Santer, Hannes Neuwirt, Gemma Marcias, Alfred Hobisch¹ and Zoran Culig

Department of Urology, Innsbruck Medical University, Anichstrasse 35, A-6020 Innsbruck, Austria

¹Department of Urology, General Hospital Feldkirch, A-6800 Feldkirch, Austria

(Correspondence should be addressed to Z Culig; Email: zoran.culig@i-med.ac.at)

Abstract

Fibroblast growth factor-2 (FGF-2) is highly expressed in prostate cancer. It promotes tumour progression through multiple pathways including those of signal transducers and activators of transcription factor 3 (STAT3), mitogen-activated protein kinases (MAPKs) and Akt. In previous studies, we have reported that STAT3 phosphorylation inversely correlates with suppressor of cytokine signalling-3 (SOCS-3) expression in prostate cancer cells. Recently, it has become evident that SOCS-3-negative regulation is not only limited to the interleukin-6 (IL-6) receptor. We hypothesised that SOCS-3 interferes with FGF signalling, thus influencing the outcome of its action in prostate cancer cells. For this purpose, we treated DU-145 and LNCaP-IL-6+ cells with increasing concentrations of FGF-2, and verified protein phosphorylation. In the presence of FGF-2, neither STAT3, STAT1, nor Akt could be phosphorylated. Solely the p44/p42 MAPK pathway was activated after FGF-2 stimulation. We show for the first time that SOCS-3 interferes with the FGF-2 signalling pathway by modulating p44 and p42 phosphorylation in prostate cancer cells. Decreased SOCS-3 protein expression results in increased MAPK phosphorylation, whereas SOCS-3 overexpression leads to a decreased cellular proliferation and migration. On the basis of the present results, we propose that SOCS-3 is a novel modulator of FGF-2-regulated cellular events in prostate cancer.

Endocrine-Related Cancer (2010) 17 525–538

Introduction

The fibroblast growth factor (FGF) family consists of 23 members, FGF-1–23. FGFs interact with a family of four tyrosine kinase receptors, FGFR1–4. Many FGFs like FGF-2 have potent mitogenic and angiogenic activity, and have been considered as promoters of tumour growth and angiogenesis (Ropiquet *et al.* 1999, 2000). FGF-2 has also been linked with an increase in motility and invasiveness of cancer cells (Wesley *et al.* 2005). Although the growth factor does not contain a signal sequence for secretion, it is secreted and localised in the extracellular matrix of many tissues, and can act in a paracrine or autocrine manner. In addition, FGF-2 also contains a nuclear

localisation sequence (Bugler *et al.* 1991). However, the role of nuclear localisation in the activity of FGF-2 remains unclear.

The expression of FGF-2 is increased in many types of carcinomas including those of prostate (Giri *et al.* 1999). FGF-2 can bind to all four FGFRs with variable binding affinities for each of them (Ornitz *et al.* 1996). The receptors consist of an extracellular part with three immunoglobulin-like domains and an intracellular part with a tyrosine kinase domain. FGFR1–3 can be alternatively spliced, which results in different isoforms with altered binding specificities. FGFs and the FGF signalling pathway appear to play not only an important role in the maintenance and growth of the

normal prostate but also during the development and progression of prostate cancer. The major source of FGFs is the stromal compartment in the normal prostate. A balanced interaction between stromal and epithelial cells through FGFs, which act as paracrine growth factors for epithelial cells, is crucial for normal development and growth of prostate epithelium (Tuxhorn *et al.* 2002, Ayala *et al.* 2003, Yang *et al.* 2008). FGF-2 supports proliferation, migration and invasion in human prostate cancer cellular models (Wesley *et al.* 2005). Oncogenic events caused by FGF-2, including proliferation and migration, could be in part mediated through heparin affinity regulatory peptide/pleiotrophin (Hatziaepostolou *et al.* 2006).

Binding of FGF-2 leads to dimerisation and transphosphorylation of tyrosine residues in the intracellular domain, which is important for receptor kinase activation. In consequence, activation of the receptors can lead to signal transduction through multiple pathways including phospholipase C γ (Mohammadi *et al.* 1991), phosphatidylinositol 3-kinase (PI3K; Hart *et al.* 2001), mitogen-activated protein kinases (MAPKs; Hadari *et al.* 2001), and signal transducers and activators of transcription (STAT; Hart *et al.* 2000, Deo *et al.* 2002, Udayakumar *et al.* 2002). The oncogenic role of STAT3 in various cancers including those of the prostate is well documented (Burke *et al.* 2001, Klein *et al.* 2003, Hodge *et al.* 2005). The activation of STAT3 is associated with either inhibition or promotion of prostate tumour growth (Spitotto & Chung 2000, Giri *et al.* 2001). In previous studies, we showed that STAT3 phosphorylation inversely correlates with the expression of suppressor of cytokine signalling-3 (SOCS-3) in prostate cancer cells, suggesting an inactivation of the Janus kinase (JAK)–STAT pathway in the presence of SOCS-3 (Bellezza *et al.* 2006). The SOCS family consists of eight members, SOCS-1–7 and CIS. All these proteins contain a N-terminal region of variable length, a central SH2 domain, and a conserved C-terminal region called SOCS box. The complex can recruit E2 ubiquitin transferases (Kamura *et al.* 2004). Thus, the protein can act as an E3 ubiquitin ligase, and mediates proteasomal degradation of associated proteins. SOCS-3 is induced especially by interleukin-6 (IL-6), a multifunctional cytokine, which plays an important role in chronic inflammation of the prostate (DeMarzo *et al.* 2003, Wong *et al.* 2008) and prostate carcinogenesis (Hammacher *et al.* 2005). IL-6-mediated JAK and STAT3 activation can be directly inhibited through the kinase inhibitory region or through high-affinity binding of SOCS-3 to the IL-6 receptor subunit gp130 (Nicholson *et al.* 2000).

Thus far, it was shown that SOCS-3 is not only a negative feedback regulator of the IL-6 receptor, but can also interact with tyrosine kinase receptors, such as insulin-like growth factor receptor 1 (IGFR1; Yadav *et al.* 2005), epidermal growth factor receptor (Xia *et al.* 2002) and FGFR (Ben-Zvi *et al.* 2006). It became evident that SOCS proteins inhibit the action of other peptides, such as IGF (Yadav *et al.* 2005). We hypothesised that SOCS-3 interferes with the action of FGF-2 in prostate cancer cells.

In this study, we demonstrate for the first time that SOCS-3 is a key regulator of the FGF pathway. SOCS-3 interferes with FGF-2 signalling by influencing the phosphorylation of the MAPK p44 and p42 in DU-145, LNCaP-IL-6+ and LNCaP prostate cancer cells. By modulating SOCS-3 expression levels, we could clearly demonstrate that SOCS-3 has an inhibitory effect on the activation of the MAPK signal transduction. Overexpression of SOCS-3 leads to decreased p44 and p42 activation and in consequence to a diminished cell proliferation and migration. Thus, in the present study, we have shown that SOCS-3 interference with FGF-2 signalling is relevant to prostate cancer progression.

Materials and methods

Cell culture and chemicals

LNCaP and DU-145 prostate cancer cells were obtained from American Type Culture Collection (Rockville, MD, USA). The LNCaP subline LNCaP-IL-6+ was derived in the presence of IL-6, as described elsewhere (Hobisch *et al.* 2001). LNCaP cells, stably expressing cmvSOCS-3, were cultured and used as already described in detail in a previous study (Neuwirt *et al.* 2007). All cell lines were cultured in RPMI 1640 (PAA Laboratories, Pasching, Austria). Media were supplemented with 10% FCS and 1% penicillin/streptomycin and glutamine. FGF-2 was a product of Cell Signaling (Danvers, MA, USA). Anisomycin, which was used to induce p38 phosphorylation, was purchased from Sigma.

Short interfering RNA transfection

The short interfering RNA (siRNA) sequence used for targeting human SOCS-3 was published previously (Komyod *et al.* 2007) as follows: sense, 5-CCAAGA ACCUGCGCAUCCAdTdT-3; antisense, 5-UGGAU GCGCAGGUUCUUGGdTdT-3 and synthesised by Genxpress (Vienna, Austria). A non-targeting siRNA pool (Cat. no.: D-001810-10-20, Dharmacon (Chicago, IL, USA)) was used as a negative control.

Transfections with siRNA were performed with Lipofectamine 2000 (Invitrogen) according to the manufacturer's protocol. DU-145 and LNCaP-IL-6+ cells were seeded at a density of 2.5×10^5 /well onto six-well plates. On the next day, the cells were transfected with either 100 nM siRNA against SOCS-3 or control siRNA for a period of 3 days, and harvested and lysed for western blot analysis.

RNA isolation and RT-PCR

Cells were grown on six-well plates. Subsequently, total RNA was isolated using the RNeasy mini kit (Qiagen) following the manufacturer's protocol. cDNA synthesis was performed using iScript select cDNA synthesis kit (Bio-Rad). Real-time PCR conditions were as follows: one cycle of denaturing at 95 °C for 10 min followed by 40 cycles of 95 °C for 15 s and 60 °C for 1 min. All amplifications were performed in triplicates. TATA-box-binding protein (TBP) was chosen as an endogenous expression standard (Savinainen *et al.* 2002). Primer and probe sequences were as follows: SOCS-3 (forward 5'-TGATCCGC-GACAGCTCG-3'; reverse 5'-TCCCAGACTGGGT-CTTGACG-3'; TaqMan probe 5'-FAM-CCAGCGC-CACTTCTTCACGCTCA-TAMRA-3'), TBP (forward 5'-CACGAACCACGGCACTGATT-3'; reverse 5'-TTTTCTGCTGCCAGTCTGGAC-3'; TaqMan probe 5'-FAM-TCTTCACTCTTGGCTCCTGTG-CACA-TAMRA-3'). Each reaction was performed in a 11- μ l volume containing 50 ng cDNA and 5.5 μ l of 2 $\times$ ABI Mastermix (Applied Biosystems, Foster City, CA, USA). The primers were added to a final concentration of 900 nmol/l, and the final probe concentration was 150 nmol/l. PCR products were measured using the ABI Prism 7500 Fast RT-PCR System (Applied Biosystems). C_t values of SOCS-3 and TBP as assessed by ABI Sequence Detection Software (version 1.3) were used to calculate the ΔC_t using Microsoft Excel. Values obtained in control cells were defined as 100%, and those from treated cells were expressed as a percentage of control.

Western blot

Cells were harvested, washed with PBS, and lysed in EORTC buffer. The total protein was quantified using the Bradford method (Bradford 1976). Fifty micrograms of protein per lane were then resolved using a 4–12% Bis–Tris gel (Invitrogen), and transferred onto a nitrocellulose membrane (Invitrogen). Western blot was carried out as described elsewhere (Puhr *et al.* 2009). The following antibodies were used for western blots: SOCS-3 (1:1000; Acris Antibodies,

Hiddenhausen, Germany), GAPDH (1:100 000; Millipore, Vienna, Austria), phospho-STAT3 (1:200; Upstate Cell Signaling Technology, Beverly, MA, USA), STAT3 (1:1000; Santa Cruz Biotechnology, Santa Cruz, CA, USA), phospho-STAT1 (1:500; Santa Cruz), STAT1 (1:1000; Santa Cruz), Akt (1:1000; Cell Signaling), pAkt (1:500; Cell Signaling), pp44/42 (1:500; Cell Signaling), p44 (1:1000; Santa Cruz), p42 (1:1000; Santa Cruz), pp38 (1:250; New England Biolabs, Ipswich, MA, USA) and p38 (1:500, New England Biolabs).

[³H]thymidine incorporation assay

DU-145 and LNCaP-IL-6+ cells were seeded at a density of 3.5×10^3 /well, and cmcSOCS-3 stably transfected LNCaP cells were seeded at a density of 1.5×10^4 /well in triplicates onto 96-well plates. On the following day, full medium was changed to starving medium and FGF-2 for DU-145 and LNCaP-IL-6+ cells, or FGF-2 and doxycycline for LNCaP cells. The cells were incubated for 5 days. At day 3, the medium was changed, and FGF-2 and doxycycline were again added to the cells. To each well, 25 μ l of [³H]thymidine (1 μ Ci/well) were added, and DNA was harvested on day 5 on 96-well filter plates (UniFilter, Perkin Elmer, Boston, MA, USA) after a 24 h incubation. Fifty microlitres of scintillation fluid were added, and radioactivity was quantified using Hidex Chameleon V liquid scintillation counter (HVD Life Sciences, Vienna, Austria).

Wound-filling assay

DU-145 and LNCaP-IL-6+ cells were seeded at a density of 3.5×10^5 /well in six-well plates and transiently transfected with a doxycycline-inducible cmcSOCS-3 plasmid. The transfection was performed with FuGENE 6 (Roche Diagnostics) according to the manufacturer's protocol. Identical scratches were made in parallel wells 24 h after the transfection with a 1000- μ l tip, and doxycycline and FGF-2 were added respectively. Pictures were taken at 0, 24 and 48 h.

Statistical analysis

For each treatment group, statistical significance was determined by Mann–Whitney *U* test. *P* values < 0.05 were defined as statistically significant and encoded in figures (**P* < 0.05 and ***P* < 0.01). Statistical analyses were performed using SPSS 12.0 software (SPSS, Chicago, IL, USA).

Results

Activation of the p44/p42 signalling pathway by FGF-2 in prostate cancer cells

Phosphorylation of ligand-bound FGFRs may lead to the activation of multiple downstream signal transduction pathways, such as those of STAT, MAPK and Akt. Because FGF-2 signalling has not been characterised in prostate cancer so far, we analysed the activation of different signalling pathways after FGF-2 stimulation in prostate cancer cells. For this purpose, we treated DU-145 and LNCaP-IL-6+ cells that represent advanced carcinoma of the prostate with increasing concentrations of FGF-2 (1–100 ng/ml) for 15 min, and verified protein phosphorylation by western blot afterwards. LNCaP cells, treated with 10 ng/ml IL-6 for 15 min, were used as a positive control. In the presence of FGF-2, neither STAT3, STAT1, p38 MAPK, nor Akt could be phosphorylated, even with the highest growth factor concentration in DU-145 cells (Fig. 1a, [Supplementary Figure 1](#), see section on [supplementary data](#) given at the end of this article). In LNCaP-IL-6+ cells, no STAT3 and Akt phosphorylation was detectable, whereas a minor STAT1 phosphorylation (Fig. 1b) could be observed. However, there was no difference in phosphorylation intensity between control and FGF-2-treated cells.

We also investigated a possible activation of the p44/p42 MAPK pathway. Therefore, we measured p44/p42 phosphorylation in both cell lines after 15 and 45 min of FGF-2 treatment. Western blot data showed a significant dose-dependent increase of p44/p42 phosphorylation in DU-145 and LNCaP-IL-6+ cells at both time points (Fig. 2).

FGF-2 differentially regulates SOCS-3 in two prostate cancer cell lines

In several cellular models, IL-6 was identified as a classical inducer of SOCS-3, which in turn limits the activation of the JAK–STAT pathway. To study the possible effects of FGF-2 on SOCS-3 expression, SOCS-3 protein levels were measured in both cell lines. We observed a moderate increase of SOCS-3 expression by FGF-2 in DU-145 cells (Fig. 2a) within 15 min. In LNCaP-IL-6+ cells, we did not observe changes in SOCS-3 expression after treatment with 1 ng/ml FGF-2. However, a decrease of SOCS-3 expression with 100 ng/ml of FGF-2 was noted (Fig. 2b). SOCS-3 mRNA levels were not altered after FGF-2 short-time treatment (data not shown).

After long-term treatments with FGF-2 (up to 5 days), we could confirm these findings (Fig. 3).

Figure 1 FGF-2 stimulation does not activate STAT3, STAT1 or Akt in prostate cancer cells. Western blot analysis for STAT3, STAT1, and Akt signal transduction pathway activation after treatment with increasing concentrations of FGF-2 for 15 min in DU-145 (a) and LNCaP-IL-6+ cells (b). LNCaP cells, treated with 10 ng/ml IL-6 for 15 min, were used as a control for the activation of the STAT3 and Akt pathways.

Figure 2 Specific activation of the MAPK pathway and regulation of SOCS-3 after FGF-2 stimulation. Western blot analysis for MAPK activation and SOCS-3 expression in DU-145 (a) and LNCaP-IL-6+ cells (b). Treatment with FGF-2 for 15 and 45 min leads to a dose-dependent phosphorylation of p44 and p42, key proteins of the MAPK pathway (mean values + S.E.M. from at least three independent experiments). **, $P < 0.01$.

No significant change of SOCS-3 mRNA levels could be observed in both cell lines after 5 days (Fig. 3a and c). At protein level, we observed a significant increase of SOCS-3 expression with 1 ng/ml of FGF-2 and a dose-dependent decline with increasing concentrations of FGF-2 in DU-145 cells (Fig. 3b). In LNCaP-IL-6+ cells, similar to the short-time treatment, a significant decrease of SOCS-3 expression could be detected with 10 and 100 ng/ml FGF-2 (Fig. 3d).

FGF-2-induced p44/p42 phosphorylation is required for prostate cancer cells' proliferation

In order to investigate the consequences of increased p44/p42 phosphorylation, DU-145 and LNCaP-IL-6+ cells were incubated with 10 and 100 ng/ml FGF-2 for 5 days (Fig. 4a and b). Cellular proliferation was measured using [3 H]thymidine incorporation. To clarify whether altered cellular proliferation is a specific consequence of p44/p42 activation, we incubated both cell lines in the presence of FGF-2 and the specific MAPK inhibitor PD98059.

We observed a significant increase in proliferation caused by 10 and 100 ng/ml FGF-2 in both cell lines after 5 days, whereas in the presence of 25 μ M PD98059 no significant alteration could be detected. Concentrations higher than 25 μ M of PD98059 inhibited the proliferation of both FGF-treated and -untreated prostate cancer cell lines (data not shown), since it is known that DU-145 and LNCaP-IL-6+ cells show up-regulated p44/p42 activity (Putz *et al.* 1999, Steiner *et al.* 2003). The data presented in Fig. 4 illustrate that p44/p42 phosphorylation is required for proliferation of DU-145 and LNCaP-IL-6+ cells.

SOCS-3 downregulation results in increased p44/p42 phosphorylation

To determine whether SOCS-3 expression has an impact on p44/p42 phosphorylation, we transfected DU-145 and LNCaP-IL-6+ cells with 100 nM control siRNA or siRNA against SOCS-3 as indicated and measured SOCS-3 protein expression levels and p44/p42 phosphorylation after 3 days (Fig. 5a and c).

Figure 3 SOCS-3 expression is regulated at protein but not at mRNA level after FGF-2 treatment. DU-145 (a and b) and LNCaP-IL-6+ cells (c and d) were treated two times with 1, 10 and 100 ng/ml FGF-2 during a period of 5 days. The cells were then harvested, and qRT-PCR and western blot were performed respectively. The results are expressed in relation to values measured in untreated cells and represent mean values $\pm$ s.e.m. from at least three independent experiments. **, $P < 0.01$.

We observed a significant 50–55% decrease of SOCS-3 protein in both cell lines. A larger decrease in SOCS-3 expression yields massive cell death which could compromise proliferation and migration experiments (Puhr et al. 2009). At the same time, a 2.5-fold increase in p44/p42 phosphorylation could be measured in DU-145 cells. In LNCaP-IL-6+ cells, an increase in phosphorylation could also be detected. Nevertheless, it was not significant compared with the negative control.

Next, we incubated both cell lines with 1 ng/ml FGF-2 for 15 and 45 min after siRNA treatment, and measured p44/p42 phosphorylation again. A significant 3.5-fold increase in p44/p42 phosphorylation was observed compared to the treated negative control, when SOCS-3 protein expression was significantly decreased in DU-145 cells (Fig. 5b). In the LNCaP-IL-6+ cell line, a similar twofold increase in p44/p42 phosphorylation was measured (Fig. 5d).

SOCS-3 overexpression yields decreased p44/p42 phosphorylation

To examine whether SOCS-3 regulation of p44/p42 phosphorylation is a general phenomenon, we stably transfected SOCS-3-negative LNCaP cells with a

doxycycline-inducible mycSOCS-3 plasmid (Neuwirt et al. 2007). After induction of mycSOCS-3 by doxycycline, the cells were incubated with 10 ng/ml IL-6 (Fig. 6a) or FGF-2 (Fig. 6b) for 15 and 45 min, and p44/p42 phosphorylation was measured afterwards. In the presence of increased SOCS-3 protein levels, a significant decrease in p44/p42 phosphorylation could be detected, compared to the doxycycline-untreated control after IL-6 or FGF-2 stimulation for both time points.

Overexpression of SOCS-3 diminished FGF-2-induced proliferation and migration

In our previous studies, it was demonstrated that SOCS-3 may exhibit different effects on proliferation and apoptosis in androgen-sensitive and -insensitive cells. Decreased p44/p42 phosphorylation, a consequence of SOCS-3 overexpression, has a significant impact on cell proliferation (Fig. 7). Proliferation of stably transfected LNCaP cells was measured after 5 days of stimulation by FGF-2 in the absence or in the presence of doxycycline, as indicated, using [3 H]thymidine incorporation assay. When SOCS-3 was not present in LNCaP cells, a dose-dependent

Figure 4 Increased p44/p42 phosphorylation is crucial for prostate cancer cell proliferation. Regulation of cellular proliferation in DU-145 (a) and LNCaP-IL-6+ (b) cells. Consecutive treatment with FGF-2 during a period of 5 days causes an increase of proliferation in both cell lines. This effect could be reversed in the presence of the MAPK inhibitor PD98059. Cellular proliferation was assessed by the measurement of [³H]thymidine incorporation (mean values $\pm$ s.e.m. from at least three independent experiments). The percentage of untreated cells was set at 100%. *, $P < 0.05$; **, $P < 0.01$.

significant increase in proliferation could be measured with 10 and 100 ng/ml FGF-2 (Fig. 7a). On the contrary, in the presence of SOCS-3, no increase in proliferation could be observed (Fig. 7b).

To get a better insight whether increased SOCS-3 expression has also an effect on migration of DU-145 and LNCaP-IL-6+ cells, we performed a wound-filling assay 3 days after transient transfection with the doxycycline-inducible cmvSOCS-3 plasmid (Fig. 8). Twenty-four hours after adding 10 ng/ml FGF-2, an obvious difference between the treatment groups could be observed in both cell lines. Thus, induction of SOCS-3 by doxycycline resulted in a reduced

migration of prostate cancer cells. After 48 h, this difference between the cells treated with FGF-2 and those treated with FGF-2 and doxycycline was even more evident.

Discussion

FGF-2 endogenous inhibitors are differently expressed in prostate cancer

To the best of our knowledge, prior to this study, there was no published evidence that SOCS-3 interacts with elements of FGF pathways in cancer or represses MAPK activation. To get more insight into the regulation of FGF-2-mediated cellular events by SOCS-3, we investigated a possible interference of SOCS-3 with FGF-2 signal transduction and examined its biological consequences in prostate cancer *in vitro*. For this purpose, we used DU-145, LNCaP and LNCaP-IL-6+ cells, which are established models for advanced prostate cancer. FGFs and their receptors influence different stages of prostate carcinogenesis (Leung *et al.* 1996, Dorkin *et al.* 1999). Overexpression of multiple FGFs, especially of FGF-2, and an altered expression of FGFRs have been reported in prostate cancer (Giri *et al.* 1999). Indeed, FGF-2 has mitogenic and angiogenic characteristics, and can promote tumour progression (Powers *et al.* 2000, Polnaszek *et al.* 2003). Therefore, it is important to identify endogenous inhibitors of FGF signalling in carcinoma of the prostate.

FGF-2 overexpression and, in consequence, altered signal transduction are also linked to a significant decrease in some of the regulators of the FGF pathway. Kwabi-Addo *et al.* (2004) could show that a member of the Sprouty family, Sprouty 1, an important negative feedback inhibitor of FGF-2, is significantly decreased in prostate cancer. Recently, it was reported that Sprouty 4, another member of this family which is involved in the regulation of cellular migration, is silenced through hypermethylation (Wang *et al.* 2006). Dipeptidyl peptidase, a serine protease which regulates the activities of several mitogens, is another example for an endogenous FGF-2 inhibitor, whose expression is decreased in prostate cancer (Wesley *et al.* 2005). We identified in the present study SOCS-3 as a novel inhibitor of FGF-2 signalling. In contrast to the Sprouty proteins or dipeptidyl peptidase, SOCS-3 is expressed in prostate cancer cells and tissues obtained from patients with prostate intraepithelial neoplasia or cancer including those who failed endocrine therapy (Puhr *et al.* 2009). For SOCS-3 inhibitory function of FGF-2 function *in vivo*, the expression of growth factor inhibitors belonging to the Sprouty family or dipeptidyl peptidase may be required.

Figure 5 Measurement of p44/p42 phosphorylation after SOCS-3 downregulation and FGF-2 stimulation. Western blot analysis for basal p44 and p42 phosphorylation in DU-145 (a) and LNCaP-IL-6+ (c) cells after SOCS-3 downregulation for 3 days. Cells were treated with specific SOCS-3 or negative control siRNA. Treatment with 1 ng/ml FGF-2 for 15 and 45 min in DU-145 (b) and LNCaP-IL-6+ cells (d) resulted in an increased p44/p42 phosphorylation. The results are expressed in relation to values measured in cells treated with negative control siRNA and represent mean values $\pm$ S.E.M. from at least three independent experiments. **, $P < 0.01$.

SOCS-3 is a regulator of receptor tyrosine kinases

There is accumulated evidence that the inhibitory role of SOCS-3 is not limited to IL-6 signalling. Furthermore, SOCS-3 can bind to tyrosine kinase receptors and interfere with their signal transduction. Ben-Zvi *et al.* (2006) could show that SOCS-3 and SOCS-1 bind to FGFR3 in HEK-293T cells. In those cells, SOCS-1 overexpression elevated MAPK phosphorylation, in contrast to the effect of SOCS-3 in prostate cancer, and repressed STAT1 phosphorylation. An interaction between the IGF1 pathway and SOCS-3 has been analysed. SOCS-3 is induced in response to IGF1 in neurons, and antagonises its effect on survival through

binding to the IGF1R (Yadav *et al.* 2005). IGF1 also upregulates SOCS-3 during differentiation of myoblasts, thus inducing transcriptional activation of the α -actin promoter (Spangenberg 2005).

In prostate cancer cells, FGF-2 acts through MAPK rather than through the JAK/STAT pathway. SOCS proteins were originally identified as main inhibitors of the STAT pathway. This effect of SOCS-3 on growth factor signalling is not limited to a single cell line. It was observed in DU-145 and LNCaP-IL-6+ cells, which express endogenous SOCS-3, and in LNCaP clones, in which SOCS-3 expression was induced after doxycycline treatment. On the basis of these results and our previous work, we conclude that SOCS-3 in human

Figure 6 Overexpression of SOCS-3 results in a decreased p44/p42 phosphorylation. Western blot analysis for p44/p42 phosphorylation in SOCS-3-overexpressing LNCaP cells. The presence of SOCS-3 leads to a significantly decreased p44/p42 phosphorylation after 15 and 45 min after IL-6 (a) and FGF-2 (b) treatment in cmvSOCS-3-transfected LNCaP cells. **, $P < 0.01$.

prostate cancer has multiple functions. Its expression in prostate cells can be induced by IL-6, cAMP and androgens (Bellezza *et al.* 2006, Neuwirt *et al.* 2007). Importantly, the implications of SOCS-3 induction by each of these compounds may be contrasting. cAMP-induced SOCS-3 expression yields increased proliferation and reduced cell death (Bellezza *et al.* 2006). In androgen-sensitive cells, SOCS-3 antagonises hormonal effect on proliferation by interfering with key regulators of cell cycle progression (Neuwirt *et al.* 2007). Further studies to assess the implications of inhibition of SOCS-3 expression in prostate cancer *in vivo* may be therefore justified.

FGF-2 activation of intracellular signalling is cell type specific

The treatment of DU-145 and LNCaP-IL-6+ cells with increasing concentrations of FGF-2 did not result in phosphorylation of STAT3 and STAT1. These findings

were consistent with our previous results, which showed that STAT3 and STAT1 phosphorylation in prostate cancer inversely correlates with SOCS-3 expression *in vitro* (Bellezza *et al.* 2006, Puhr *et al.* 2009). In SOCS-3-positive cell lines, STAT3 cannot be phosphorylated. Treatment of prostate cancer cells with SOCS-3 siRNA led to increased phosphorylation of STAT factors. The results showing a lack of effect of FGF-2 on STAT phosphorylation are in line with those of Ishikawa *et al.* (2005). They could not detect phosphorylation of STAT3 and STAT1 after FGF-2 stimulation in a myeloma cell line. On the other hand, Deo *et al.* (2002) reported STAT3 phosphorylation after FGF-2 stimulation in human umbilical vein endothelial cells. Thus, the phosphorylation of STATs after FGF-2 stimulation depends on cellular context. As we were not able to detect the activation of the PI3K pathway after FGF-2 treatment, we focused on MAPK in the present study. Indeed, the treatment with increasing concentrations of FGF-2

Figure 7 Cellular proliferation is decreased in the presence of SOCS-3. Regulation of cellular proliferation in mycSOCS-3-transfected LNCaP cells in the presence or in the absence of doxycycline. Treatment with 10 and 100 ng/ml FGF-2 during a period of 5 days causes a significant increase of proliferation. This effect could be reversed in the presence of 2 µg/ml doxycycline. Cellular proliferation was assessed by the measurement of [³H]thymidine incorporation (mean values + S.E.M. from at least three independent experiments). The percentage of untreated cells was set at 100%. *, $P < 0.05$.

resulted in a dose-dependent phosphorylation of the MAPK p44 and p42. These data are again in accordance with those of Ishikawa *et al.* (2005) as well as with those of two other groups who showed that the FGF-2 effect is mediated through this pathway in prostate cancer cellular models (Wesley *et al.* 2005, Hatziaepostolou *et al.* 2006).

SOCS-3 inhibition of p44/p42 MAPK antagonises FGF-2-induced proliferation and migration

MAPKs have an important role in cellular proliferation, and its phosphorylation is increased in a number of prostate cancer patient cases (Gioeli *et al.* 1999). To determine whether FGF-2-increased activation of p44 and p42 has an effect on proliferation of DU-145 and LNCaP-IL-6+ cells, they were treated with FGF-2

in the presence or in the absence of the specific MAPK inhibitor PD98059. Our results confirm that increased p44/p42 phosphorylation is required for the FGF-2-induced cellular proliferation in prostate cancer. These findings are supported by those of Giri *et al.* (1999) who showed an increased proliferation of primary prostate epithelial cells after FGF-2 stimulation. Recombinant FGF-2 also stimulated DNA synthesis in DU-145 prostate cancer cells in a previous study. However, the relative effect was lower than that observed in LNCaP cells which do not produce endogenous FGF-2 (Nakamoto *et al.* 1992). It should be noted that in prostate cancer cell lines used in the present study, FGFs may exhibit autocrine activity. LNCaP-IL-6+ cells, in contrast to the parental cells, upregulate IL-6 and SOCS-3 (Hobisch *et al.* 2001, Bellezza *et al.* 2006). This observation may provide an explanation why addition of exogenous FGF-2 does not elevate SOCS-3 protein expression in the LNCaP-IL-6+ subline.

Importantly, we show that SOCS-3 interference with the p44/p42 MAPK pathway has consequences on proliferation and migration of prostate cancer cells. In LNCaP cells, overexpression of SOCS-3 inhibits FGF-2-induced proliferation. DU-145 and LNCaP-IL-6+ cells show a decreased migrative potential in the presence of elevated SOCS-3 protein levels. We could thus extend the results of a previous work, in which it was revealed that FGF-2 increases the expression and secretion of heparin affini regulatory peptide/pleiotrophin, cellular proliferation and migration (Hatziaepostolou *et al.* 2006). One could hypothesise that expression of FGF-2 targets such as cyclin E, retinoblastoma or matrix metalloproteinases is affected through the effect of SOCS-3 on p44/p42 MAPK phosphorylation (Muddasani *et al.* 2007, Takeuchi *et al.* 2007). We demonstrate for the first time that SOCS-3 may interfere with oncogenic events caused by the p44/p42 MAPK pathway in cancer. Our findings extend previous knowledge since SOCS proteins were identified as classic inhibitors of the JAK/STAT signal transduction. It remains to be determined which elements of the FGF signalling pathway, such as receptors, interact with SOCS-3. Possible binding partners of SOCS-3 could be members of the MAPK family or the FGFR. FGFR1 is considered to be the most important receptor of the FGFR family for FGF-2 signalling and a known oncoprotein in prostate cancer (Freeman *et al.* 2003). However, we could not detect an interaction between SOCS-3 and FGFR1 in DU-145 or LNCaP-IL-6+ cells (data not shown). FGFR2, FGFR3, and FGFR4 are also involved in transduction of FGF-2 signalling in malignancies. However, in

Figure 8 SOCS-3 overexpression neutralises the migratory effect of FGF-2 in a wound-filling assay. SOCS-3 expression was induced with 2 μ g/ml doxycycline as indicated. Identical scratches were made in parallel wells using a 1000- μ l pipette tip 48 h after transfection. Pictures (40 $\times$) were taken at 0, 24, and 48 h. The strongest differences between the FGF-2 treatment and the FGF-2 + doxycycline treatment were observed at 48 h for DU-145 cells (a) and LNCaP-IL-6+ cells (b).

contrast to FGFR1 which is oncogenic, re-expression of FGFR2 in human prostate cancer cells led to the inhibition of growth and tumorigenicity (Yasumoto *et al.* 2004, Acevedo *et al.* 2007). The role of FGFR3 and FGFR4 in human prostate cancer is at present less clear, and the implications of receptor mutations should be further investigated (Wang *et al.* 2004, Hernandez *et al.* 2009). We hypothesise that the action of SOCS-3 on the FGF-2 signalling pathway is more complex than its interference with IGF1 signalling. In the latter case, it was documented that SOCS-3 interacts with the IGFRI (Dey *et al.* 2000).

Two different approaches were used to examine the regulation of MAPK by SOCS-3 in prostate cancer: i) specific siRNA against SOCS-3 and ii) doxycycline-regulated SOCS-3 expression to manipulate SOCS-3 protein expression in prostate cancer cell lines. Downregulation of SOCS-3 resulted in a significant increase of basal p44/p42 phosphorylation in DU-145 cells but not in LNCaP-IL-6+ cells. The basal p44/p42 phosphorylation in LNCaP-IL-6+ cells increased as a consequence of chronic IL-6 treatment as shown by our group previously (Steiner *et al.* 2003, Cavarretta *et al.* 2007). There may be several explanations for this apparent discrepancy. One should keep in mind that other cytokines compensate for loss of FGF-2 expression. Nevertheless, after treatment with FGF-2, both cell lines responded with a significant increase in p44/p42 phosphorylation compared to the treated control in the presence of decreased SOCS-3 protein levels. On the other hand, LNCaP cells overexpressing SOCS-3 showed a decreased p44/p42 phosphorylation after FGF-2 treatment.

The findings of our study may have considerable clinical implications. Both FGF-2 and SOCS-3 were

detected in prostate cancer, and it was shown in experiments using cell lines and patients' material that their expression increases during progression to therapy resistance (Nakamoto *et al.* 1992, Cronauer *et al.* 1997, Bellezza *et al.* 2006, Puhr *et al.* 2009).

SOCS-3 mRNA and protein are differentially regulated by FGF-2

FGF-2 regulated SOCS-3 protein expression levels, although mRNA level remained unchanged. One plausible explanation for this finding could be cell line-specific protein stabilisation or increased proteasomal degradation as a consequence of FGF-2 treatment. In order to elucidate the importance of this observation, we extended our treatment time up to 5 days. Consistently, the data were similar and confirmed short-time treatment results. A transient downregulation of SOCS-3 mRNA by IGF1 was also reported (Evans *et al.* 2007). Thus, peptide growth factors may display different effects on SOCS mRNA and protein expression respectively.

In summary, in the present study, we clearly showed that SOCS-3, which is in part regulated by FGF-2, antagonises the effect of the growth factor by interfering with the activation of the p44/p42 MAPK pathway. SOCS-3 is a multifunctional key protein in many pathways, and there are indications that it has different functions depending on the tumour origin. In a previous study, we showed that sustained downregulation of SOCS-3 leads to increased apoptosis through the activation of the extrinsic and intrinsic apoptosis pathways in prostate cancer cells (Puhr *et al.* 2009). We also demonstrated that SOCS-3 is highly expressed in castration-resistant prostate cancer patients (Bellezza *et al.* 2006, Puhr *et al.* 2009).

Taken together, due to the involvement of SOCS-3 in many different signalling pathways, it can act as an oncogene or a tumour suppressor, depending on the cellular context. In future studies, the interaction of SOCS-3 with other pathways must be investigated in detail to better understand the role of this pleiotropic protein in cancer development and progression.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1677/ERC-10-0007>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

Austrian Science Fund FWF – grants W1101, 19933, and 18193 (to Z Culig).

Acknowledgements

The authors acknowledge Dr Ilaria T Cavarretta for helpful discussions, Gertraud Sierek for excellent technical assistance and Robert Schober for the editorial assistance. The authors thank Dr Wolfgang Doppler for kindly providing the cmycSOCS-3 construct used in the present study.

References

Acevedo VD, Gangula RD, Freeman KW, Li R, Zhang Y, Wang F, Ayala GE, Peterson LE, Ittmann M & Spencer DM 2007 Inducible FGFR-1 activation leads to irreversible prostate adenocarcinoma and an epithelial-to-mesenchymal transition. *Cancer Cell* **12** 559–571.

Ayala G, Tuxhorn JA, Wheeler TM, Frolov A, Scardino PT, Ohori M, Wheeler M, Spittler J & Rowley DR 2003 Reactive stroma as a predictor of biochemical-free recurrence in prostate cancer. *Clinical Cancer Research* **9** 4792–4801.

Bellezza I, Neuwirt H, Nemes C, Cavarretta IT, Puhr M, Steiner H, Minelli A, Bartsch G, Offner F, Hobisch A *et al.* 2006 Suppressor of cytokine signaling-3 antagonizes cAMP effects on proliferation and apoptosis and is expressed in human prostate cancer. *American Journal of Pathology* **169** 2199–2208.

Ben-Zvi T, Yayon A, Gertler A & Monsonego-Ornan E 2006 Suppressors of cytokine signaling (SOCS)1 and SOCS3 interact with and modulate fibroblast growth factor receptor signaling. *Journal of Cell Science* **119** 380–387.

Bradford MM 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72** 248–254.

Bugler B, Amalric F & Prats H 1991 Alternative initiation of translation determines cytoplasmic or nuclear localization of basic fibroblast growth factor. *Molecular and Cellular Biology* **11** 573–577.

Burke WM, Jin X, Lin HJ, Huang M, Liu R, Reynolds RK & Lin J 2001 Inhibition of constitutively active Stat3 suppresses growth of human ovarian and breast cancer cells. *Oncogene* **20** 7925–7934.

Cavarretta IT, Neuwirt H, Untergasser G, Moser PL, Zaki MH, Steiner H, Rumpold H, Fuchs D, Hobisch A, Nemeth JA *et al.* 2007 The antiapoptotic effect of IL-6 autocrine loop in a cellular model of advanced prostate cancer is mediated by Mcl-1. *Oncogene* **26** 2822–2832.

Cronauer MV, Hittmair A, Eder IE, Hobisch A, Culig Z, Ramoner R, Zhang J, Bartsch G, Reissigl A, Radmayr C *et al.* 1997 Basic fibroblast growth factor levels in cancer cells and in sera of patients suffering from proliferative disorders of the prostate. *Prostate* **31** 223–233.

DeMarzo AM, Nelson WG, Isaacs WB & Epstein JI 2003 Pathological and molecular aspects of prostate cancer. *Lancet* **361** 955–964.

Deo DD, Axelrad TW, Robert EG, Marcheselli V, Bazan NG & Hunt JD 2002 Phosphorylation of STAT-3 in response to basic fibroblast growth factor occurs through a mechanism involving platelet-activating factor, JAK-2, and Src in human umbilical vein endothelial cells. Evidence for a dual kinase mechanism. *Journal of Biological Chemistry* **277** 21237–21245.

Dey BR, Furlanetto RW & Nissley P 2000 Suppressor of cytokine signaling (SOCS)-3 protein interacts with the insulin-like growth factor-I receptor. *Biochemical and Biophysical Research Communications* **278** 38–43.

Dorkin TJ, Robinson MC, Marsh C, Neal DE & Leung HY 1999 aFGF immunoreactivity in prostate cancer and its co-localization with bFGF and FGF8. *Journal of Pathology* **189** 564–569.

Evans MK, Yu CR, Lohani A, Mahdi RM, Liu X, Trzeciak AR & Egwuagu CE 2007 Expression of SOCS1 and SOCS3 genes is differentially regulated in breast cancer cells in response to proinflammatory cytokine and growth factor signals. *Oncogene* **26** 1941–1948.

Freeman KW, Gangula RD, Welm BE, Ozen M, Foster BA, Rosen JM, Ittmann M, Greenberg NM & Spencer DM 2003 Conditional activation of fibroblast growth factor receptor (FGFR)1, but not FGFR2, in prostate cancer cells leads to increased osteopontin induction, extracellular signal-regulated kinase activation, and *in vivo* proliferation. *Cancer Research* **63** 6237–6243.

Gioeli D, Mandell JW, Petroni GR, Frierson HF Jr & Weber MJ 1999 Activation of mitogen-activated protein kinase associated with prostate cancer progression. *Cancer Research* **59** 279–284.

- Giri D, Ropiquet F & Ittmann M 1999 Alterations in expression of basic fibroblast growth factor (FGF) 2 and its receptor FGFR-1 in human prostate cancer. *Clinical Cancer Research* **5** 1063–1071.
- Giri D, Ozen M & Ittmann M 2001 Interleukin-6 is an autocrine growth factor in human prostate cancer. *American Journal of Pathology* **159** 2159–2165.
- Hadari YR, Gotoh N, Kouhara H, Lax I & Schlessinger J 2001 Critical role for the docking-protein FRS2 alpha in FGF receptor-mediated signal transduction pathways. *PNAS* **98** 8578–8583.
- Hammacher A, Thompson EW & Williams ED 2005 Interleukin-6 is a potent inducer of S100P, which is up-regulated in androgen-refractory and metastatic prostate cancer. *International Journal of Biochemistry & Cell Biology* **37** 442–450.
- Hart KC, Robertson SC, Kanemitsu MY, Meyer AN, Tynan JA & Donoghue DJ 2000 Transformation and Stat activation by derivatives of FGFR1, FGFR3, and FGFR4. *Oncogene* **19** 3309–3320.
- Hart KC, Robertson SC & Donoghue DJ 2001 Identification of tyrosine residues in constitutively activated fibroblast growth factor receptor 3 involved in mitogenesis, Stat activation, and phosphatidylinositol 3-kinase activation. *Molecular Biology of the Cell* **12** 931–942.
- Hatzia Apostolou M, Polytaichou C, Katsoris P, Courty J & Papadimitriou E 2006 Heparin affinity regulatory peptide/pleiotrophin mediates fibroblast growth factor 2 stimulatory effects on human prostate cancer cells. *Journal of Biological Chemistry* **281** 32217–32226.
- Hernandez S, de Muga S, Agell L, Juanpere N, Esgueva R, Lorente JA, Mojal S, Serrano S & Lloreta J 2009 FGFR3 mutations in prostate cancer: association with low-grade tumors. *Modern Pathology* **22** 848–856.
- Hobisch A, Ramoner R, Fuchs D, Godoy-Tundidor S, Bartsch G, Klocker H & Culig Z 2001 Prostate cancer cells (LNCaP) generated after long-term interleukin 6 (IL-6) treatment express IL-6 and acquire an IL-6 partially resistant phenotype. *Clinical Cancer Research* **7** 2941–2948.
- Hodge DR, Hurt EM & Farrar WL 2005 The role of IL-6 and STAT3 in inflammation and cancer. *European Journal of Cancer* **41** 2502–2512.
- Ishikawa H, Tsuyama N, Liu S, Abroun S, Li FJ, Otsuyama K, Zheng X, Ma Z, Maki Y, Iqbal MS *et al.* 2005 Accelerated proliferation of myeloma cells by interleukin-6 cooperating with fibroblast growth factor receptor 3-mediated signals. *Oncogene* **24** 6328–6332.
- Kamura T, Maenaka K, Kotoshiba S, Matsumoto M, Kohda D, Conaway RC, Conaway JW & Nakayama KI 2004 VHL-box and SOCS-box domains determine binding specificity for Cul2-Rbx1 and Cul5-Rbx2 modules of ubiquitin ligases. *Genes and Development* **18** 3055–3065.
- Klein B, Tarte K, Jourdan M, Mathouk K, Moreaux J, Jourdan E, Legouffe E, De Vos J & Rossi JF 2003 Survival and proliferation factors of normal and malignant plasma cells. *International Journal of Hematology* **78** 106–113.
- Komyod W, Bohm M, Metz D, Heinrich PC & Behrmann I 2007 Constitutive suppressor of cytokine signaling 3 expression confers a growth advantage to a human melanoma cell line. *Molecular Cancer Research* **5** 271–281.
- Kwabi-Addo B, Wang J, Erdem H, Vaid A, Castro P, Ayala G & Ittmann M 2004 The expression of Sproutyl, an inhibitor of fibroblast growth factor signal transduction, is decreased in human prostate cancer. *Cancer Research* **64** 4728–4735.
- Leung HY, Dickson C, Robson CN & Neal DE 1996 Over-expression of fibroblast growth factor-8 in human prostate cancer. *Oncogene* **12** 1833–1835.
- Mohammadi M, Honegger AM, Rotin D, Fischer R, Bellot F, Li W, Dionne CA, Jaye M, Rubinstein M & Schlessinger J 1991 A tyrosine-phosphorylated carboxy-terminal peptide of the fibroblast growth factor receptor (Flg) is a binding site for the SH2 domain of phospholipase C-gamma 1. *Molecular and Cellular Biology* **11** 5068–5078.
- Muddasani P, Norman JC, Ellman M & van Wijnen AJ 2007 Basic fibroblast growth factor activates the MAPK and NFkappa B pathways that converge on Elk-1 to control production of matrix metalloproteinase-13 by human adult articular chondrocytes. *Journal of Biological Chemistry* **282** 31409–31421.
- Nakamoto T, Chang CS, Li AK & Chodak GW 1992 Basic fibroblast growth factor in human prostate cancer cells. *Cancer Research* **52** 571–577.
- Neuwirt H, Pühr M, Cavarretta IT, Mitterberger M, Hobisch A & Culig Z 2007 Suppressor of cytokine signalling-3 is up-regulated by androgen in prostate cancer cell lines and inhibits androgen-mediated proliferation and secretion. *Endocrine-Related Cancer* **14** 1007–1019.
- Nicholson SE, De Souza D, Fabri LJ, Corbin J, Willson TA, Zhang JG, Silva A, Asimakis M, Farley A, Nash AD *et al.* 2000 Suppressor of cytokine signaling-3 preferentially binds to the SHP-2-binding site on the shared cytokine receptor subunit gp130. *PNAS* **97** 6493–6498.
- Ornitz DM, Xu J, Colvin JS, McEwen DG, MacArthur CA, Coulier F, Gao C & Goldfarb M 1996 Receptor specificity of the fibroblast growth factor family. *Journal of Biological Chemistry* **271** 15292–15297.
- Polnaszek N, Kwabi-Addo B, Peterson LE, Ozen M, Greenberg NM, Ortega S, Basilico C & Ittmann M 2003 Fibroblast growth factor 2 promotes tumor progression in an autochthonous mouse model of prostate cancer. *Cancer Research* **63** 5754–5760.
- Powers CJ, McLeskey SW & Wellstein A 2000 Fibroblast growth factors, their receptors and signaling. *Endocrine-Related Cancer* **7** 165–197.

- Puhr M, Santer FR, Neuwirt H, Susani M, Nemeth JA, Hobisch A, Kenner L & Culig Z 2009 Down-regulation of suppressor of cytokine signaling-3 causes prostate cancer cell death through activation of the extrinsic and intrinsic apoptosis pathways. *Cancer Research* **69** 7375–7384.
- Putz T, Culig Z, Eder IE, Nessler-Menardi C, Bartsch G, Grunicke H, Überall F & Klocker H 1999 Epidermal growth factor (EGF) receptor blockade inhibits the action of EGF, insulin-like growth factor I, and a protein kinase A activator on the mitogen-activated protein kinase pathway in prostate cancer cell lines. *Cancer Research* **59** 227–233.
- Ropiquet F, Giri D, Lamb DJ & Ittmann M 1999 FGF7 and FGF2 are increased in benign prostatic hyperplasia and are associated with increased proliferation. *Journal of Urology* **162** 595–599.
- Ropiquet F, Giri D, Kwabi-Addo B, Mansukhani A & Ittmann M 2000 Increased expression of fibroblast growth factor 6 in human prostatic intraepithelial neoplasia and prostate cancer. *Cancer Research* **60** 4245–4250.
- Savinainen KJ, Saramaki OR, Linja MJ, Bratt O, Tammela TL, Isola JJ & Visakorpi T 2002 Expression and gene copy number analysis of ERBB2 oncogene in prostate cancer. *American Journal of Pathology* **160** 339–345.
- Spangenburg EE 2005 SOCS-3 induces myoblast differentiation. *Journal of Biological Chemistry* **280** 10749–10758.
- Spiotto MT & Chung TD 2000 STAT3 mediates IL-6-induced growth inhibition in the human prostate cancer cell line LNCaP. *Prostate* **42** 88–98.
- Steiner H, Godoy-Tundidor S, Rogatsch H, Berger AP, Fuchs D, Comuzzi B, Bartsch G, Hobisch A & Culig Z 2003 Accelerated *in vivo* growth of prostate tumors that up-regulate interleukin-6 is associated with reduced retinoblastoma protein expression and activation of the mitogen-activated protein kinase pathway. *American Journal of Pathology* **162** 655–663.
- Takeuchi R, Matsumoto H, Okada H, Hori M, Gunji A, Hakoziaki K, Akimoto Y & Fujii A 2007 Differences of cell growth and cell cycle regulators induced by basic fibroblast growth factor between nifedipine responders and non-responders. *Journal of Pharmacological Sciences* **103** 168–174.
- Tuxhorn JA, Ayala GE, Smith MJ, Smith VC, Dang TD & Rowley DR 2002 Reactive stroma in human prostate cancer: induction of myofibroblast phenotype and extracellular matrix remodeling. *Clinical Cancer Research* **8** 2912–2923.
- Udayakumar TS, Stratton MS, Nagle RB & Bowden GT 2002 Fibroblast growth factor-1 induced promatrilysin expression through the activation of extracellular-regulated kinases and STAT3. *Neoplasia* **4** 60–67.
- Wang J, Stockton DW & Ittmann M 2004 The fibroblast growth factor receptor-4 Arg388 allele is associated with prostate cancer initiation and progression. *Clinical Cancer Research* **10** 6169–6178.
- Wang J, Thompson B, Ren C, Ittmann M & Kwabi-Addo B 2006 Sprouty4, a suppressor of tumor cell motility, is down regulated by DNA methylation in human prostate cancer. *Prostate* **66** 613–624.
- Wesley UV, McGroarty M & Homoyouni A 2005 Dipeptidyl peptidase inhibits malignant phenotype of prostate cancer cells by blocking basic fibroblast growth factor signaling pathway. *Cancer Research* **65** 1325–1334.
- Wong CP, Bray TM & Ho E 2008 Induction of proinflammatory response in prostate cancer epithelial cells by activated macrophages. *Cancer Letters* **27** 38–46.
- Xia L, Wang L, Chung AS, Ivanov SS, Ling MY, Dragoi AM, Platt A, Gilmer TM, Fu XY & Chin YE 2002 Identification of both positive and negative domains within the epidermal growth factor receptor COOH-terminal region for signal transducer and activator of transcription (STAT) activation. *Journal of Biological Chemistry* **277** 30716–30723.
- Yadav A, Kalita A, Dhillon S & Banerjee K 2005 JAK/STAT3 pathway is involved in survival of neurons in response to insulin-like growth factor and negatively regulated by suppressor of cytokine signaling-3. *Journal of Biological Chemistry* **280** 31830–31840.
- Yang F, Strand DW & Rowley DR 2008 Fibroblast growth factor-2 mediates transforming growth factor-beta action in prostate cancer reactive stroma. *Oncogene* **27** 450–459.
- Yasumoto H, Matsubara A, Mutaguchi K, Usui T & McKeenan WL 2004 Restoration of fibroblast growth factor receptor 2 suppresses growth and tumorigenicity of malignant human prostate carcinoma PC-3 cells. *Prostate* **61** 236–242.