The renin–angiotensin system in the breast and breast cancer

Gavin P Vinson, Stewart Barker and John R Puddfoot

School of Biological and Chemical Sciences, Queen Mary University of London, London E1 4NS, UK

(Correspondence should be addressed to G P Vinson; Email: g.p.vinson@qmul.ac.uk)

Abstract

Much evidence now suggests that angiotensin II has roles in normal functions of the breast that may be altered or attenuated in cancer. Both angiotensin type 1 (AT1) and type 2 (AT2) receptors are present particularly in the secretory epithelium. Additionally, all the elements of a tissue renin–angiotensin system, angiotensinogen, prorenin and angiotensin-converting enzyme (ACE), are also present and distributed in different cell types in a manner suggesting a close relationship with sites of angiotensin II activity. These findings are consistent with the concept that stromal elements and myoepithelium are instrumental in maintaining normal epithelial structure and function. In disease, this system becomes disrupted, particularly in invasive carcinoma. Both AT1 and AT2 receptors are present in tumours and may be up-regulated in some. Experimentally, angiotensin II, acting via the AT1 receptor, increases tumour cell proliferation and angiogenesis, both these are inhibited by blocking its production or function. Epidemiological evidence on the effect of expression levels of ACE or the distribution of ACE or AT1 receptor variants in many types of cancer gives indirect support to these concepts. It is possible that there is a case for the therapeutic use of high doses of ACE inhibitors and AT1 receptor blockers in breast cancer, as there may be for AT2 receptor agonists, though this awaits full investigation. Attention is drawn to the possibility of blocking specific AT1-mediated intracellular signalling pathways, for example by AT1-directed antibodies, which exploit the possibility that the extracellular N-terminus of the AT1 receptor may have previously unsuspected signalling roles.

Endocrine-Related Cancer (2012) 19 R1–R19

Introduction

In the treatment of breast cancer, the various ways of removing the effects of oestrogen, first by surgery and then by the use of drugs, such as tamoxifen and the aromatase inhibitors, which block the actions of oestrogens or prevent their formation, have been hugely successful (Barnes *et al.* 2004, Howell & Dowsett 2004, Jones & Buzdar 2004). Indeed, the critical association between oestrogens, oestrogen receptor (ER) expression and cancer is so entrenched in relation to the breast that the terms ‘receptor-positive’ or ‘receptor-negative’ tumours are a widely accepted shorthand for ER alone (e.g. Yaren *et al.* (2007)).

This long-established connection between oestrogen and breast tumours preceded the more general realisation that the misdirection of normal growth regulatory processes underlies many cancers. Subversion of growth factor receptor structure and function is a well-understood mechanism of oncogene action

(Ross *et al.* 2004, Bianco *et al.* 2005, Hynes & Lane 2005, Pal & Pegram 2005, Zhang *et al.* 2005). In the breast, mechanisms that regulate tissue and tumour growth are multifactorial, and many hormones, growth factors and intracellular signalling pathways are involved (Haagensen 1986, Dickson *et al.* 1992, Hansen & Bissell 2000, Tucker 2000, Pollard 2001, Goffin *et al.* 2002, Singer *et al.* 2003, Lamote *et al.* 2004, Nicolini *et al.* 2006, Cheng *et al.* 2008). Several of these have been targeted for drug development, particularly in tumours that either do not contain ER or are unresponsive to anti-oestrogens.

The systemic renin–angiotensin system (RAS) and the generation of angiotensin II (Fig. 1) has as major roles the regulation of blood pressure, and the adrenal secretion of aldosterone (Mulrow 1999, de Gasparo *et al.* 2000, Kaschina & Unger 2003). The actions of angiotensin II in the regulation of vasoconstriction have even been used to facilitate better accessibility of

Figure 1 The renin–angiotensin system. In normal tissue, and in cancer, the major active hormone is usually considered to be angiotensin II, though angiotensins III and IV, and angiotensin 1–7 have also been implicated (see text).

chemotherapeutic drugs to tumours (Noguchi *et al.* 1988, Goldberg *et al.* 1990, Yamaue *et al.* 1990, Anderson *et al.* 1991). Angiotensins III and IV and angiotensin 1–7 may also be produced, and may act through the same two receptor types as angiotensin II, i.e. angiotensin type 1 and 2 (AT1 and AT2) receptors though with varying effectiveness (de Gasparo *et al.* 2000, Le *et al.* 2002). Angiotensin IV also acts through an insulin-regulated transmembrane enzyme designated as the AT4 receptor (Thomas & Mendelsohn 2003, Chai *et al.* 2004), and angiotensin 1–7 primarily through MasR, the product of the *Mas* oncogene (Neo *et al.* 2010).

Angiotensin receptors AT1 and AT2 are widespread, and they uniformly occur in secretory epithelia. In addition to its functions in the maintenance of blood pressure and hypertension, angiotensin II has also well-studied actions on electrolyte and water transport in the kidney, and elsewhere, including across other epithelial surfaces (Wong *et al.* 1990, Norris *et al.* 1991, Lees *et al.* 1993, Quan & Baum 1996, Wang & Giebisch 1996, Leung *et al.* 1997, Mahmood *et al.* 2002) where it also affects ciliary beat frequency (Saridogan *et al.* 1996a).

Importantly, in many tissues, including the cardiovascular system, adrenal cortex, kidney, liver and perhaps muscle and connective tissue, angiotensin II regulates cell turnover by promoting both cell

proliferation and programmed cell death, perhaps predominantly through differential actions via the AT1 and AT2 receptors (Linz *et al.* 1989, Millan *et al.* 1989, Weber *et al.* 1991, Johnston 1992, Motz *et al.* 1992, Natarajan *et al.* 1992, Wolf & Neilson 1993, Booz & Baker 1995, Schorb *et al.* 1995, Quan & Baum 1996, Vinson & Ho 1998, Kaschina & Unger 2003, Carl-McGrath *et al.* 2007, Billet *et al.* 2008, Moreno & Bataller 2008, Kaschina *et al.* 2009).

Accordingly, it is appropriate to consider angiotensin II among the growth promoting and tissue modelling factors that may be subverted in cancer.

Angiotensin in cancer

Epidemiological evidence

Because angiotensin receptors are widely distributed in epithelia, their possible relevance to cancer, particularly carcinoma, is clear. It is now known that several different types of cancer express angiotensin receptors, and in particular, AT1 and AT2 receptors are expressed in breast cancer (Vinson *et al.* 1995, Marsigliante *et al.* 1996, Inwang *et al.* 1997, Kucerova *et al.* 1998, De Paepe *et al.* 2001, Fujimoto *et al.* 2001, Suganuma *et al.* 2005, Uemura *et al.* 2005b, Gonzalez-Zuloeta Ladd *et al.* 2007, Dolley-Hitze *et al.* 2010, George *et al.* 2010).

Indirect patient evidence supports the role in cancer (Deshayes & Nahmias 2005). Thus, the AT1 receptor has been reported to be up-regulated in various hyperplastic and cancer tissues (De Paepe *et al.* 2001), though not according to all reports (Dinh *et al.* 2002). Additionally, polymorphisms in angiotensinogen, AT1 receptors and angiotensin-converting enzyme (ACE) have been associated with breast cancer risk (Koh *et al.* 2003, 2005, Arima *et al.* 2006, Gonzalez-Zuloeta Ladd *et al.* 2007, Yaren *et al.* 2007, van der Knaap *et al.* 2008, Mendizabal-Ruiz *et al.* 2011). Such polymorphisms have recently been extensively reviewed and discussed (Xi *et al.* 2011). One widely studied polymorphism is a 278 bp *Alu* insertion/deletion (I/D) polymorphism in intron 16 of the *ACE* gene that apparently accounts for 50% of the variability in circulating ACE levels (Rigat *et al.* 1990). Though earlier studies suggested that this had no strong predictive value (Haiman *et al.* 2003), more recently the DD phenotype has been associated with increased risk and poor prognosis in breast cancer (Gonzalez-Zuloeta Ladd *et al.* 2005, Yaren *et al.* 2007, van der Knaap *et al.* 2008, Rosenthal & Gavras 2009). Interestingly, the same ACE polymorphism may also increase risk in benign prostatic hyperplasia (BHP) and prostate cancer, whereas the A1166C substitution in the AT1 receptor increases BHP risk alone (Sierra Diaz *et al.* 2009). Conversely, the C allele carriers have reduced breast cancer risk (Mendizabal-Ruiz *et al.* 2011). Three further AT1 receptor substitutions (A168G, C535T and T825A) have also been associated with reduced breast cancer risk (Koh *et al.* 2005).

More direct patient evidence has been elusive. The first report of the potential utility of ACE inhibitors in preventing cancer development was that of Lever *et al.* (1998) who surveyed data from patients receiving these medications for other reasons, but their findings were not confirmed by others (Meier *et al.* 2000, Li *et al.* 2003, Gonzalez-Perez *et al.* 2004, Ronquist *et al.* 2004, Fryzek *et al.* 2006, Rosenthal & Gavras 2009), nor, in similar patient studies was the use of angiotensin II antagonists in any way linked with the disease (Fryzek *et al.* 2006, Teo 2011). One report suggests that candesartan, an AT1 receptor blocker, when used at a dose similar to that used in patients for other reasons, has beneficial effects in prostate cancer, in that circulating prostate-specific antigen is reduced (Uemura *et al.* 2005a), though this study does not appear to have been repeated. Others have even suggested a modest increase in cancers of all types in patients receiving angiotensin receptor blockers (Sipahi *et al.* 2010), though this too has been contested (Volpe *et al.* 2011). The failure to make such

associations may depend on variations in gene expression, and patients with low ACE expression phenotype may have poorer breast cancer outcomes than high ACE-expressing subjects (Yaren *et al.* 2007), though again there appear to be conflicting findings (Yaren *et al.* 2007, van der Knaap *et al.* 2008). Nevertheless, RAS inhibiting drugs may benefit high ACE-expressing patients but not others (van der Knaap *et al.* 2008). It is possible that anti-RAS drugs are more effective in combination. In patients with advanced pancreatic cancer receiving the nucleoside analogue gemcitabine, lower doses of losartan and other RAS inhibitors were effective in improving outcomes (Nakai *et al.* 2010). One way in which this might occur has been described by Diop-Frimpong *et al.* (2011). Drawing on previous work (Stylianopoulos *et al.* 2010) demonstrating the effect of collagen fibre networks, such as those that occur in connective tissue, on the diffusion of drugs, Diop-Frimpong *et al.* (2011) demonstrated that losartan blocks collagen I production by breast carcinoma-associated fibroblasts, potentially facilitating drug accessibility.

In contrast to the patient data, both ACE inhibitors and AT1 receptor antagonists are effective *in vitro*: they inhibit growth in many different types of tumour cells, including breast cancer cells (Chen *et al.* 1991, Reddy *et al.* 1995, Small *et al.* 1997, Rivera *et al.* 2001, Uemura *et al.* 2008, Inigo *et al.* 2009, Ino *et al.* 2011). In experimental animals *in vivo*, on the other hand, for example in xenografts of SKOV-3 ovarian tumour in mice or of C6 rat glioma cell tumours in rats, much higher doses of candesartan and losartan, respectively, were needed to demonstrate tumour regression than those generally used in patients (Rivera *et al.* 2001, Suganuma *et al.* 2005). It is perhaps because high doses are required when these drugs alone are used that the epidemiological studies on patients receiving anti-hypertensive treatment for cardiovascular disease show no benefit in incidence of cancer.

The discovery of the zinc metalloprotease ACE2 introduced a new aspect of angiotensin signalling (Donoghue *et al.* 2000). ACE2 preferentially hydrolyses angiotensin I to angiotensin 1–9, and angiotensin II to angiotensin 1–7 (Fleming *et al.* 2006; see Fig. 1). Angiotensin 1–7 has properties different from those of angiotensin II and may oppose angiotensin II functions. In particular, it is anti-proliferative and reduces fibrosis in breast tumours, and angiogenesis in lung tumours (Menon *et al.* 2007, Soto-Pantoja *et al.* 2009, Cook *et al.* 2010, Gallagher *et al.* 2011), and has been used with benefit in phase I patient trials (Petty *et al.* 2009).

Angiogenesis

Additionally, although both ACE inhibitors and AT1 receptor antagonists may be effective on animal tumours *in vivo*, the results are more ambiguous than *in vitro*, and at least in part could be due to their anti-angiogenic actions (Volpert *et al.* 1996, Fujimoto *et al.* 2001, Fujita *et al.* 2002, 2005, Yoshiji *et al.* 2004, Kosaka *et al.* 2007, De Paepe 2009, Miyajima *et al.* 2009). The important part played by angiogenesis in the development of cancer has frequently been emphasised. There is considerable evidence that cancer growth and spread is angiogenesis dependent, tumour cells themselves can produce angiogenic factors and inhibition of angiogenesis can limit tumour growth (Weidner 2004, Sharma *et al.* 2005, Clapp *et al.* 2009), including in the breast (Heffelfinger 2007, Groves *et al.* 2011). It is difficult to assess the importance of this process in relation to the direct effects of angiotensin II on tumour growth and cell proliferation. Certainly, angiotensin II is involved in angiogenesis. Several *in vitro* studies have shown that vascular endothelial growth factor (VEGF) expression is stimulated by angiotensin II or inhibited by ACE or angiotensin blockers in tumour cells, including squamous cell (Yasumatsu *et al.* 2004) ovarian (Suganuma *et al.* 2005), prostate (Kosaka *et al.* 2007) and rat pituitary tumour cells (Ptasinska-Wnuk *et al.* 2007). Similar conclusions have been reached from *in vivo* studies. Thus, AT1 receptor expression and angiogenesis were correlated in ovarian tumours and in astrocytomas (Ino *et al.* 2006, Arrieta *et al.* 2008). Angiotensin II supported VEGF production and angiogenesis in xenografts of ovarian cancer cells (Suganuma *et al.* 2005) and AT1 receptor blockade inhibited both of these actions in xenografts of ovarian and gastric tumour cells (Suganuma *et al.* 2005, Huang *et al.* 2008). AT1 receptor blockade also inhibited angiogenesis in murine Lewis lung tumours (Imai *et al.* 2007) and through this means enhanced the effectiveness of radiation treatment in murine melanoma (Ohnuma *et al.* 2009, Otake *et al.* 2009) and in murine renal tumours (Miyajima *et al.* 2009). However, in *in vivo* studies in which S-180 murine sarcoma cell tumours were developed in AT1a receptor null mice, angiogenesis, along with VEGF expression, was both reduced and partially refractory to AT1 receptor blockade when compared with normal tissue. Hence, host angiotensin II activity is instrumental in supporting angiogenesis in host stromal cells in addition to any effect it has on the cancer cells themselves (Fujita *et al.* 2002, 2005, Imai *et al.* 2007).

Actions of angiotensin II on breast cancer cells

As in other tissues, angiotensin II acts on the AT1 receptor to promote cell proliferation in breast cancer cells (Muscella *et al.* 2002). The AT1-mediated signalling involves the protein kinase C (PKC, zeta and iota)/Ca²⁺/inositol trisphosphate (IP3) pathways, and also extracellular signal-related kinase (ERK) activation (Greco *et al.* 2002a,b, 2003, Muscella *et al.* 2003, 2005). Angiotensin II also activates Na⁺/K⁺ ATPase (Muscella *et al.* 2002, 2005).

Angiotensin II has further possible roles involved in cell adhesion and invasion. Specifically, again acting via the AT1 receptor, it inhibits expression of integrin subtypes $\alpha 3$ and $\beta 1$ and also binding to and invasion through components of the extracellular matrix. In contrast to its actions on proliferation, these effects of angiotensin II may be regarded as potentially beneficial (Puddefoot *et al.* 2006). Consequently, RAS blockade may not always be entirely an appropriate therapy in cancer, perhaps also explaining its apparent lack of benefit in patients. Conflicting evidence on the efficacy of anti-RAS treatment has also been discussed in the context of cardiovascular disease (Magy *et al.* 2005).

Angiotensin II, ER and growth factors

Because of the well-known importance of ER and growth factors and their interrelationship in breast cancer, it is relevant to examine their interactions with the RAS. The interrelationship between the RAS and ER is complex. Depending on the tissue, oestrogen has varyingly been reported to down-regulate AT1 receptors, in rat pituitary and hypothalamus (Seltzer *et al.* 1992, Kisley *et al.* 1999) and dog kidney, myocardium, liver and adrenal (Owonikoko *et al.* 2004; see also Fischer *et al.* (2002), but to up-regulate them at other sites, including rat kidney (Baiardi *et al.* 2005) and sheep uterine artery endothelium (Sullivan *et al.* 2005). Consistent with RAS up-regulation by oestrogen, intensity of AT1 receptor staining is most intense in the periovulatory period in human fallopian tube and uterine epithelia (Saridogan *et al.* 1996a,b) and the AT2 receptor is also high during the proliferative phase in human myometrium (Pucell *et al.* 1987, Mancina *et al.* 1996), as it is in the rat ovary (Pucell *et al.* 1987, Mancina *et al.* 1996). However, such changes do not necessarily reflect the functions of the RAS as a whole and other RAS components may respond independently, for example renin and ACE activities are reduced in various tissues by oestrogen (Fischer

et al. 2002), though angiotensinogen is increased (Gordon *et al.* 1992, Klett *et al.* 1993, Fischer *et al.* 2002). Oestrogen stimulates plasma renin activity (PRA) and RAS activity in sheep (Magness *et al.* 1993), though in women high PRA is associated with the luteal phase (Sealey *et al.* 1994, Chapman *et al.* 1997, Chidambaram *et al.* 2002). In breast duct cancer cells, angiotensin II treatment *in vitro* reduces ER and increases PR (Small *et al.* 1997).

The relationship between ER and AT1 is thus incompletely resolved. It may be that angiotensin II signalling is more significant in ER-negative breast tumours (Herr *et al.* 2008), in which a role has been postulated for AT1 receptors in the non-genomic response to oestrogen (Lim *et al.* 2006), though there is a subset of ER-positive (and ERBB2-negative) tumours that shows marked overexpression of AT1 receptors (Rhodes *et al.* 2009). This appears to contrast with vascular smooth muscle cells in which the ER blocker raloxifene (in the presence of oestradiol) inhibited angiotensin II-stimulated proliferation (Wang *et al.* 2007).

Angiotensin receptor signalling also interacts with growth factors in breast cancer cells. Thus, ERKs are activated by angiotensin II directly via PKC and indirectly via epidermal growth factor receptor (EGFR)-mediated phosphatidylinositol-3 kinase/serine-threonine protein kinases (PI3-kinase/Akt/mTOR/p70S6K1) signalling pathways (Greco *et al.* 2002b, 2003, Chiu *et al.* 2005, Han *et al.* 2007). In more detail, the AT1 receptor, linked to Gq/11, signals both by Ca^{2+} /IP3 and by diacylglycerol-linked events, and also by tyrosine kinase activation, including via EGFR-linked PI3-kinase and Akt signalling, with subsequent activation of ERK1 and ERK2 (Greco *et al.* 2003, Shah *et al.* 2004, Han *et al.* 2007, Kim *et al.* 2009). Such EGFR activation is at least in part mediated via angiotensin II-stimulated metalloproteinase activity (Liebmann 2011, Smith *et al.* 2011; see below). There is extensive crosstalk with other receptors, including insulin and growth factor signalling pathways (Shah *et al.* 2006, Redondo *et al.* 2007, Escano *et al.* 2008, Muscogiuri *et al.* 2008, Olivares-Reyes *et al.* 2009, Arellano-Plancarte *et al.* 2010). Conversely, the AT2 receptor is thought to activate phosphatase activity and block AT1 receptor-mediated intracellular signalling events, including phospholipase activation and the phosphorylation of signalling components. These pathways have been extensively discussed elsewhere (de Gasparo *et al.* 2000, de Gasparo 2002, Kaschina & Unger 2003, Deshayes & Nahmias 2005, Louis *et al.* 2010, Zhao *et al.* 2010).

The local RAS in the breast

Tissue remodelling and matrix metalloproteinases

In the normal cycle of events in the breast, the ductal system, which begins to develop in puberty, stabilises in the adult but proliferates extensively during pregnancy to enable production of a high level of secretory activity during lactation. After lactation ceases, the ducts undergo apoptotic involution (Fig. 2; Wiseman & Werb 2002, Boutinaud *et al.* 2004, Green & Streuli 2004). Because of the relationship between the stage of the cycle and the incidence of metaplastic change, Villadsen (2005) and Russo *et al.* (2006) postulated that there are at least two types, or a hierarchy, of stem cells. The whole process does not involve the ducts alone, and stromal cells and their products, including growth factors and integrins, are also strongly implicated (Chrenek *et al.* 2001, Pollard 2001, Wiseman & Werb 2002, Barcellos-Hoff & Medina 2005, Zechmann *et al.* 2007). Because of its sites of origin, described below, and the location of its receptors, it is appropriate to consider angiotensin II among these factors and that, perhaps acting through both receptor types, it is instrumental in both proliferative and apoptotic phases of the normal cycle.

The breast cycle (Fig. 2) and its sequence of development and resorption reflect, among other things, synthesis and proteolysis of proteins of the extracellular matrix and the basement membrane, such as collagen, in a balanced manner (Morini *et al.* 2000, Sun *et al.* 2006). Hydrolysis of extracellular matrix proteins is catalysed at the basement membrane by the zinc-dependent matrix metalloproteinases (MMPs) present in stromal and secretory cells of normal and diseased tissue (Werb *et al.* 1996, Lebeau *et al.* 1999, Bodey *et al.* 2001). Accordingly, these enzymes are also involved in the invasive process (Ambili *et al.* 1998, Rudolph-Owen & Matrisian 1998) and high MMP levels are associated with poor outcomes (Duffy *et al.* 2000). Because epithelial cells depend on the functions of the basement membrane and their constituents, protein breakdown contributes to epithelial dysfunction. In many tissues, angiotensin II plays a key part in such tissue remodelling, and it affects both MMP activity and collagen synthesis (Gack *et al.* 1994, Ford *et al.* 1999, Dzau 2001, Galis & Khatri 2002, Shah *et al.* 2004, Chiu *et al.* 2005, Yang *et al.* 2005, Karakiulakis *et al.* 2007, Kim *et al.* 2007). As MMPs are located in myoepithelial cells, like prorenin (see below), it is clear that locally produced angiotensin II may have such a role in the breast.

Figure 2 The breast cycle: note particularly extensive duct and gland development during pregnancy and lactation, followed by apoptotic involution when lactation ceases (cf. [Wiseman & Werb \(2002\)](#), [Boutinaud et al. \(2004\)](#) and [Green & Streuli \(2004\)](#)). Drawing by Bronwen Vinson. Reproduced from [Vinson et al. \(2007\)](#) with kind permission from Springer Science and Business Media.

All the functions of angiotensin II described so far acquire an additional perspective in the light of our understanding of the tissue-based RAS. This is because the significant factor in both normal function and in disease may not be the angiotensin II in the blood, but that which is locally produced, within the tissue.

That many organ systems contain discrete RASs has been well understood for some time: such localised systems have been described in many tissues, including the kidney, liver and adrenal ([Phillips et al. 1993](#), [Gupta et al. 1995](#), [Zimmerman & Dunham 1997](#), [Mulrow 1998](#), [Vinson & Ho 1998](#), [Neo et al. 2010](#)), brain, pituitary and reproductive system ([Hagemann et al. 1994](#), [Ganong 1995](#), [Nielsen et al. 1995](#), [Vinson et al. 1997](#), [Vila-Porcile & Corvol 1998](#), [McKinley et al. 2003](#), [Li et al. 2004](#), [Dean et al. 2006](#)), pancreas ([Tahmasebi et al. 1999](#), [Leung & Carlsson 2001, 2005](#), [Lau & Leung 2011](#)), lung ([Feng et al. 2010](#)) and heart ([Okura et al. 1992](#), [Bader 2002](#), [Dean et al. 2006](#)).

These tissue RASs may be perturbed in cancer. For example, in a mouse model of colorectal cancer metastases, ACE expression was increased (though ACE2 was decreased) in tumour-bearing livers, as well as in the tumours themselves. Tumour volume was

reduced by the ACE inhibitor captopril. Liver angiotensinogen was unaffected by the tumours and decreased in captopril treatment, whereas ACE in both liver and tumour tissues was further increased. AT1 receptor expression was elevated by tumour induction and reduced by captopril: MasR, the putative receptor for angiotensin 1–7, was increased by captopril ([Neo et al. 2010](#)). The possibility that angiotensin III may have a specific role has also been suggested in studies on rats with *N*-methyl nitrosourea-induced breast tumours, in which soluble and membrane-bound aspartyl and glutamyl aminopeptidase activities are increased whereas soluble aminopeptidase N and B activities are decreased, both of which potentially increase angiotensin III production, with reduced angiotensins II and IV ([del Pilar Carrera et al. 2010](#)).

Localisation of RAS components

In studies on the sites of (pro)renin gene transcription, (pro)renin mRNA was found in most of the breast samples examined, invariably in close proximity to the ductal epithelium but not within the epithelium itself. Prorenin mRNA was abundant in the stroma immediately adjacent to the ducts, in myoepithelial cells in

normal tissue and in early cancer stages but tended to be lost from both sites in more advanced disease, paralleling the partial loss of AT1 receptors (Tahmasebi *et al.* 1998; Fig. 3). Confirmation of these findings, and evidence for other RAS components, was obtained using quantitative RT-PCR and the presence of RNA coding for angiotensinogen, prorenin, ACE and both AT1 and AT2 receptors was demonstrated in normal and diseased breast tissues, supporting the hypothesis that a tissue RAS is present in the breast. As in the *in situ* hybridisation data (Tahmasebi *et al.* 1998), there was significantly less (pro)renin mRNA in carcinoma than in normal tissue, and indeed, ACE and angiotensinogen mRNAs were also reduced in carcinoma compared with normal tissue (Tahmasebi *et al.* 2006). This reflects the earlier finding that AT1 receptors are reduced in advanced tumours.

mRNA coding for prorenin was distributed between myoepithelium and, most extensively in fibroblasts and connective tissue close to the ducts. Conversely, prorenin protein itself was mostly present in myoepithelial cells and absent from the connective tissue. Of course, this distribution could represent differences in

mRNA translation between the two cell types, but a rather different picture emerges in cancer. Although the distribution of prorenin and its mRNA in ductal and in lobular carcinoma *in situ* was similar to normal, in more advanced conditions, as the myoepithelium was lost, prorenin protein was only sparsely present in the epithelium, but it was located in fibroblasts. Here, though always present, it appeared to decrease in amount as malignancy advanced (Tahmasebi *et al.* 1998; Fig. 3). Two possibilities present themselves, one is that the prorenin mRNA that is ever present in breast fibroblasts is translated only in cancer. Alternatively, it is always translated, even in normal tissue, but the prorenin formed is normally transported elsewhere, to the myoepithelium or to the epithelium (though this latter is not frequently observed). Whatever the explanation, it is evident that the functions of the breast RAS may be greatly perturbed in cancer. Similar processes may well occur in other types of cancer, for example in the pancreas (Lau & Leung 2011).

There is a difficulty in testing this concept of an entirely localised RAS in any tissue – what can *in situ*

Figure 3 Both angiotensin II receptors and ACE are present in epithelial cells and in cancer cells. Sites of (pro)renin mRNA transcription (dark shading) are shown in (i) normal breast ducts, (ii) intraductal carcinoma *in situ* and (iii) invasive carcinoma. The myoepithelial source of (pro)renin transcription is lost as cancer develops. As in normal tissue this lies in close proximity to the epithelium, the configuration strongly suggests that angiotensin II can be produced at its epithelial site of action. This tightly linked system is lost in cancer, suggesting that the AT1 and AT2 receptor-containing carcinoma eventually becomes deprived of its source of angiotensin II. Adapted from Tahmasebi *et al.* (1998, 2006). e, epithelium; m, myoepithelium; f, fibroblast; t, tumour; s, stroma. Drawing by Bronwen Vinson. Reproduced from Vinson *et al.* (2007) with kind permission from Springer Science and Business Media.

hybridisation or immunocytochemistry reveal about the state of activation of any of the components? Prorenin provides a key example here: the methods used in the papers cited above do not distinguish between the cleaved or activated forms. The primary mechanism for prorenin activation has been considered to be through cleavage by prohormone convertases (Benjannet *et al.* 1992), which may be highly expressed in cancer, including breast tumours, and this is associated with greater oestrogen dependency (Cheng *et al.* 1997, 2001). Of course, prohormone convertases may be involved in tumorigenic processes that do not involve either prorenin or angiotensin II (Siegfried *et al.* 2003, Scamuffa *et al.* 2008). Alternatively, the discovery of a specific prorenin receptor that binds prorenin and activates intracellular signalling pathways while at the same time activating its enzymic activity in the absence of cleavage opens new possibilities (Nguyen & Contrepas 2008, Nguyen 2011). These and related questions of activation of the breast RAS will need to be addressed in future.

Like the AT1 receptor, ACE is present in the secretory epithelium of the normal breast and also in diseased breast tissue (Tahmasebi *et al.* 2006), suggesting that angiotensin II may be formed directly in the cells on which it acts. However, in cancer, the overall loss and changes in the distribution of prorenin described above may mean that as the disease progresses, neither substrate for the enzyme nor ligand for the receptor remains available. Malignancy is thus correlated with the deregulation of RAS function.

These findings and this proposed mechanism have considerable resonance with other authors' concepts of the role of the stromal and myoepithelial interaction with the secretory epithelium and with cancer. Kalluri & Weinberg (2009) have proposed that one class of epithelial–mesenchymal transformations (EMTs), which they call type 3 EMT (to distinguish from implantation and wound-healing forms, types 1 and 2), is characteristic of the transformation of polarised and highly differentiated epithelial cells into mesenchymal cells. Such cells secrete extracellular matrix components and are highly mobile and invasive, though there is a reverse transformation (MET) at sites of metastatic colonisation (Kalluri 2009, Kalluri & Weinberg 2009). EMTs, of whatever type, are initiated as the epithelial cells that invade through the basement membrane. A number of factors are thought to be involved, including insulin-like growth factor, transforming growth factor β , platelet-derived growth factor, integrins and the signalling pathways they evoke, with all of which AT1 and AT2 receptors may interact.

So what retains epithelia in their normal functional state? Here, the focus is on myoepithelial cells. These too are now known to be important in cancer progression. First rather overlooked, as they only infrequently produce tumours, they are now thought to be natural tumour suppressors because of their role in maintaining epithelial cell polarity and cell cycle progression and inhibiting cell migration and invasion (Lakhani & O'Hare 2001, Polyak & Hu 2005). This has been postulated to be due to the secretion of proteinase and angiogenic inhibitors (Barsky & Karlin 2005). Additionally, as well as inflammatory cells, fibroblasts have also been thought to be the source of factors affecting tumour development (Tlsty & Coussens 2006). These concepts received direct experimental support when MCF7 breast cancer cells were grown *in vitro* in an environment of extracellular matrices of type 1 collagen, or reconstituted basement membrane proteins, together with human fibroblasts. Surviving cells in the presence of collagen organised into clusters, while the further addition of basement membrane proteins induced MCF7 cell polarisation and the formation of lumina, and the presence of fibroblasts induced the formation of elongated structures (Krause *et al.* 2010). Furthermore, differences in gene expression between core biopsies of breast tumours with varying degrees of stromal content were taken to indicate the influence of the stroma (Cleator *et al.* 2006). So the stromal and myoepithelial localisation of RAS components strongly suggests that angiotensin II may have an important, possibly crucial role in this context.

Implications for therapy

One way in which beneficial advances have been made despite initially discouraging data has been to identify subsets of patients who may benefit where others may not. A key example here is in the identification of a subgroup of breast tumours that overexpress the ERBB2 (HER2) tyrosine kinase receptor and are thus sensitive to the monoclonal antibody trastuzumab (Nahta *et al.* 2006, Nahta & Esteva 2007). More and more it becomes clear that patient profiling in this way yields benefit, and this may well be true for the RAS in breast. It is known that a significant subset of breast tumours overexpress the AT1 receptor, and although there are various mechanisms for this, one way may be that AT1 receptor expression is directly controlled by ER, leading to a subset of ER-positive, ERBB2-negative tumours that overexpress AT1 receptor (Ateeq *et al.* 2009, Rhodes *et al.* 2009).

Because of the possibility of both beneficial and disadvantageous effects of AT1 receptor inhibition, it is worth exploring whether means exist to selectively inhibit individual signalling events. This possibility has been discussed in a recent review, in which the signalling roles of individual domains of the receptor were explored, though the possibility that the extra-cellular N-terminal domain might be involved was not considered (Aplin *et al.* 2009). There may, however, be good reasons to consider the N-terminus in this light because there appear to be ligand binding or signalling determinants in this region (Hjorth *et al.* 1994, Oliveira *et al.* 2007), and a particular role for Arg23 has been identified (Santos *et al.* 2004).

In this respect, the activity of monoclonal antibody 6313/G2 directed against a sequence in the N-terminal domain of the AT1 receptor has provided further information, as it appears to enhance some signalling pathways while inhibiting others. Though not affecting angiotensin II binding to the receptor (Barker *et al.* 1993) the antibody directly stimulates aldosterone secretion via the IP3 pathway in rat glomerulosa cells *in vitro*, though it also blocks PKC activation, apparently by interrupting receptor internalization (Kapas *et al.* 1994, Vinson *et al.* 1994). In other studies on rat vascular smooth muscle cells, basal and angiotensin-stimulated tritiated thymidine incorporation into rat arterial smooth muscle cells was inhibited by 6313/G2, inducing a transient increase in intracellular calcium in cultured rat arterial smooth muscle cells, but reducing PKC and MAPK signal transduction (Xiao *et al.* 2008). A short-chain fragment variable of this antibody also blocked AT1 receptor-mediated caspase-3/7 inhibition in breast cancer cells and dose dependently gave significant tumour regression in breast cell xenografts *in vivo*. These data support the view that differential inhibition of angiotensin II-stimulated signalling pathways may be achieved in this way (Redondo-Muller *et al.* 2008).

Conclusions

There can now be no doubt that the RAS is involved both in the normal physiology (and perhaps development) of the breast and in the ontogeny of breast carcinoma, and possibly other cancers. There is strong evidence that blocking the pathways of AT1 receptor-mediated angiotensin signalling can have beneficial effects. However, in view of the multiple actions of angiotensin II on breast cancer cells, some of which themselves may be considered to be beneficial, this is not without potential cost. In identifying the AT1 receptor as a new target for breast cancer therapy,

development of agents that more precisely discriminate between individual signalling pathways is an important goal. The monoclonal antibody 6313/G2 and its recombinant counterpart demonstrate that this kind of approach may be entirely feasible.

Declaration of interest

Queen Mary, University of London owns IP related to antibodies against the AT1 receptor, currently licensed to Oncobiopharm Ltd.

Funding

This review did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

References

- Ambili M, Jayasree K & Sudhakaran PR 1998 60K gelatinase involved in mammary gland involution is regulated by beta-oestradiol. *Biochimica et Biophysica Acta* **1403** 219–231. (doi:10.1016/S0167-4889(98)00030-5)
- Anderson JH, Willmott N, Bessent R, Angerson WJ, Kerr DJ & McArdle CS 1991 Regional chemotherapy for inoperable renal carcinoma: a method of targeting therapeutic microspheres to tumour. *British Journal of Cancer* **64** 365–368. (doi:10.1038/bjc.1991.308)
- Aplin M, Bonde MM & Hansen JL 2009 Molecular determinants of angiotensin II type 1 receptor functional selectivity. *Journal of Molecular and Cellular Cardiology* **46** 15–24. (doi:10.1016/j.yjmcc.2008.09.123)
- Arellano-Plancarte A, Hernandez-Aranda J, Catt KJ & Olivares-Reyes JA 2010 Angiotensin-induced EGF receptor transactivation inhibits insulin signaling in C9 hepatic cells. *Biochemical Pharmacology* **79** 733–745. (doi:10.1016/j.bcp.2009.10.014)
- Arima H, Kiyohara Y, Tanizaki Y, Nakabeppu Y, Kubo M, Kato I, Sueishi K, Tsuneyoshi M, Fujishima M & Iida M 2006 Angiotensin I-converting enzyme gene polymorphism modifies the smoking-cancer association: the Hisayama study. *European Journal of Cancer Prevention* **15** 196–201. (doi:10.1097/01.cej.0000199506.15571.37)
- Arrieta O, Pineda-Olvera B, Guevara-Salazar P, Hernandez-Pedro N, Morales-Espinosa D, Ceron-Lizarraga TL, Gonzalez-De la Rosa CH, Rembao D, Segura-Pacheco B & Sotelo J 2008 Expression of AT1 and AT2 angiotensin receptors in astrocytomas is associated with poor prognosis. *British Journal of Cancer* **99** 160–166. (doi:10.1038/sj.bjc.6604431)
- Ateeq B, Tomlins SA & Chinnaiyan AM 2009 AGTR1 as a therapeutic target in ER-positive and ERBB2-negative breast cancer cases. *Cell Cycle* **8** 3794–3795. (doi:10.4161/cc.8.23.9976)

- Bader M 2002 Role of the local renin–angiotensin system in cardiac damage: a minireview focussing on transgenic animal models. *Journal of Molecular and Cellular Cardiology* **34** 1455–1462. (doi:10.1006/jmcc.2002.2077)
- Baiardi G, Macova M, Armando I, Ando H, Tyurmin D & Saavedra JM 2005 Estrogen upregulates renal angiotensin II AT1 and AT2 receptors in the rat. *Regulatory Peptides* **124** 7–17. (doi:10.1016/j.regpep.2004.06.021)
- Barcellos-Hoff MH & Medina D 2005 New highlights on stroma–epithelial interactions in breast cancer. *Breast Cancer Research* **7** 33–36. (doi:10.1186/bcr972)
- Barker S, Marchant W, Ho MM, Puddefoot JR, Hinson JP, Clark AJL & Vinson GP 1993 A monoclonal antibody to a conserved sequence in the extracellular domain recognizes the angiotensin II AT1 receptor in mammalian tissues. *Journal of Molecular Endocrinology* **11** 241–245. (doi:10.1677/jme.0.0110241)
- Barnes DM, Millis RR, Gillett CE, Ryder K, Skilton D, Fentiman IS & Rubens RD 2004 The interaction of oestrogen receptor status and pathological features with adjuvant treatment in relation to survival in patients with operable breast cancer: a retrospective study of 2660 patients. *Endocrine-Related Cancer* **11** 85–96. (doi:10.1677/erc.0.0110085)
- Barsky SH & Karlin NJ 2005 Myoepithelial cells: autocrine and paracrine suppressors of breast cancer progression. *Journal of Mammary Gland Biology and Neoplasia* **10** 249–260. (doi:10.1007/s10911-005-9585-5)
- Benjannet S, Reudelhuber T, Mercure C, Rondeau N, Chretien M & Seidah NG 1992 Proprotein conversion is determined by a multiplicity of factors including convertase processing, substrate specificity, and intracellular environment. Cell type-specific processing of human prorenin by the convertase PC1. *Journal of Biological Chemistry* **267** 11417–11423.
- Bianco R, Daniele G, Ciardiello F & Tortora G 2005 Monoclonal antibodies targeting the epidermal growth factor receptor. *Current Drug Targets* **6** 275–287. (doi:10.2174/1389450053765842)
- Billet S, Aguilar F, Baudry C & Clauser E 2008 Role of angiotensin II AT1 receptor activation in cardiovascular diseases. *Kidney International* **74** 1379–1384. (doi:10.1038/ki.2008.358)
- Bodey B, Bodey B Jr, Siegel SE & Kaiser HE 2001 Matrix metalloproteinases in neoplasm-induced extracellular matrix remodeling in breast carcinomas. *Anticancer Research* **21** 2021–2028.
- Booz GW & Baker KM 1995 Molecular signalling mechanisms controlling growth and function of cardiac fibroblasts. *Cardiovascular Research* **30** 537–543.
- Boutinaud M, Guinard-Flament J & Jammes H 2004 The number and activity of mammary epithelial cells, determining factors for milk production. *Reproduction, Nutrition, Development* **44** 499–508. (doi:10.1051/rnd:2004054)
- Carl-McGrath S, Ebert MP, Lendeckel U & Rocken C 2007 Expression of the local angiotensin II system in gastric cancer may facilitate lymphatic invasion and nodal spread. *Cancer Biology & Therapy* **6** 1218–1226.
- Chai SY, Fernando R, Peck G, Ye SY, Mendelsohn FA, Jenkins TA & Albiston AL 2004 The angiotensin IV/AT4 receptor. *Cellular and Molecular Life Sciences* **61** 2728–2737. (doi:10.1007/s00018-004-4246-1)
- Chapman AB, Zamudio S, Woodmansee W, Merouani A, Osorio F, Johnson A, Moore LG, Dahms T, Coffin C, Abraham WT et al. 1997 Systemic and renal hemodynamic changes in the luteal phase of the menstrual cycle mimic early pregnancy. *American Journal of Physiology. Renal Physiology* **42** F777–F782.
- Chen L, Re RN, Prakash O & Mondal D 1991 Angiotensin-converting enzyme-inhibition reduces neuroblastoma cell-growth rate. *Proceedings of the Society for Experimental Biology and Medicine* **196** 280–283.
- Cheng M, Watson PH, Paterson JA, Seidah N, Chretien M & Shiu RP 1997 Pro-protein convertase gene expression in human breast cancer. *International Journal of Cancer* **71** 966–971. (doi:10.1002/(SICI)1097-0215(19970611)71:6<966::AID-IJC10>3.0.CO;2-4)
- Cheng M, Xu N, Iwasio B, Seidah N, Chretien M & Shiu RP 2001 Elevated expression of proprotein convertases alters breast cancer cell growth in response to estrogen and tamoxifen. *Journal of Molecular Endocrinology* **26** 95–105. (doi:10.1677/jme.0.0260095)
- Cheng N, Chytil A, Shyr Y, Joly A & Moses HL 2008 Transforming growth factor-beta signaling-deficient fibroblasts enhance hepatocyte growth factor signaling in mammary carcinoma cells to promote scattering and invasion. *Molecular Cancer Research* **6** 1521–1533. (doi:10.1158/1541-7786.MCR-07-2203)
- Chidambaram M, Duncan JA, Lai VS, Cattran DC, Floras JS, Scholey JW & Miller JA 2002 Variation in the renin angiotensin system throughout the normal menstrual cycle. *Journal of the American Society of Nephrology* **13** 446–452.
- Chiu T, Santiskulvong C & Rozengurt E 2005 EGF receptor transactivation mediates ANG II-stimulated mitogenesis in intestinal epithelial cells through the PI3-kinase/Akt/mTOR/p70S6K1 signaling pathway. *American Journal of Physiology. Gastrointestinal and Liver Physiology* **288** G182–G194. (doi:10.1152/ajpgi.00200.2004)
- Chrenek MA, Wong P & Weaver VM 2001 Tumour–stromal interactions. Integrins and cell adhesions as modulators of mammary cell survival and transformation. *Breast Cancer Research* **3** 224–229. (doi:10.1186/bcr300)
- Clapp C, Thebault S, Jezierski MC & Martinez De La Escalera G 2009 Peptide hormone regulation of angiogenesis. *Physiological Reviews* **89** 1177–1215. (doi:10.1152/physrev.00024.2009)
- Cleator SJ, Powles TJ, Dexter T, Fulford L, Mackay A, Smith IE, Valgeirsson H, Ashworth A & Dowsett M 2006

- The effect of the stromal component of breast tumours on prediction of clinical outcome using gene expression microarray analysis. *Breast Cancer Research* **8** R32. (doi:10.1186/bcr1506)
- Cook KL, Metheny-Barlow LJ, Tallant EA & Gallagher PE 2010 Angiotensin-(1–7) reduces fibrosis in orthotopic breast tumors. *Cancer Research* **70** 8319–8328. (doi:10.1158/0008-5472.CAN-10-1136)
- Dean SA, Tan J, White R, O'Brien ER & Leenen FH 2006 Regulation of components of the brain and cardiac renin-angiotensin systems by 17 β -estradiol following myocardial infarction in female rats. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* **291** R155–R162. (doi:10.1152/ajpregu.00497.2005)
- De Paepe B 2009 Anti-angiogenic agents and cancer: current insights and future perspectives. *Recent Patents on Anti-Cancer Drug Discovery* **4** 180–185. (doi:10.2174/157489209788452821)
- De Paepe B, Verstraeten VL, De Potter CR, Vakaet LA & Bullock GR 2001 Growth stimulatory angiotensin II type-1 receptor is upregulated in breast hyperplasia and *in situ* carcinoma but not in invasive carcinoma. *Histochemistry and Cell Biology* **116** 247–254.
- Deshayes F & Nahmias C 2005 Angiotensin receptors: a new role in cancer? *Trends in Endocrinology and Metabolism* **16** 293–299. (doi:10.1016/j.tem.2005.07.009)
- Dickson RB, Johnson MD, Bano M, Shi E, Kurebayashi J, Ziff B, Martinezlacaci I, Amundadottir LT & Lippman ME 1992 Growth-factors in breast-cancer – mitogenesis to transformation. *Journal of Steroid Biochemistry and Molecular Biology* **43** 69–78. (doi:10.1016/0960-0760(92)90189-P)
- Dinh DT, Frauman AG, Somers GR, Ohishi M, Zhou J, Casley DJ, Johnston CI & Fabiani ME 2002 Evidence for activation of the renin-angiotensin system in the human prostate: increased angiotensin II and reduced AT(1) receptor expression in benign prostatic hyperplasia. *Journal of Pathology* **196** 213–229. (doi:10.1002/path.1021)
- Diop-Frimpong B, Chauhan VP, Krane S, Boucher Y & Jain RK 2011 Losartan inhibits collagen I synthesis and improves the distribution and efficacy of nanotherapeutics in tumors. *PNAS* **108** 2909–2914. (doi:10.1073/pnas.1018892108)
- Dolley-Hitze T, Jouan F, Martin B, Mottier S, Edeline J, Moranne O, Le Pogamp P, Belaud-Rotureau MA, Patard JJ, Rioux-Leclercq N *et al.* 2010 Angiotensin-2 receptors (AT1-R and AT2-R), new prognostic factors for renal clear-cell carcinoma? *British Journal of Cancer* **103** 1698–1705. (doi:10.1038/sj.bjc.6605866)
- Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R *et al.* 2000 A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1–9. *Circulation Research* **87** E1–E9.
- Duffy MJ, Maguire TM, Hill A, McDermott E & O'Higgins N 2000 Metalloproteinases: role in breast carcinogenesis, invasion and metastasis. *Breast Cancer Research* **2** 252–257. (doi:10.1186/bcr65)
- Dzau VJ 2001 Theodore Cooper Lecture: tissue angiotensin and pathobiology of vascular disease: a unifying hypothesis. *Hypertension* **37** 1047–1052.
- Escano CS Jr, Keever LB, Gutweiler AA & Andresen BT 2008 Angiotensin II activates extracellular signal-regulated kinase independently of receptor tyrosine kinases in renal smooth muscle cells: implications for blood pressure regulation. *Journal of Pharmacological and Experimental Therapeutics* **324** 34–42. (doi:10.1124/jpet.107.126300)
- Feng Y, Wan H, Liu J, Zhang R, Ma Q, Han B, Xiang Y, Che J, Cao H, Fei X *et al.* 2010 The angiotensin-converting enzyme 2 in tumor growth and tumor-associated angiogenesis in non-small cell lung cancer. *Oncology Reports* **23** 941–948.
- Fischer M, Baessler A & Schunkert H 2002 Renin angiotensin system and gender differences in the cardiovascular system. *Cardiovascular Research* **53** 672–677. (doi:10.1016/S0008-6363(01)00479-5)
- Fleming I, Kohlstedt K & Busse R 2006 The tissue renin-angiotensin system and intracellular signalling. *Current Opinion in Nephrology and Hypertension* **15** 8–13. (doi:10.1097/01.mnh.0000196146.65330.ea)
- Ford CM, Li S & Pickering JG 1999 Angiotensin II stimulates collagen synthesis in human vascular smooth muscle cells. Involvement of the AT(1) receptor, transforming growth factor-beta, and tyrosine phosphorylation. *Arteriosclerosis, Thrombosis, and Vascular Biology* **19** 1843–1851. (doi:10.1161/01.ATV.19.8.1843)
- Fryzek JP, Poulsen AH, Lipworth L, Pedersen L, Norgaard M, McLaughlin JK & Friis S 2006 A cohort study of antihypertensive medication use and breast cancer among Danish women. *Breast Cancer Research and Treatment* **97** 231–236. (doi:10.1007/s10549-005-9091-x)
- Fujimoto Y, Sasaki T, Tsuchida A & Chayama K 2001 Angiotensin II type 1 receptor expression in human pancreatic cancer and growth inhibition by angiotensin II type 1 receptor antagonist. *FEBS Letters* **495** 197–200. (doi:10.1016/S0014-5793(01)02377-8)
- Fujita M, Hayashi I, Yamashina S, Itoman M & Majima M 2002 Blockade of angiotensin AT1a receptor signaling reduces tumor growth, angiogenesis, and metastasis. *Biochemical and Biophysical Research Communications* **294** 441–447. (doi:10.1016/S0006-291X(02)00496-5)
- Fujita M, Hayashi I, Yamashina S, Fukamizu A, Itoman M & Majima M 2005 Angiotensin type 1a receptor signaling-dependent induction of vascular endothelial growth factor in stroma is relevant to tumor-associated angiogenesis and tumor growth. *Carcinogenesis* **26** 271–279. (doi:10.1093/carcin/bgh324)
- Gack S, Vallon R, Schaper J, Ruther U & Angel P 1994 Phenotypic alterations in fos-transgenic mice correlate with changes in Fos/Jun-dependent collagenase type I

- expression. Regulation of mouse metalloproteinases by carcinogens, tumor promoters, cAMP, and Fos oncoprotein. *Journal of Biological Chemistry* **269** 10363–10369.
- Galis ZS & Khatri JJ 2002 Matrix metalloproteinases in vascular remodeling and atherogenesis: the good, the bad, and the ugly. *Circulation Research* **90** 251–262.
- Gallagher PE, Cook K, Soto-Pantoja D, Menon J & Tallant EA 2011 Angiotensin peptides and lung cancer. *Current Cancer Drug Targets* **11** 394–404. (doi:10.2174/156800911795538048)
- Ganong WF 1995 Reproduction and the renin–angiotensin system. *Neuroscience and Biobehavioral Reviews* **19** 241–250. (doi:10.1016/0149-7634(94)00056-7)
- de Gasparo M 2002 AT(1) and AT(2) angiotensin II receptors: key features. *Drugs* **62** 1–10. (doi:10.2165/00003495-200262991-00001)
- de Gasparo M, Catt KJ, Inagami T, Wright JW & Unger T 2000 International union of pharmacology, XXIII. The angiotensin II receptors. *Pharmacological Reviews* **52** 415–472.
- George AJ, Thomas WG & Hannan RD 2010 The renin–angiotensin system and cancer: old dog, new tricks. *Nature Reviews. Cancer* **10** 745–759. (doi:10.1038/nrc2945)
- Goffin V, Binart N, Touraine P & Kelly PA 2002 Prolactin: the new biology of an old hormone. *Annual Review of Physiology* **64** 47–67. (doi:10.1146/annurev.physiol.64.081501.131049)
- Goldberg JA, Kerr DJ, Wilmott N, McKillop JH & McArdle CS 1990 Regional chemotherapy for colorectal liver metastases: a phase II evaluation of targeted hepatic arterial 5-fluorouracil for colorectal liver metastases. *British Journal of Surgery* **77** 1238–1240. (doi:10.1002/bjs.1800771114)
- Gonzalez-Perez A, Ronquist G & Rodriguez LAG 2004 Breast cancer incidence and use of antihypertensive medication in women. *Pharmacoepidemiology and Drug Safety* **13** 581–585. (doi:10.1002/pds.910)
- Gonzalez-Zuloeta Ladd AM, Arias Vasquez A, Sayed-Tabatabaei FA, Coebergh JW, Hofman A, Njajou O, Stricker B & van Duijn C 2005 Angiotensin-converting enzyme gene insertion/deletion polymorphism and breast cancer risk. *Cancer Epidemiology, Biomarkers & Prevention* **14** 2143–2146. (doi:10.1158/1055-9965.EPI-05-0045)
- Gonzalez-Zuloeta Ladd AM, Arias Vasquez A, Siemes C, Yazdanpanah M, Coebergh JW, Hofman A, Stricker BH & van Duijn CM 2007 Differential roles of angiotensinogen and angiotensin receptor type I polymorphisms in breast cancer risk. *Breast Cancer Research and Treatment* **101** 299–304. (doi:10.1007/s10549-006-9290-0)
- Gordon MS, Chin WW & Shupnik MA 1992 Regulation of angiotensinogen gene-expression by estrogen. *Journal of Hypertension* **10** 361–366. (doi:10.1097/00004872-199204000-00007)
- Greco S, Elia MG, Muscella A, Storelli C & Marsigliante S 2002a AT1 angiotensin II receptor mediates intracellular calcium mobilization in normal and cancerous breast cells in primary culture. *Cell Calcium* **32** 1–10. (doi:10.1016/S0143-4160(02)00077-5)
- Greco S, Muscella A, Elia MG, Salvatore P, Storelli C & Marsigliante S 2002b Activation of angiotensin II type I receptor promotes protein kinase C translocation and cell proliferation in human cultured breast epithelial cells. *Journal of Endocrinology* **174** 205–214. (doi:10.1677/joe.0.1740205)
- Greco S, Muscella A, Elia MG, Salvatore P, Storelli C, Mazzotta A, Manca C & Marsigliante S 2003 Angiotensin II activates extracellular signal regulated kinases via protein kinase C and epidermal growth factor receptor in breast cancer cells. *Journal of Cellular Physiology* **196** 370–377. (doi:10.1002/jcp.10313)
- Green KA & Streuli CH 2004 Apoptosis regulation in the mammary gland. *Cellular and Molecular Life Sciences* **61** 1867–1883. (doi:10.1007/s00018-004-3366-y)
- Groves AM, Shastry M, Rodriguez-Justo M, Malhotra A, Endozo R, Davidson T, Kelleher T, Miles KA, Ell PJ & Keshtgar MR 2011 (1)F-FDGPET and biomarkers for tumour angiogenesis in early breast cancer. *European Journal of Nuclear Medicine and Molecular Imaging* **38** 46–52. (doi:10.1007/s00259-010-1590-2)
- Gupta P, Francosaenz R & Mulrow PJ 1995 Locally generated angiotensin-II in the adrenal-gland regulates basal, corticotropin-stimulated, and potassium-stimulated aldosterone secretion. *Hypertension* **25** 443–448.
- Haagensen C 1986 Diseases of the breast. Philadelphia, PA, USA: Saunders.
- Hagemann A, Nielsen AH & Poulsen K 1994 The uteroplacental renin–angiotensin system – a review. *Experimental and Clinical Endocrinology* **102** 252–261. (doi:10.1055/s-0029-1211289)
- Haiman CA, Henderson SO, Bretsky P, Kolonel LN & Henderson BE 2003 Genetic variation in angiotensin I-converting enzyme (ACE) and breast cancer risk: the multiethnic cohort. *Cancer Research* **63** 6984–6987.
- Han HJ, Han JY, Heo JS, Lee SH, Lee MY & Kim YH 2007 ANGII-stimulated DNA synthesis is mediated by ANG II receptor-dependent Ca(2+)/PKC as well as EGF receptor-dependent PI3K/Akt/mTOR/p70S6K1 signal pathways in mouse embryonic stem cells. *Journal of Cellular Physiology* **211** 618–629. (doi:10.1002/jcp.20967)
- Hansen RK & Bissell MJ 2000 Tissue architecture and breast cancer: the role of extracellular matrix and steroid hormones. *Endocrine-Related Cancer* **7** 95–113. (doi:10.1677/erc.0.0070095)
- Heffelfinger SC 2007 The renin angiotensin system in the regulation of angiogenesis. *Current Pharmaceutical Design* **13** 1215–1229. (doi:10.2174/138161207780618858)
- Herr D, Rodewald M, Fraser HM, Hack G, Konrad R, Kreienberg R & Wulff C 2008 Potential role of renin–angiotensin-system for tumor angiogenesis in receptor negative breast cancer. *Gynecological Oncology* **109** 418–425. (doi:10.1016/j.ygyno.2008.02.019)

- Hjorth SA, Schambye HT, Greenlee WJ & Schwartz TW 1994 Identification of peptide binding residues in the extracellular domains of the AT1 receptor. *Journal of Biological Chemistry* **269** 30953–30959.
- Howell A & Dowsett M 2004 Endocrinology and hormone therapy in breast cancer – aromatase inhibitors versus antioestrogens. *Breast Cancer Research* **6** 269–274. (doi:10.1186/bcr945)
- Huang W, Wu YL, Zhong J, Jiang FX, Tian XL & Yu LF 2008 Angiotensin II type 1 receptor antagonist suppress angiogenesis and growth of gastric cancer xenografts. *Digestive Diseases and Sciences* **53** 1206–1210. (doi:10.1007/s10620-007-0009-9)
- Hynes NE & Lane HA 2005 ERBB receptors and cancer: the complexity of targeted inhibitors. *Nature Reviews. Cancer* **5** 341–354. (doi:10.1038/nrc1609)
- Imai N, Hashimoto T, Kihara M, Yoshida S, Kawana I, Yazawa T, Kitamura H & Umemura S 2007 Roles for host and tumor angiotensin II type 1 receptor in tumor growth and tumor-associated angiogenesis. *Laboratory Investigation* **87** 189–198. (doi:10.1038/labinvest.3700504)
- Inigo SD, Lopez-Jorge CE, Gomez-Casares MT, Castellano AL, Cabrera PM, Brito JL, Cabrera AS & Labarta TM 2009 Induction of apoptosis in leukemic cell lines treated with captopril, trandolapril and losartan: a new role in the treatment of leukaemia for these agents. *Leukemia Research* **33** 810–816. (doi:10.1016/j.leukres.2008.09.029)
- Ino K, Shibata K, Kajiyama H, Yamamoto E, Nagasaka T, Nawa A, Nomura S & Kikkawa F 2006 Angiotensin II type 1 receptor expression in ovarian cancer and its correlation with tumour angiogenesis and patient survival. *British Journal of Cancer* **94** 552–560. (doi:10.1038/sj.bjc.6602961)
- Ino K, Shibata K, Yamamoto E, Kajiyama H, Nawa A, Mabuchi Y, Yagi S, Minami S, Tanizaki Y, Kobayashi A *et al.* 2011 Role of the renin–angiotensin system in gynecologic cancers. *Current Cancer Drug Targets* **11** 405–411. (doi:10.2174/156800911795538057)
- Inwang ER, Puddefoot JR, Brown CL, Goode AW, Marsigliante S, Ho MM, Payne JG & Vinson GP 1997 Angiotensin II type 1 receptor expression in human breast tissues. *British Journal of Cancer* **75** 1279–1283. (doi:10.1038/bjc.1997.217)
- Johnston CI 1992 Franz Volhard Lecture. Renin–angiotensin system: a dual tissue and hormonal system for cardiovascular control. *Journal of Hypertension* **10** S13–S26.
- Jones KL & Buzdar AU 2004 A review of adjuvant hormonal therapy in breast cancer. *Endocrine-Related Cancer* **11** 391–406. (doi:10.1677/erc.1.00594)
- Kalluri R 2009 EMT: when epithelial cells decide to become mesenchymal-like cells. *Journal of Clinical Investigation* **119** 1417–1419. (doi:10.1172/JCI39675)
- Kalluri R & Weinberg RA 2009 The basics of epithelial–mesenchymal transition. *Journal of Clinical Investigation* **119** 1420–1428. (doi:10.1172/JCI39104)
- Kapas S, Hinson JP, Puddefoot JR, Ho MM & Vinson GP 1994 Internalization of the type-I angiotensin-II receptor (AT1) is required for protein-kinase-C activation but not for inositol trisphosphate release in the angiotensin-II stimulated rat adrenal zona glomerulosa cell. *Biochemical and Biophysical Research Communications* **204** 1292–1298. (doi:10.1006/bbrc.1994.2603)
- Karakulakis G, Papakonstantinou E, Aletras AJ, Tamm M & Roth M 2007 Cell type specific effect of hypoxia and PDGF-BB on extracellular matrix turnover and its consequences for lung remodelling. *Journal of Biological Chemistry* **282** 908–925. (doi:10.1074/jbc.M602178200)
- Kaschina E & Unger T 2003 Angiotensin AT1/AT2 receptors: regulation, signalling and function. *Blood Pressure* **12** 70–88. (doi:10.1080/08037050310001057)
- Kaschina E, Scholz H, Steckelings UM, Sommerfeld M, Kemnitz UR, Artuc M, Schmidt S & Unger T 2009 Transition from atherosclerosis to aortic aneurysm in humans coincides with an increased expression of RAS components. *Atherosclerosis* **205** 396–403. (doi:10.1016/j.atherosclerosis.2009.01.003)
- Kim IY, Jeong SJ, Kim ES, Kim SH & Moon A 2007 Type I collagen-induced pro-MMP-2 activation is differentially regulated by H-Ras and N-Ras in human breast epithelial cells. *Journal of Biochemistry and Molecular Biology* **40** 825–831. (doi:10.5483/BMBRep.2007.40.5.825)
- Kim J, Ahn S, Rajagopal K & Lefkowitz RJ 2009 Independent beta-arrestin2 and Gq/protein kinase C pathways for ERK stimulated by angiotensin type 1A receptors in vascular smooth muscle cells converge on transactivation of the epidermal growth factor receptor. *Journal of Biological Chemistry* **284** 11953–11962. (doi:10.1074/jbc.M808176200)
- Kisley LR, Sakai RR & Fluharty SJ 1999 Estrogen decreases hypothalamic angiotensin II AT(1) receptor binding and mRNA in the female rat. *Brain Research* **844** 34–42. (doi:10.1016/S0006-8993(99)01815-6)
- Klett C, Hellmann W, Hackenthal E & Ganten D 1993 Modulation of tissue angiotensinogen gene expression by glucocorticoids, estrogens, and androgens in SHR and WKY rats. *Clinical and Experimental Hypertension* **15** 683–708. (doi:10.3109/10641969309041637)
- van der Knaap R, Siemes C, Coebergh JW, van Duijn CM, Hofman A & Stricker BH 2008 Renin–angiotensin system inhibitors, angiotensin I-converting enzyme gene insertion/deletion polymorphism, and cancer: the Rotterdam Study. *Cancer* **112** 748–757. (doi:10.1002/encr.23215)
- Koh WP, Yuan JM, Sun CL, van den Berg D, Seow A, Lee HP & Yu MC 2003 Angiotensin I-converting enzyme (ACE) gene polymorphism and breast cancer risk among Chinese women in Singapore. *Cancer Research* **63** 573–578.
- Koh WP, Yuan JM, Van Den Berg D, Lee HP & Yu MC 2005 Polymorphisms in angiotensin II type 1 receptor and angiotensin I-converting enzyme genes and breast cancer risk among Chinese women in Singapore. *Carcinogenesis* **26** 459–464. (doi:10.1093/carcin/bgh309)

- Kosaka T, Miyajima A, Takayama E, Kikuchi E, Nakashima J, Ohigashi T, Asano T, Sakamoto M, Okita H, Murai M *et al.* 2007 Angiotensin II type 1 receptor antagonist as an angiogenic inhibitor in prostate cancer. *Prostate* **67** 41–49. (doi:10.1002/pros.20486)
- Krause S, Maffini MV, Soto AM & Sonnenschein C 2010 The microenvironment determines the breast cancer cells' phenotype: organization of MCF7 cells in 3D cultures. *BMC Cancer* **10** 263. (doi:10.1186/1471-2407-10-263)
- Kucerova D, Zelezna B, Sloncova E & Sovova V 1998 Angiotensin II receptors on colorectal carcinoma cells. *International Journal of Molecular Medicine* **2** 593–595.
- Lakhani SR & O'Hare MJ 2001 The mammary myoepithelial cell—Cinderella or ugly sister? *Breast Cancer Research* **3** 1–4. (doi:10.1186/bcr260)
- Lamote I, Meyer E, Massart-Leen AM & Burvenich C 2004 Sex steroids and growth factors in the regulation of mammary gland proliferation, differentiation, and involution. *Steroids* **69** 145–159. (doi:10.1016/j.steroids.2003.12.008)
- Lau ST & Leung PS 2011 Role of the RAS in pancreatic cancer. *Current Cancer Drug Targets* **11** 412–420. (doi:10.2174/156800911795538110)
- Le MT, Vanderheyden PM, Szaszak M, Hunyady L & Vauquelin G 2002 Angiotensin IV is a potent agonist for constitutive active human AT1 receptors. Distinct roles of the N- and C-terminal residues of angiotensin II during AT1 receptor activation. *Journal of Biological Chemistry* **277** 23107–23110. (doi:10.1074/jbc.C200201200)
- Lebeau A, Nerlich AG, Sauer U, Lichtinghagen R & Lohrs U 1999 Tissue distribution of major matrix metalloproteinases and their transcripts in human breast carcinomas. *Anticancer Research* **19** 4257–4264.
- Lees KR, MacFadyen RJ, Doig JK & Reid JL 1993 Role of angiotensin in the extravascular system. *Journal of Human Hypertension* **7** S7–S12.
- Leung PS & Carlsson PO 2001 Tissue renin–angiotensin system: its expression, localization, regulation and potential role in the pancreas. *Journal of Molecular Endocrinology* **26** 155–164. (doi:10.1677/jme.0.0260155)
- Leung PS & Carlsson PO 2005 Pancreatic islet renin angiotensin system – its novel roles in islet function and in diabetes mellitus. *Pancreas* **30** 293–298. (doi:10.1097/01.mpa.0000158028.76666.76)
- Leung PS, Chan HC, Fu LX, Zhou WL & Wong PY 1997 Angiotensin II receptors, AT1 and AT2 in the rat epididymis, Immunocytochemical and electrophysiological studies. *Biochimica et Biophysica Acta* **1357** 65–72. (doi:10.1016/S0167-4889(97)00015-3)
- Lever AF, Hole DJ, Gillis CR, McCallum IR, McInnes GT, MacKinnon PL, Meredith PA, Murray LS, Reid JL & Robertson JWK 1998 Do inhibitors of angiotensin-I-converting enzyme protect against risk of cancer? *Lancet* **352** 179–184. (doi:10.1016/S0140-6736(98)03228-0)
- Li CI, Malone KE, Weiss NS, Boudreau DM, Cushing-Haugen KL & Daling JR 2003 Relation between use of anti hypertensive medications and risk of breast carcinoma among women ages 65–79 years. *Cancer* **98** 1504–1513. (doi:10.1002/cncr.11663)
- Li YH, Jiao LH, Liu RH, Chen XL, Wang H & Wang WH 2004 Localization of angiotensin II in pig ovary and its effects on oocyte maturation *in vitro*. *Theriogenology* **61** 447–459. (doi:10.1016/S0093-691X(03)00246-2)
- Liebmann C 2011 EGF receptor activation by GPCRs: an universal pathway reveals different versions. *Molecular and Cellular Endocrinology* **331** 222–231. (doi:10.1016/j.mce.2010.04.008)
- Lim KT, Cosgrave N, Hill AD & Young LS 2006 Nongenomic oestrogen signalling in oestrogen receptor negative breast cancer cells: a role for the angiotensin II receptor AT1. *Breast Cancer Research* **8** R33. (doi:10.1186/bcr1509)
- Linz W, Scholkens BA & Ganten D 1989 Converting enzyme inhibition specifically prevents the development and induces regression of cardiac hypertrophy in rats. *Clinical and Experimental Hypertension* **11** 1325–1350. (doi:10.3109/10641968909038172)
- Louis SN, Chow L, Rezmann L, Krezel MA, Catt KJ, Tikellis C, Frauman AG & Louis WJ 2010 Expression and function of ATIP/MTUS1 in human prostate cancer cell lines. *Prostate* **70** 1563–1574. (doi:10.1002/pros.21192)
- Magness RR, Parker CR & Rosenfeld CR 1993 Systemic and uterine responses to chronic infusion of estradiol-17-beta. *American Journal of Physiology* **265** E690–E698.
- Magy L, Vincent F, Faure S, Messerli FH, Wang JG, Achard JM & Fournier A 2005 The renin-angiotensin systems: evolving pharmacological perspectives for cerebroprotection. *Current Pharmaceutical Design* **11** 3275–3291.
- Mahmood T, Djahanbakhch O, Burleigh DE, Puddefoot JR, O'Mahony OA & Vinson GP 2002 Effect of angiotensin II on ion transport across human Fallopian tube epithelial cells *in vitro*. *Reproduction* **124** 573–579. (doi:10.1530/rep.0.1240573)
- Mancina R, Susini T, Renzetti A, Forti G, Razzoli E, Serio M & Maggi M 1996 Sex steroid modulation of AT(2) receptors in human myometrium. *Journal of Clinical Endocrinology and Metabolism* **81** 1753–1757. (doi:10.1210/jc.81.5.1753)
- Marsigliante S, Resta L, Muscella A, Vinson GP, Marzullo A & Storelli C 1996 AT1 angiotensin II receptor subtype in the human larynx and squamous laryngeal carcinoma. *Cancer Letters* **110** 19–27. (doi:10.1016/S0304-3835(96)04449-7)
- McKinley MJ, Albiston AL, Allen AM, Mathai ML, May CN, McAllen RM, Oldfield BJ, Mendelsohn FA & Chai SY 2003 The brain renin–angiotensin system: location and physiological roles. *International Journal of Biochemistry & Cell Biology* **35** 901–918. (doi:10.1016/S1357-2725(02)00306-0)
- Meier CR, Derby LE, Jick SS & Jick H 2000 Angiotensin-converting enzyme inhibitors, calcium channel blockers, and breast cancer. *Archives of Internal Medicine* **160** 349–353. (doi:10.1001/archinte.160.3.349)

- Mendizabal-Ruiz AP, Morales J, Castro Martinez X, Gutierrez Rubio SA, Valdez L, Vasquez-Camacho JG, Sanchez Corona J & Moran Moguel MC 2011 RAS polymorphisms in cancerous and benign breast tissue. *Journal of Renin-Angiotensin-Aldosterone System* **12** 85–92. (doi:10.1177/1470320310383735)
- Menon J, Soto-Pantoja DR, Callahan MF, Cline JM, Ferrario CM, Tallant EA & Gallagher PE 2007 Angiotensin-(1–7) inhibits growth of human lung adenocarcinoma xenografts in nude mice through a reduction in cyclooxygenase-2. *Cancer Research* **67** 2809–2815. (doi:10.1158/0008-5472.CAN-06-3614)
- Millan MA, Carvallo P, Izumi S, Zemel S, Catt KJ & Aguilera G 1989 Novel sites of expression of functional angiotensin II receptors in the late gestation fetus. *Science* **244** 1340–1342. (doi:10.1126/science.2734613)
- Miyajima A, Kikuchi E, Kosaka T & Oya M 2009 Angiotensin II type 1 receptor antagonist as an angiogenic inhibitor in urogenital cancer. *Reviews on Recent Clinical Trials* **4** 75–78. (doi:10.2174/157488709788185996)
- Moreno M & Bataller R 2008 Cytokines and renin–angiotensin system signaling in hepatic fibrosis. *Clinics in Liver Disease* **12** 825–852. (doi:10.1016/j.cld.2008.07.013)
- Morini M, Mottolèse M, Ferrari N, Ghiorzo F, Buglioni S, Mortarini R, Noonan DM, Natali PG & Albini A 2000 The alpha 3 beta 1 integrin is associated with mammary carcinoma cell metastasis, invasion, and gelatinase B (MMP-9) activity. *International Journal of Cancer* **87** 336–342. (doi:10.1002/1097-0215(20000801)87:3<336::AID-IJC>3.0.CO;2-3)
- Motz WH, Scheler S & Strauer BE 1992 Medical repair of hypertensive left ventricular remodeling. *Journal of Cardiovascular Pharmacology* **20** (Suppl 1) S32–S36.
- Mulrow PJ 1998 Renin–angiotensin system in the adrenal. *Hormone and Metabolic Research* **30** 346–349. (doi:10.1055/s-2007-978896)
- Mulrow PJ 1999 Angiotensin II and aldosterone regulation. *Regulatory Peptides* **80** 27–32. (doi:10.1016/S0167-0115(99)00004-X)
- Muscella A, Greco S, Elia MG, Storelli C & Marsigliante S 2002 Angiotensin II stimulation of Na⁺/K⁺ATPase activity and cell growth by calcium-independent pathway in MCF-7 breast cancer cells. *Journal of Endocrinology* **173** 315–323. (doi:10.1677/joe.0.1730315)
- Muscella A, Greco S, Elia MG, Storelli C & Marsigliante S 2003 PKC-zeta is required for angiotensin II-induced activation of ERK and synthesis of C-FOS in MCF-7 cells. *Journal of Cellular Physiology* **197** 61–68. (doi:10.1002/jcp.10336)
- Muscella A, Storelli C & Marsigliante S 2005 Atypical PKC-zeta and PKC-iota mediate opposing effects on MCF-7 Na⁺/K⁺ATPase activity. *Journal of Cellular Physiology* **205** 278–285. (doi:10.1002/jcp.20396)
- Muscogiuri G, Chavez AO, Gastaldelli A, Perego L, Tripathy D, Saad MJ, Velloso L & Folli F 2008 The crosstalk between insulin and renin–angiotensin–aldosterone signaling systems and its effect on glucose metabolism and diabetes prevention. *Current Vascular Pharmacology* **6** 301–312. (doi:10.2174/157016108785909715)
- Nahta R & Esteva FJ 2007 Trastuzumab: triumphs and tribulations. *Oncogene* **26** 3637–3643. (doi:10.1038/sj.onc.1210379)
- Nahta R, Yu D, Hung MC, Hortobagyi GN & Esteva FJ 2006 Mechanisms of disease: understanding resistance to HER2-targeted therapy in human breast cancer. *Nature Clinical Practice. Oncology* **3** 269–280. (doi:10.1038/ncponc0509)
- Nakai Y, Isayama H, Ijichi H, Sasaki T, Sasahira N, Hirano K, Kogure H, Kawakubo K, Yagioka H, Yashima Y *et al.* 2010 Inhibition of renin–angiotensin system affects prognosis of advanced pancreatic cancer receiving gemcitabine. *British Journal of Cancer* **103** 1644–1648. (doi:10.1038/sj.bjc.6605955)
- Natarajan R, Gonzales N, Hornsby PJ & Nadler J 1992 Mechanism of angiotensin II-induced proliferation in bovine adrenocortical cells. *Endocrinology* **131** 1174–1180. (doi:10.1210/en.131.3.1174)
- Neo JH, Ager EI, Angus PW, Zhu J, Herath CB & Christophi C 2010 Changes in the renin angiotensin system during the development of colorectal cancer liver metastases. *BMC Cancer* **10** 134. (doi:10.1186/1471-2407-10-134)
- Nguyen G 2011 Renin, (pro)renin and receptor: an update. *Clinical Science* **120** 169–178. (doi:10.1042/CS20100432)
- Nguyen G & Contrepas A 2008 Physiology and pharmacology of the (pro)renin receptor. *Current Opinion in Pharmacology* **8** 127–132. (doi:10.1016/j.coph.2007.12.009)
- Nicolini A, Carpi A & Tarro G 2006 Biomolecular markers of breast cancer. *Frontiers in Bioscience* **11** 1818–1843. (doi:10.2741/1926)
- Nielsen AH, Hagemann A & Poulson K 1995 The tissue renin–angiotensin system in the female reproductive tissues. In *Tissue Renin–Angiotensin Systems*, pp 253–268. Eds AK Mukhopadhyay & MK Raizada. New York & London: Plenum.
- Noguchi S, Miyauchi K, Nishizawa Y, Sasaki Y, Imaoka S, Iwanaga T, Koyama H & Terasawa T 1988 Augmentation of anticancer effect with angiotensin II in intraarterial infusion chemotherapy for breast carcinoma. *Cancer* **62** 467–473. (doi:10.1002/1097-0142(19880801)62:3<467::AID-CNCR2820620304>3.0.CO;2-Y)
- Norris B, Gonzalez C, Concha J, Palacios S & Contreras G 1991 Stimulatory effect of angiotensin-II on electrolyte transport in canine tracheal epithelium. *General Pharmacology* **22** 527–531.
- Ohnuma Y, Toda M, Fujita M, Hosono K, Suzuki T, Ogawa Y, Amano H, Kitasato H, Hayakawa K & Majima M 2009 Blockade of an angiotensin type I receptor enhances effects of radiation on tumor growth and tumor-associated angiogenesis by reducing vascular endothelial growth factor expression. *Biomedicine & Pharmacotherapy* **63** 136–145. (doi:10.1016/j.biopha.2007.11.005)

- Okura T, Kitami Y, Wakamiya R, Marumoto K, Iwata T & Hiwada K 1992 Renal and extra-renal renin gene expression in spontaneously hypertensive rats. *Blood Pressure* **3** 6–11.
- Olivares-Reyes JA, Arellano-Plancarte A & Castillo-Hernandez JR 2009 Angiotensin II and the development of insulin resistance: implications for diabetes. *Molecular and Cellular Endocrinology* **302** 128–139. (doi:10.1016/j.mce.2008.12.011)
- Oliveira L, Costa-Neto CM, Nakaie CR, Schreier S, Shimuta SI & Paiva AC 2007 The angiotensin II AT1 receptor structure-activity correlations in the light of rhodopsin structure. *Physiological Reviews* **87** 565–592. (doi:10.1152/physrev.00040.2005)
- Otake AH, Mattar AL, Freitas HC, Machado CM, Nonogaki S, Fujihara CK, Zatz R & Chammas R 2009 Inhibition of angiotensin II receptor 1 limits tumor-associated angiogenesis and attenuates growth of murine melanoma. *Cancer Chemotherapy and Pharmacology*.
- Owonikoko TK, Fabucci ME, Brown PR, Nisar N, Hilton J, Mathews WB, Ravert HT, Rauseo P, Sandberg K, Dannals RF et al. 2004 *In vivo* investigation of estrogen regulation of adrenal and renal angiotensin (AT1) receptor expression by PET. *Journal of Nuclear Medicine* **45** 94–100.
- Pal SK & Pegram M 2005 Epidermal growth factor receptor and signal transduction: potential targets for anti-cancer therapy. *Anti-Cancer Drugs* **16** 483–494. (doi:10.1097/00001813-200506000-00003)
- Petty WJ, Miller AA, McCoy TP, Gallagher PE, Tallant EA & Torti FM 2009 Phase I and pharmacokinetic study of angiotensin-(1–7), an endogenous antiangiogenic hormone. *Clinical Cancer Research* **15** 7398–7404. (doi:10.1158/1078-0432.CCR-09-1957)
- Phillips MI, Speakman EA & Kimura B 1993 Levels of angiotensin and molecular biology of the tissue renin angiotensin systems. *Regulatory Peptides* **43** 1–20. (doi:10.1016/0167-0115(93)90403-U)
- del Pilar Carrera M, Ramirez-Exposito MJ, Mayas MD, Garcia MJ & Martinez-Martos JM 2010 Mammary renin–angiotensin system-regulating aminopeptidase activities are modified in rats with breast cancer. *Tumour Biology* **31** 583–588. (doi:10.1007/s13277-010-0072-2)
- Pollard JW 2001 Tumour–stromal interactions – transforming growth factor-beta isoforms and hepatocyte growth factor/scatter factor in mammary gland ductal morphogenesis. *Breast Cancer Research* **3** 230–237. (doi:10.1186/bcr301)
- Polyak K & Hu M 2005 Do myoepithelial cells hold the key for breast tumor progression? *Journal of Mammary Gland Biology and Neoplasia* **10** 231–247. (doi:10.1007/s10911-005-9584-6)
- Ptasinska-Wnuk D, Lawnicka H, Fryczak J, Kunert-Radek J & Pawlikowski M 2007 Angiotensin peptides regulate angiogenic activity in rat anterior pituitary tumour cell cultures. *Endokrynologia Polska* **58** 478–486.
- Pucell AG, Bumpus FM & Husain A 1987 Rat ovarian angiotensin-II receptors – characterization and coupling to estrogen secretion. *Journal of Biological Chemistry* **262** 7076–7080.
- Puddefoot JR, Udeozo UKI, Barker S & Vinson GP 2006 The role of angiotensin II in the regulation of breast cancer cell adhesion and invasion. *Endocrine-Related Cancer* **13** 895–903. (doi:10.1677/erc.1.01136)
- Quan A & Baum M 1996 Endogenous production of angiotensin-II modulates rat proximal tubule transport. *Journal of Clinical Investigation* **97** 2878–2882. (doi:10.1172/JCI118745)
- Reddy MK, Baskaran K & Molteni A 1995 Inhibitors of angiotensin-converting enzyme modulate mitosis and gene expression in pancreatic cancer cells. *Proceedings of the Society for Experimental Biology and Medicine* **210** 221–226.
- Redondo S, Ruiz E, Padilla E, Gordillo-Moscoso A & Tejerina T 2007 Role of TGF-beta1 in vascular smooth muscle cell apoptosis induced by angiotensin II. *European Journal of Pharmacology* **556** 36–44. (doi:10.1016/j.ejphar.2006.10.060)
- Redondo-Muller MA, Stevanovic-Walker M, Barker S, Puddefoot JR & Vinson GP 2008 Anti-cancer actions of a recombinant antibody (R6313/G2) against the angiotensin II AT1 receptor. *Endocrine-Related Cancer* **15** 277–288. (doi:10.1677/ERC-07-0068)
- Rhodes DR, Ateeq B, Cao Q, Tomlins SA, Mehra R, Laxman B, Kalyana-Sundaram S, Lonigro RJ, Helgeson BE, Bhojani MS et al. 2009 AGTR1 overexpression defines a subset of breast cancer and confers sensitivity to losartan, an AGTR1 antagonist. *PNAS* **106** 10284–10289. (doi:10.1073/pnas.0900351106)
- Rigat B, Hubert C, Alhenc-Gelas F, Cambien F, Corvol P & Soubrier F 1990 An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *Journal of Clinical Investigation* **86** 1343–1346. (doi:10.1172/JCI114844)
- Rivera E, Arrieta O, Guevara P, Duarte-Rojo A & Sotelo J 2001 AT1 receptor is present in glioma cells; its blockage reduces the growth of rat glioma. *British Journal of Cancer* **85** 1396–1399. (doi:10.1054/bjoc.2001.2102)
- Ronquist G, Rodriguez LA, Ruigomez A, Johansson S, Wallander MA, Frithz G & Svardsudd K 2004 Association between captopril, other antihypertensive drugs and risk of prostate cancer. *Prostate* **58** 50–56. (doi:10.1002/pros.10294)
- Rosenthal T & Gavras I 2009 Angiotensin inhibition and malignancies: a review. *Journal of Human Hypertension* **23** 623–635. (doi:10.1038/jhh.2009.21)
- Ross JS, Schenkein DP, Pietrusko R, Rolfe M, Linette GP, Stec J, Stagliano NE, Ginsburg GS, Symmans WF, Pusztai L et al. 2004 Targeted therapies for cancer 2004. *American Journal of Clinical Pathology* **122** 598–609. (doi:10.1309/5CWPU41AFR1VYM3F)

- Rudolph-Owen LA & Matrisian LM 1998 Matrix metalloproteinases in remodeling of the normal and neoplastic mammary gland. *Journal of Mammary Gland Biology and Neoplasia* **3** 177–189. (doi:10.1023/A:1018746923474)
- Russo J, Balogh GA, Chen JQ, Fernandez SV, Fernbaugh R, Heulings R, Mailo DA, Moral R, Russo PA, Sheriff F *et al.* 2006 The concept of stem cell in the mammary gland and its implication in morphogenesis, cancer and prevention. *Frontiers in Bioscience* **11** 151–172. (doi:10.2741/1788)
- Santos EL, Pesquero JB, Oliveira L, Paiva AC & Costa-Neto CM 2004 Mutagenesis of the AT1 receptor reveals different binding modes of angiotensin II and (Sar1)-angiotensin II. *Regulatory Peptides* **119** 183–188. (doi:10.1016/j.regpep.2004.02.009)
- Saridogan E, Djahanbakhch O, Puddefoot JR, Demetroulis C, Collingwood K, Mehta JG & Vinson GP 1996a Angiotensin II receptors and angiotensin II stimulation of ciliary activity in human fallopian tube. *Journal of Clinical Endocrinology and Metabolism* **81** 2719–2725. (doi:10.1210/jc.81.7.2719)
- Saridogan E, Djahanbakhch O, Puddefoot JR, Demetroulis C, Dawda R, Hall AJ & Vinson GP 1996b Type 1 angiotensin II receptors in human endometrium. *Molecular Human Reproduction* **2** 659–664. (doi:10.1093/molehr/2.9.659)
- Scamuffa N, Siegfried G, Bontemps Y, Ma L, Basak A, Cherel G, Calvo F, Seidah NG & Khatib AM 2008 Selective inhibition of proprotein convertases represses the metastatic potential of human colorectal tumor cells. *Journal of Clinical Investigation* **118** 352–363. (doi:10.1172/JCI32040)
- Schorb W, Conrad KM, Singer HA, Dostal DE & Baker KM 1995 Angiotensin-II is a potent stimulator of MAP-kinase activity in neonatal rat cardiac fibroblasts. *Journal of Molecular and Cellular Cardiology* **27** 1151–1160. (doi:10.1016/0022-2828(95)90051-9)
- Sealey JE, Itskovitzeldor J, Rubattu S, James GD, August P, Thaler I, Levrone J & Laragh JH 1994 Estradiol-related and progesterone-related increases in the renin–aldosterone system – studies during ovarian stimulation and early-pregnancy. *Journal of Clinical Endocrinology and Metabolism* **79** 258–264. (doi:10.1210/jc.79.1.258)
- Seltzer A, Pinto JEB, Viglione PN, Correa FMA, Libertun C, Tsutsumi K, Steele MK & Saavedra JM 1992 Estrogens regulate angiotensin-converting enzyme and angiotensin receptors in female rat anterior-pituitary. *Neuroendocrinology* **55** 460–467. (doi:10.1159/000126157)
- Shah BH, Yesilkaya A, Olivares-Reyes JA, Chen HD, Hunyady L & Catt KJ 2004 Differential pathways of angiotensin II-induced extracellularly regulated kinase 1/2 phosphorylation in specific cell types: role of heparin-binding epidermal growth factor. *Molecular Endocrinology* **18** 2035–2048. (doi:10.1210/me.2003-0476)
- Shah BH, Neithardt A, Chu DB, Shah FB & Catt KJ 2006 Role of EGF receptor transactivation in phosphoinositide 3-kinase-dependent activation of MAP kinase by GPCRs. *Journal of Cellular Physiology* **206** 47–57. (doi:10.1002/jcp.20423)
- Sharma S, Sharma MC & Sarkar C 2005 Morphology of angiogenesis in human cancer: a conceptual overview, histoprosthetic perspective and significance of neoangiogenesis. *Histopathology* **46** 481–489. (doi:10.1111/j.1365-2559.2005.02142.x)
- Siegfried G, Basak A, Cromlish JA, Benjannet S, Marcinkiewicz J, Chretien M, Seidah NG & Khatib AM 2003 The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. *Journal of Clinical Investigation* **111** 1723–1732.
- Sierra Diaz E, Sanchez Corona J, Rosales Gomez RC, Gutierrez Rubio SA, Vazquez Camacho JG, Solano Moreno H & Moran Moguel MC 2009 Angiotensin-converting enzyme insertion/deletion and angiotensin type 1 receptor A1166C polymorphisms as genetic risk factors in benign prostatic hyperplasia and prostate cancer. *Journal of Renin-Angiotensin-Aldosterone System* **10** 241–246. (doi:10.1177/1470320309352800)
- Singer CF, Hudelist G, Galid A & Kubista E 2003 Pharmacological modulation of local feedback mechanisms as a therapeutic approach in breast cancer treatment. *Drugs of Today* **39** 917–926. (doi:10.1358/dot.2003.39.12.799410)
- Sipahi I, Debanne SM, Rowland DY, Simon DI & Fang JC 2010 Angiotensin-receptor blockade and risk of cancer: meta-analysis of randomised controlled trials. *Lancet Oncology* **11** 627–636. (doi:10.1016/S1470-2045(10)70106-6)
- Small W Jr, Molteni A, Kim YT, Taylor JM, Chen Z & Ward WF 1997 Captopril modulates hormone receptor concentration and inhibits proliferation of human mammary ductal carcinoma cells in culture. *Breast Cancer Research and Treatment* **44** 217–224. (doi:10.1023/A:1005827119296)
- Smith NJ, Chan HW, Qian H, Bourne AM, Hannan KM, Warner FJ, Ritchie RH, Pearson RB, Hannan RD & Thomas WG 2011 Determination of the exact molecular requirements for type 1 angiotensin receptor epidermal growth factor receptor transactivation and cardiomyocyte hypertrophy. *Hypertension* **57** 973–980. (doi:10.1161/HYPERTENSIONAHA.110.166710)
- Soto-Pantoja DR, Menon J, Gallagher PE & Tallant EA 2009 Angiotensin-(1–7) inhibits tumor angiogenesis in human lung cancer xenografts with a reduction in vascular endothelial growth factor. *Molecular Cancer Therapeutics* **8** 1676–1683. (doi:10.1158/1535-7163.MCT-09-0161)
- Stylianopoulos T, Diop-Frimpong B, Munn LL & Jain RK 2010 Diffusion anisotropy in collagen gels and tumors: the effect of fiber network orientation. *Biophysical Journal* **99** 3119–3128. (doi:10.1016/j.bpj.2010.08.065)
- Suganuma T, Ino K, Shibata K, Kajiyama H, Nagasaka T, Mizutani S & Kikkawa F 2005 Functional expression of the angiotensin II type 1 receptor in human ovarian

- carcinoma cells and its blockade therapy resulting in suppression of tumor invasion, angiogenesis, and peritoneal dissemination. *Clinical Cancer Research* **11** 2686–2694. (doi:10.1158/1078-0432.CCR-04-1946)
- Sullivan JA, Rupnow HL, Cale JM, Magness RR & Bird IM 2005 Pregnancy and ovarian steroid regulation of angiotensin II type 1 and type 2 receptor expression in ovine uterine artery endothelium and vascular smooth muscle. *Endothelium* **12** 41–56. (doi:10.1080/10623320590933752)
- Sun SZ, Wang Y, Li Q, Tian YJ, Liu MH & Yu YH 2006 Effects of benazepril on renal function and kidney expression of matrix metalloproteinase-2 and tissue inhibitor of metalloproteinase-2 in diabetic rats. *Chinese Medical Journal* **119** 814–821.
- Tahmasebi M, Puddefoot JR, Inwang ER, Goode AW, Carpenter R & Vinson GP 1998 Transcription of the prorenin gene in normal and diseased breast. *European Journal of Cancer* **34** 1777–1782. (doi:10.1016/S0959-8049(98)00173-7)
- Tahmasebi M, Puddefoot JR, Inwang ER & Vinson GP 1999 The tissue renin–angiotensin system in human pancreas. *Journal of Endocrinology* **161** 317–322. (doi:10.1677/joe.0.1610317)
- Tahmasebi M, Barker S, Puddefoot JR & Vinson GP 2006 Localisation of renin–angiotensin system (RAS) components in breast. *British Journal of Cancer* **95** 67–74. (doi:10.1038/sj.bjc.6603213)
- Teo KK 2011 Effects of telmisartan, irbesartan, valsartan, candesartan, and losartan on cancers in 15 trials enrolling 138 769 individuals The ARB Trialists Collaboration. *Journal of Hypertension* **29** 623–635. (doi:10.1097/HJH.0b013e32834928ff)
- Thomas WG & Mendelsohn FAO 2003 Angiotensin receptors: form and function and distribution. *International Journal of Biochemistry & Cell Biology* **35** 774–779. (doi:10.1016/S1357-2725(02)00263-7)
- Tlsty TD & Coussens LM 2006 Tumor stroma and regulation of cancer development. *Annual Review of Pathology* **1** 119–150. (doi:10.1146/annurev.pathol.1.110304.100224)
- Tucker HA 2000 Hormones, mammary growth, and lactation: a 41-year perspective. *Journal of Dairy Science* **83** 874–884. (doi:10.3168/jds.S0022-0302(00)74951-4)
- Uemura H, Hasumi H, Kawahara T, Sugiura S, Miyoshi Y, Nakaigawa N, Teranishi J, Noguchi K, Ishiguro H & Kubota Y 2005a Pilot study of angiotensin II receptor blocker in advanced hormone-refractory prostate cancer. *International Journal of Clinical Oncology* **10** 405–410. (doi:10.1007/s10147-005-0520-y)
- Uemura H, Nakaigawa N, Ishiguro H & Kubota Y 2005b Antiproliferative efficacy of angiotensin II receptor blockers in prostate cancer. *Current Cancer Drug Targets* **5** 307–323. (doi:10.2174/1568009054629663)
- Uemura H, Ishiguro H & Kubota Y 2008 Pharmacology and new perspectives of angiotensin II receptor blocker in prostate cancer treatment. *International Journal of Urology* **15** 19–26. (doi:10.1111/j.1442-2042.2007.01937.x)
- Vila-Porcile E & Corvol P 1998 Angiotensinogen, prorenin, and renin are Co-localized in the secretory granules of all glandular cells of the rat anterior pituitary: an immunoultrastructural study. *Journal of Histochemistry and Cytochemistry* **46** 301–311. (doi:10.1177/002215549804600303)
- Villadsen R 2005 In search of a stem cell hierarchy in the human breast and its relevance to breast cancer evolution. *APMIS* **113** 903–921. (doi:10.1111/j.1600-0463.2005.apm_344.x)
- Vinson GP & Ho MM 1998 The adrenal renin/angiotensin system in the rat. *Hormone and Metabolic Research* **30** 355–359. (doi:10.1055/s-2007-978898)
- Vinson GP, Ho MM, Puddefoot JR, Teja R & Barker S 1994 Internalisation of the type I angiotensin II receptor (AT1) and angiotensin II function in the rat adrenal zona glomerulosa cell. *Journal of Endocrinology* **141** R5–R9. (doi:10.1677/joe.0.141R005)
- Vinson GP, Puddefoot JR, Ho MM, Barker S, Mehta J, Saridogan E & Djahanbakhch O 1995 Type 1 angiotensin II (AT1) receptors in sperm. *Journal of Endocrinology* **144** 369–378. (doi:10.1677/joe.0.1440369)
- Vinson GP, Saridogan E, Puddefoot JR & Djahanbakhch O 1997 Tissue renin–angiotensin systems and reproduction. *Human Reproduction* **12** 651–662. (doi:10.1093/humrep/12.4.651)
- Vinson GP, Barker S, Puddefoot JR & Tahmasebi M 2007 The renin–angiotensin system in the breast. In *Frontiers in Research of the Renin–Angiotensin System on Human Disease*, pp 135–153. Ed PS Leung. Dordrecht: Springer.
- Volpe M, Azizi M, Danser AH, Nguyen G & Ruilope LM 2011 Twisting arms to angiotensin receptor blockers/antagonists: the turn of cancer. *European Heart Journal* **32** 19–22. (doi:10.1093/eurheartj/ehq382)
- Volpert OV, Ward WF, Lingen MW, Chesler L, Solt DB, Johnson MD, Molteni A, Polverini PJ & Bouck NP 1996 Captopril inhibits angiogenesis and slows the growth of experimental tumors in rats. *Journal of Clinical Investigation* **98** 671–679. (doi:10.1172/JCI118838)
- Wang T & Giebisch G 1996 Effects of angiotensin II on electrolyte transport in the early and late distal tubule in rat kidney. *American Journal of Physiology* **271** F143–F149.
- Wang TH, Xiang QL, Chen JW, Pan H & Cui YH 2007 Raloxifene plus 17beta-estradiol inhibits proliferation of primary cultured vascular smooth muscle cells and human mammary endothelial cells via the janus kinase/signal transducer and activator of transcription3 cascade. *European Journal of Pharmacology* **561** 7–13. (doi:10.1016/j.ejphar.2007.01.026)
- Weber KT, Brilla CG & Janicki JS 1991 Signals for the remodeling of the cardiac interstitium in systemic hypertension. *Journal of Cardiovascular Pharmacology* **17** (Suppl 2) S14–S19. (doi:10.1097/00005344-199117002-00004)

- Weidner N 2004 The importance of tumor angiogenesis: the evidence continues to grow. *American Journal of Clinical Pathology* **122** 675–677. (doi:10.1309/KY6EH0LGY6D6PGP5)
- Werb Z, Ashkenas J, MacAuley A & Wiesen JF 1996 Extracellular matrix remodeling as a regulator of stromal–epithelial interactions during mammary gland development, involution and carcinogenesis. *Brazilian Journal of Medical and Biological Research* **29** 1087–1097.
- Wiseman BS & Werb Z 2002 Stromal effects on mammary gland development and breast cancer. *Science* **296** 1046–1049. (doi:10.1126/science.1067431)
- Wolf G & Neilson EG 1993 Angiotensin II as a renal growth factor. *Journal of the American Society of Nephrology* **3** 1531–1540.
- Wong PYD, Fu WO, Huang SJ & Law WK 1990 Effect of angiotensins on electrogenic anion transport in monolayer-cultures of rat epididymis. *Journal of Endocrinology* **125** 449–456. (doi:10.1677/joe.0.1250449)
- Xi B, Zeng T, Liu L, Liang Y, Liu W, Hu Y & Li J 2011 Association between polymorphisms of the renin–angiotensin system genes and breast cancer risk: a meta-analysis. *Breast Cancer Research and Treatment* **130** 561–568. (doi:10.1007/s10549-011-1602-3)
- Xiao F, Puddefoot JR, Barker S & Vinson GP 2008 Changes in angiotensin II type I receptor signalling pathways evoked by a monoclonal antibody raised to the N-terminus. *Journal of Endocrinology* **197** 25–33. (doi:10.1677/JOE-07-0498)
- Yamaue H, Tanimura H, Terashita S, Iwahashi M, Tani M, Tsunoda T, Tamai M & Mori K 1990 Clinical evaluation of chemotherapy under angiotensin II-induced hypertension in patients with advanced cancer. *Nihon Geka Hokan* **59** 302–309.
- Yang X, Zhu MJ, Sreejayan N, Ren J & Du M 2005 Angiotensin II promotes smooth muscle cell proliferation and migration through release of heparin-binding epidermal growth factor and activation of EGF-receptor pathway. *Molecular Cell* **20** 263–270. (doi:10.1016/j.molcel.2005.10.023)
- Yaren A, Turgut S, Kursunluoglu R, Oztop I, Turgut G, Degirmencioglu S, Kelten C & Erdem E 2007 Insertion/deletion polymorphism of the angiotensin I-converting enzyme gene in patients with breast cancer and effects on prognostic factors. *Journal of Investigative Medicine* **55** 255–261. (doi:10.2310/6650.2007.00006)
- Yasumatsu R, Nakashima T, Masuda M, Ito A, Kuratomi Y, Nakagawa T & Komune S 2004 Effects of the angiotensin-I converting enzyme inhibitor perindopril on tumor growth and angiogenesis in head and neck squamous cell carcinoma cells. *Journal of Cancer Research and Clinical Oncology* **130** 567–573. (doi:10.1007/s00432-004-0582-7)
- Yoshiji H, Kuriyama S, Noguchi R & Fukui H 2004 Angiotensin-I converting enzyme inhibitors as potential anti-angiogenic agents for cancer therapy. *Current Cancer Drug Targets* **4** 555–567. (doi:10.2174/1568009043332790)
- Zechmann CM, Woenne EC, Brix G, Radzwill N, Ilg M, Bachert P, Peschke P, Kirsch S, Kauczor HU, Delorme S *et al.* 2007 Impact of stroma on the growth, micro-circulation, and metabolism of experimental prostate tumors. *Neoplasia* **9** 57–67. (doi:10.1593/neo.06688)
- Zhang Z, Li M, Rayburn ER, Hill DL, Zhang RW & Wang H 2005 Oncogenes as novel targets for cancer therapy – (part I) – growth factors and protein tyrosine kinases. *American Journal of Pharmacogenomics* **5** 173–190. (doi:10.2165/00129785-200505030-00004)
- Zhao Y, Chen X, Cai L, Yang Y, Sui G & Fu S 2010 Angiotensin II/angiotensin II type I receptor (AT1R) signaling promotes MCF-7 breast cancer cells survival via PI3-kinase/Akt pathway. *Journal of Cellular Physiology* **225** 168–173. (doi:10.1002/jcp.22209)
- Zimmerman BG & Dunham EW 1997 Tissue renin–angiotensin system: a site of drug action? *Annual Review of Pharmacology and Toxicology* **37** 53–69. (doi:10.1146/annurev.pharmtox.37.1.53)

Received in final form 10 December 2011

Accepted 15 December 2011

Made available online as an Accepted Preprint
16 December 2011