Clinical and functional impact of TARBP2 over-expression in adrenocortical carcinoma

Stefano Caramuta^{1,3}, Linkiat Lee^{1,3}, Deniz M Özata^{1,3}, Pinar Akçakaya^{1,3}, Hong Xie^{1,3}, Anders Höög³, Jan Zedenius^{2,4}, Martin Bäckdahl^{2,4}, Catharina Larsson^{1,3} and Weng-Onn Lui^{1,3}

Departments of ¹Oncology-Pathology ²Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

³Cancer Center Karolinska (CCK), R8:04, Karolinska University Hospital-Solna, SE-17176 Stockholm, Sweden

⁴Department of Breast and Endocrine Surgery, Karolinska University Hospital, Stockholm, Sweden

Correspondence should be addressed to S Caramuta or W-O Lui
Email
stefano.caramuta@ki.se or
weng-onn.lui@ki.se

Abstract

Deregulation of microRNA (miRNA) expression in adrenocortical carcinomas (ACCs) has been documented to have diagnostic, prognostic, as well as functional implications. Here, we evaluated the mRNA expression of *DROSHA*, *DGCR8*, *DICER* (*DICER1*), *TARBP2*, and *PRKRA*, the core components in the miRNA biogenesis pathway, in a cohort of 73 adrenocortical tumors (including 43 adenomas and 30 carcinomas) and nine normal adrenal cortices using a RT-qPCR approach. Our results show a significant over-expression of *TARBP2*, *DICER*, and *DROSHA* in the carcinomas compared with adenomas or adrenal cortices ($P < 0.001$ for all comparisons). Using western blot and immunohistochemistry analyses, we confirmed the higher expression of TARBP2, DICER, and DROSHA at the protein level in carcinoma cases. Furthermore, we demonstrate that mRNA expression of *TARBP2*, but not *DICER* or *DROSHA*, is a strong molecular predictor to discriminate between adenomas and carcinomas. Functionally, we showed that inhibition of TARBP2 expression in human NCI-H295R ACC cells resulted in a decreased cell proliferation and induction of apoptosis. TARBP2 over-expression was not related to gene mutations; however, copy number gain of the *TARBP2* gene was observed in 57% of the carcinomas analyzed. In addition, we identified that *miR-195* and *miR-497* could directly regulate TARBP2 and DICER expression in ACC cells. This is the first study to demonstrate the deregulation of miRNA-processing factors in adrenocortical tumors and to show the clinical and biological impact of TARBP2 over-expression in this tumor type.

Key Words

- TARBP2
- over-expression
- adrenocortical cancer
- diagnostic
- function

Endocrine-Related Cancer
(2013) 20, 551–564

Introduction

Adrenocortical tumors are detected in up to 7% of the population and are often found as incidental tumors (Hammarstedt *et al.* 2010). The majority of these tumors are adenomas while only a small proportion are classified as adrenocortical carcinomas (ACCs) with an annual

incidence of two cases per million among adults worldwide (Schteingart 2007).

The prognosis for ACCs is poor with reported 5-year survival rates between 20 and 45%, and treatment alternatives are very limited (Allolio & Fassnacht 2006,

Bertherat *et al.* 2006). In addition, discrimination of ACCs from adrenocortical adenomas can be challenging. Generally, a tumor is unambiguously classified as ACC when it presents signs of invasion and/or distant metastases. In the absence of metastasis, the diagnosis relies on the histopathological evaluation, and the histological Weiss score classification is still the most commonly used one (Weiss *et al.* 1989). Given the limitation of conventional histopathology in adrenocortical tumor diagnostics, the identification of additional molecular markers with diagnostic and prognostic potential for clinical management of ACC is needed.

Previous reports identified over-expression of IGF2, loss-of-heterozygosity at 11p15 and 17p13, and increased MIB-1 proliferation index as molecular biomarkers to distinguish adrenocortical adenomas and carcinomas (Yano *et al.* 1989, Ilvesmaki *et al.* 1993, Gicquel *et al.* 1994, Stojadinovic *et al.* 2003). Several studies have shown the potential of mRNA and microRNA (miRNA) profiling to correctly classify adenomas or carcinomas and to identify subgroups of ACC patients with different survival and outcome (de Fraipont *et al.* 2005, Giordano *et al.* 2009, Laurell *et al.* 2009, de Reynies *et al.* 2009, Soon *et al.* 2009, Tombol *et al.* 2009, Özata *et al.* 2011, Patterson *et al.* 2011, Schmitz *et al.* 2011).

While the role of miRNAs in cancer development and progression is established, recent studies are also focusing on the deregulation of miRNA-processing factors, which are needed for miRNA maturation. This pathway has been shown to play an important role in tumor initiation and progression and to have a prognostic potential in several cancer types (Roh *et al.* 2005, Sugito *et al.* 2006, Chiosea *et al.* 2007, Merritt *et al.* 2008, Melo *et al.* 2009).

In this study, we evaluated the expression of the miRNA machinery components as potential diagnostic and prognostic markers in adrenocortical tumors. We further investigated the effect of TARBP2 deregulation on cell growth and apoptosis and explored possible mechanisms responsible for its deregulation in this tumor type.

Materials and methods

Clinical material

This study included 73 snap-frozen primary sporadic adrenocortical tumors from 72 patients and nine histopathologically verified normal adrenal cortices. All tissue samples were stored at -80°C until use. Representative

sections from all specimens were subjected to histopathological evaluation to confirm more than 80% tumor cells in the samples. Tumors were histopathologically classified according to the WHO classification (DeLellis *et al.* 2004). The diagnosis of ACC was based on vascular invasion, nuclear grade, mitotic index, tumor necrosis, invasion of surrounding organs, and/or presence of distant metastasis. Clinical details of tumor cases have been partially published in previous studies (Laurell *et al.* 2009, Özata *et al.* 2011) and are summarized in Table 1. All samples were collected with informed consent, and the study of tissue material was approved by the local ethics committee (Dnr 01-353,01-136).

Cell line

The ACC cell line NCI-H295R was purchased from the American Type Culture Collection (LGC Standards, Middlesex, UK) and maintained in culture as described previously (Özata *et al.* 2011). Authentication of the cell line was verified by short tandem repeat profiling, as described in our recent study (Özata *et al.* 2011).

Table 1 Clinical features for the adrenocortical tumor cases studied.

Clinical features	Adenomas	Carcinomas
No. of cases	43	29
Gender		
Male	12	13
Female	31	16
Age (years)		
Median	55	61
Min-max	16–81	28–84
Subtypes		
Cushing	13	NA
Aldosteronoma	16	NA
Non-hyperfunctioning	14	NA
Tumor size (cm) ^a		
Median	3.5	12
Min-max	0.9–6.5	7–21
Metastasis		
Yes	0	16
No	43	13
Follow-up (months)		
Median	39	42
Min-max	12–258	2–195
Follow-up (outcome)		
Alive	38	13
Dead of other causes ^b	5	4
DOD	0	12

NA, not available; DOD, dead of disease.

^aNo information was available for one of the carcinomas.

^bPatients who died for causes not related to the disease.

DNA and RNA extraction

Genomic DNA was extracted using DNeasy Blood and Tissue Kit (Qiagen) and total RNA was isolated using mirVana miRNA Isolation Kit (Applied Biosystems/Ambion, Austin, TX, USA). Quantification of isolated DNA or RNA was performed by NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA).

RT quantitative real-time PCR (RT-qPCR) analysis

cDNA was synthesized from 100 ng of total RNA using High Capacity cDNA RT Kit (Applied Biosystems), and the mRNA expression was quantified for *DICER* (*DICER1*) (ID_00998578_m1), *DROSHA* (ID_01095029_m1), *TARBP2* (ID_00998379_m1), *DGCR8* (ID_00987089_m1), *PRKRA* (ID_00269379_m1), *IGF2* (ID_00277496_s1), and *H19* (ID_00399293_g1) using a 7900HT Real-Time PCR System (Applied Biosystems). Expression of *18S* rRNA (ID_99999901_s1) was evaluated in parallel for normalization purpose. *18S* was chosen as a reference gene for its stable expression in all samples (Supplementary Figure S1, see section on supplementary data given at the end of this article). Expression levels of *miR-497* (ID_001043) and *miR-195* (ID_000494) were quantified in the NCI-H295R cells after transfection experiments to evaluate transfection efficiency. The expression of these two miRNAs has previously been evaluated in a subset of the adrenocortical tumors (Özata et al. 2011). Here, we analyzed the levels of *miR-195* and *miR-497* in additional five ACC cases included in this study. Normalization for miRNA expression was done using *RNU6B* (*RNU6-6P*) (ID_001093). All reactions were performed in triplicate, and relative expression levels were reported as $2^{-\Delta CT}$.

Western blot analysis

Whole cell lysates were prepared as described previously (Özata et al. 2011) for detection of miRNA machinery proteins using primary antibody anti-TARBP2 (sc-100909; Santa Cruz Biotechnology, Inc.) at 1:1000 dilution, anti-DICER (ab14601; Abcam, Cambridge, UK) at 1:200 dilution, or anti-DROSHA (ab12286; Abcam) at 1:400 dilution. Anti-GAPDH antibody (sc-47724; Santa Cruz Biotechnology, Inc.) diluted at 1:5000 was used for normalization. Protein levels were quantified on X-ray films from immunoblots using ImageJ software (<http://rsb.info.nih.gov/ij/>).

Immunohistochemistry

Immunohistochemistry of TARBP2 was performed in 34 adrenocortical tumors (17 adenomas and 17 ACCs) and

five normal adrenal tissues using anti-TARBP2 (ab72547, Abcam). The details are available in the [Supplementary Materials and methods](#), see section on [supplementary data](#) given at the end of this article. The TARBP2 immunostaining and subcellular localization were evaluated by a pathologist (A H). The cytoplasmic immunoreactivity was expressed in most tumor cells with the most obvious difference in the intensity of the immunoreactivity. Therefore, the clinical specimens were scored as having strong, moderate, or weak/negative TARBP2 immunoreactivity. The nuclear immunoreactivity, on the other hand, did not appear in all nuclei and the cases were classified as positive, mixed, or negative. In the adenomas, two different cell types were identified (oxyphilic and lipid-rich cells) and evaluated for both cytoplasmic and nuclear TARBP2 staining.

Validation of potential molecular biomarkers in independent cohorts

Expression data from two previously published microarray data sets were downloaded from ArrayExpress database. Giordano et al.'s study included a total of 55 adrenocortical tumors (22 adenomas and 33 carcinomas) and ten adrenal cortices (<http://www.ebi.ac.uk/arrayexpress/experiments/E-GEOD-10927>), while 92 adrenocortical tumors (58 adenomas and 34 carcinomas) were analyzed in de Reynies et al.'s study (<http://www.ebi.ac.uk/arrayexpress/experiments/E-TABM-311>). Expression values for *TARBP2* (HG-U133_Plus_2.0 probe set: 203677_s_at), *DICER* (213229_at, 206061_s_at), *DROSHA* (2218269_at), *IGF2* (202409_at, 202410_x_at, 210881_s_at), and *H19* (224646_x_at, 224997_x_at) were extracted from the microarray data. Geometrical mean was used to obtain a single expression value for genes that were represented by more than one probe set on the Affymetrix chip. Comparison of mRNA expression among sample groups for each gene and evaluation of their predictive value in ACC classification was performed as described in the 'Statistical analyses' section.

Transfection experiments in NCI-H295R cells

NCI-H295R cells were transfected using Amaxa Nucleofector technology (Lonza, Basel, Switzerland) with pre-miR-195, pre-miR-497 (PM10827 and PM10490 respectively; Applied Biosystems/Ambion), or siTARBP2 (sc-106846; Santa Cruz Biotechnology, Inc.), as described previously (Özata et al. 2011). Pre-miR Negative control#1 (4464058; Applied Biosystems/Ambion) or siCTR (sc-36869; Santa Cruz Biotechnology, Inc.) were used as

negative controls. All transfection experiments were repeated at least three times.

WST-1 colorimetric assay

Cell viability was evaluated using WST-1 colorimetric assay (Roche Applied Science), as described previously (Özata *et al.* 2011). All experiments were conducted in eight wells for each condition and replicated at least three times independently. Cell viability was calculated by comparing the absorbance values of the samples after background subtraction and normalized to the siCTR-treated cells.

Apoptosis caspase-3 colorimetric assay

After 72 h of transfection, the effect on apoptosis was evaluated using caspase-3 colorimetric assay (Genscript, Piscataway, NJ, USA), as described previously (Özata *et al.* 2011). Relative caspase-3 activity was calculated by comparing the absorbance values of the siTARBP2-treated cells with the respective siCTR-treated cells. All experiments were replicated three times.

Mutation analysis

All coding exons and flanking exon–intron junctions of the *TARBP2* gene (NM_134323) were sequenced in 23 ACC specimens. The PCR products were purified using ExoSAP-IT (USB Corporation/Affymetrix, Cleveland, OH, USA) and sequenced at the KIGene facility. Primer sequences and PCR conditions used for the analysis are detailed in [Supplementary Table S1](#), see section on [supplementary data](#) given at the end of this article.

TaqMan copy number assay

TaqMan copy number assay (Applied Biosystems) was used to evaluate the changes of *TARBP2* copy number in adrenal cortices and adrenocortical tumors. The target gene *TARBP2* (ID_02091089_cn) and the reference gene *RNaseP* (ID_4403326) were analyzed in parallel. The calculated relative copy numbers and the predicted copy numbers were estimated using CopyCaller software (Applied Biosystems). DNA from normal adrenal cortex was used as calibrator for the analysis. All reactions were performed in triplicate.

Argonaute 2 co-immunoprecipitation and analysis of argonaute 2-associated mRNAs

Cells (2×10^6 cells/dish) were transfected with pre-miR Negative control#1, pre-miR-195 or pre-miR-497, and

seeded in six tissue culture plates (10 cm). After 72 h of transfection, the cells were used for co-immunoprecipitation (co-IP) experiments using protein G Sepharose 4 Fast Flow beads (17-0618-01; GE Healthcare, Uppsala, Sweden) coated with mouse anti-human argonaute 2 (Ago2) antibody (ab57113; Abcam), as described previously (Xie *et al.* 2012). Ago2-bound RNA was extracted with TRIzol reagent (Invitrogen). *DICER* and *TARBP2* mRNA expression levels were measured by RT-qPCR and normalized to *miR-483-3p* for input and IP samples. This miRNA was chosen as an internal control due to its high abundance in the NCI-H295R cells. Enrichment of *DICER* and *TARBP2* mRNAs bound to Ago2 was calculated from the relative amount of mRNA detected in IP samples divided by the relative amount of mRNA in the corresponding input samples.

Statistical analyses

Statistica 8.0 (StatSoft, Inc., Tulsa, OK, USA) or MS Office Excel was used for statistical calculations. One-way ANOVA and unpaired Student's *t*-test were used to compare mRNA and protein expression levels among or between sample groups, and paired Student's *t*-test was performed to analyze transfection and Ago2-IP experiments. The patient population was divided into two groups based on high or low expression of *DICER*, *DROSHA*, *TARBP2*, *IGF2*, or *H19* according to the median expression levels among the adrenocortical tumors. Correlations between *TARBP2*, *DICER*, and *DROSHA* mRNA and protein expression levels were assessed by Pearson's correlation analyses and *P* values were estimated by permuting the samples 1000 times. The association between mRNA gene expression levels and clinical–histopathological parameters was analyzed using χ^2 test. Kaplan–Meier analysis was carried out to obtain survival curves. The survival curves of different patient groups were compared using log-rank test. Overall survival was the period from the time of diagnosis until the end of follow-up or death of the patient. For overall survival analysis, survival was censored if patients were still alive at the end of the follow-up. Disease-free survival was the time between the initial diagnosis and the end of follow-up or documented recurrence/death. For disease-free survival analysis, the data were censored if patients did not show any recurrence or died for other causes not related to disease. In the survival analyses, the recurrent ACC (Ca30) was not included. All the analyses were two tailed and *P* values <0.05 were considered significant.

Results

Deregulation of miRNA-processing factors in adrenocortical tumors and clinical associations

We analyzed the mRNA expression levels of five miRNA-processing genes (*DROSHA*, *DGCR8*, *DICER*, *TARBP2*, and *PRKRA*) in 73 adrenocortical tumors and nine adrenal cortices using RT-qPCR. In comparison to adrenal cortices, 29 of 30 ACCs (96%) showed a significantly increased expression of *TARBP2* ($P=0.0001$). Twenty-six carcinomas (87%) also exhibited significant over-expression of *DICER* ($P=0.024$) and 24 (80%) of the carcinomas presented a significantly increased *DROSHA* expression ($P=0.03$) (Fig. 1A). By contrast, abnormal mRNA expression of *TARBP2*, *DICER*, and *DROSHA* was less frequent in adenomas (*TARBP2*, 32% (14/43); *DICER*, 37% (16/43); and *DROSHA*, 40% (17/43)) (Fig. 1A). Significant over-expression of *TARBP2* (1.3- to 4-fold, $P<0.0001$), *DICER* (0.4- to 5-fold, $P<0.0001$), and *DROSHA* (0.2- to 4.5-fold, $P<0.0001$) was also observed in carcinomas when compared with adenomas (Fig. 1A). However, we did not observe any significant differences for *DGCR8* or *PRKRA* between the three sample groups (Supplementary Figure S2, see section on supplementary data given at the end of this article).

The adenoma samples included in the study consisted of 13 Cushing, 16 aldosteronoma, and 14 non-hyperfunctional tumor samples. Comparing the mRNA expression levels of the five miRNA machinery genes analyzed, we did not detect any significant differences between the three adenoma subgroups (Supplementary Figure S3). Moreover, for carcinoma cases, we did not observe any significant association between *DROSHA*, *DGCR8*, *DICER*, *TARBP2*, or *PRKRA* mRNA levels and clinical-histopathological parameters such as gender, age, tumor size, or presence of metastasis (data not shown).

To establish whether the increased mRNA expressions were reflected on the protein level, *TARBP2*, *DICER*, and *DROSHA* were examined by western blot analysis in a subset of nine adrenal cortices, 23 adenomas, and 19 carcinomas. Concordantly with gene expression results, carcinoma cases showed a significant increased expression of the three proteins analyzed when compared with adenomas and adrenal cortices ($P<0.01$ for all comparisons) (Fig. 1B and Supplementary Figure S4, S5, and S6). In addition, Pearson's correlation analysis showed a significant correlation between western blot and RT-qPCR results for *TARBP2* ($Cor=0.75$, $P<0.001$), *DICER* ($Cor=0.67$, $P<0.001$), and *DROSHA* ($Cor=0.5$, $P<0.01$) (Fig. 1C).

Sixteen of the ACC cases included in the study were diagnosed with distant metastases (seven at time of diagnosis and nine during follow-up). Among the ACCs, univariate analysis identified presence of distant metastasis as a predictive factor of shorter overall survival (log-rank, $P=0.014$) and disease-free survival (log-rank, $P=0.001$) (Supplementary Figure S7). However, we did not find any significant association between size, gender, age, or expression levels of miRNA-processing factors with survival among the ACC patients (data not shown).

Immunohistochemical analysis of TARBP2 expression

Based on the RT-qPCR and western blot analyses, *TARBP2* showed the strongest upregulation in ACC. To further confirm this over-expression, we evaluated the protein level of *TARBP2* in a subset of adrenocortical tumors (17 adenomas and 17 carcinomas) and five normal adrenal glands using immunohistochemistry. The immunostaining results are summarized in Supplementary Table S2 and exemplified in Fig. 2.

RNA interference (RNAi) activity mainly occurs in the cytoplasm, and the essential RNAi-processing factors, such as *DICER*, *TARBP2*, and argonautes, are usually localized in the cytoplasm (Chendrimada et al. 2005). Interestingly, Kim et al. (2006) reported that *TARBP2* is required for siRNA-mediated transcriptional silencing in human cells, suggesting its role in the nuclear compartment. We therefore sought to determine subcellular localization and expression of *TARBP2* in adrenocortical tumors. Our results showed that *TARBP2* was mainly expressed in the cytoplasm, although a weaker nuclear staining was also observed. In the normal adrenal tissues, positive cytoplasmic staining for *TARBP2* was present in cells of the zona glomerulosa while cells of zona fasciculata and reticularis showed weak/negative cytoplasmic staining (Fig. 2A). In the adenomas, *TARBP2* staining was mainly confined to oxyphilic cells while the preponderance of lipid-rich cells was negative for *TARBP2* immunoreactivity. Twelve of the 17 adenomas (70%) showed a moderate cytoplasmic staining, four cases (24%) demonstrated weak or negative *TARBP2* expression while only one case presented strong *TARBP2* immunoreactivity (Fig. 2B and Supplementary Table S2). Concordantly with the results obtained by RT-qPCR and western blot, most of the ACCs (14/17, 82%) showed strong cytoplasmic *TARBP2* staining in nearly all tumor cells, while moderate *TARBP2* expression was detected in only three ACCs (18%) (Fig. 2C and Supplementary Table S2).

Figure 1

Relative expression levels of miRNA-processing machinery components in adrenocortical tumors and adrenal cortices. (A) Box plots show relative mRNA expression levels for *TARBP2*, *DICER*, and *DROSHA* in nine normal adrenal cortices, 43 adenomas, and 30 carcinomas measured by RT-qPCR. *18S* was used as a reference gene. The expression levels of mRNA between groups were compared using one-way ANOVA models and $P < 0.05$ was

considered significant. (B) Comparison of *TARBP2*, *DICER*, and *DROSHA* protein expression levels among different groups. GAPDH was used for normalization. P values were calculated using one-way ANOVA models and $P < 0.05$ was considered significant. (C) Scatter plots showing correlation between mRNA and protein levels for *TARBP2*, *DICER*, and *DROSHA* as assessed by Pearson's correlation analysis.

At nuclear level, among the five normal adrenal cortices, only a single case showed a mixed nuclear staining in the cells of the zona glomerulosa, while all the other samples exhibited negative *TARBP2* nuclear immunoreactivity. Among the adenomas, most of the cases (12/17) presented mixed or positive nuclear *TARBP2* expression in oxyphilic cells while the remaining five cases

showed negative staining in the nuclei of the same cell type. On the other hand, lipid-rich cells had mixed or positive nuclear staining in five adenomas and negative in 12 adenomas. The majority of ACC tumor cells were positive for *TARBP2* nuclear expression (10/17), while five showed mixed nuclear staining and two cases were negative (Fig. 2 and Supplementary Table S2). However,

Figure 2

Immunohistochemical detection of TARBP2 expression in normal adrenal glands (A, $\times 40$), adrenocortical adenomas (B, $\times 160$), and adrenocortical carcinomas (C, $\times 160$). (A) In the normal adrenal gland, TARBP2 staining is mainly present in the zona glomerulosa and, at lower intensity, in the zona fasciculata and reticularis (inserts A₁ and A₂, $\times 160$). (B) The adenomas show a moderate (insert B₁, $\times 400$) or weak/negative (insert B₂, $\times 400$) TARBP2 expression in the cytoplasm of oxyphilic cells while lipid-rich cells

only demonstrate weak/negative cytoplasmic TARBP2 staining. Most of the adenomas showed positive or mixed nuclear TARBP2 immunoreactivity among the oxyphilic cells and negative TARBP2 expression in the nuclei of lipid-rich cells. (C) The majority of carcinomas show a strong TARBP2 expression in the cytoplasm and mainly positive (insert C₁, $\times 400$) or mixed (insert C₂, $\times 400$) nuclear staining.

positive, mixed, or negative TARBP2 nuclear expression did not show any significant correlation with histopathological parameters and survival among the ACC cases (data not shown).

TARBP2 as a novel molecular predictor of carcinoma

Given the observed over-expression of *TARBP2*, *DICER*, and *DROSHA* in ACCs, we sought to investigate whether the ACCs could be accurately classified based on the expression levels of these three genes. Adrenocortical tumor patients were divided into high or low mRNA levels of *TARBP2*, *DICER*, and *DROSHA* according to their median expression levels. Among the three genes analyzed, *TARBP2* showed the highest sensitivity and specificity to discriminate between ACCs and adenomas. Twenty-seven patients with ACCs were correctly predicted based on the expression of *TARBP2* mRNA (27/29 patients, 93% sensitivity); however, 10/43 adenoma cases were misclassified, leading to a specificity of 76%. Classification of ACCs based on *DICER* or *DROSHA* expression levels resulted in a much lower sensitivity and specificity (76 and 67% respectively).

Previous studies have clearly shown over-expression of IGF2 (Gicquel *et al.* 2001, Giordano *et al.* 2009, Laurell *et al.* 2009) or decreased expression of *H19* (Gao *et al.* 2002,

Giordano *et al.* 2009) in ACCs and demonstrated the potential of IGF2 as a biomarker for ACC diagnosis (Gicquel *et al.* 2001). We therefore compared the predictive values of *IGF2*, *H19*, *TARBP2*, *DICER*, and *DROSHA* for detection of ACC. As shown in Supplementary Table S3, see section on supplementary data given at the end of this article, the classification based on *H19* and *TARBP2* expression had similar levels of sensitivity (SEN=93% for both *H19* and *TARBP2*) and specificity (SPE=79% for both *H19* and *TARBP2*), while *IGF2* showed a slightly lower sensitivity (89%) and specificity (77%) when compared with *TARBP2*. The combination of *TARBP2*, *IGF2*, and *H19* expressions had an additive effect in the predictive value for ACC (SEN=97% and SPE=81%). On the other hand, both *DICER* and *DROSHA* had lower prediction power than the three genes described earlier. The sensitivity, specificity, positive predictive value, negative predictive value, and overall accuracy for the five genes analyzed are summarized in Supplementary Table S3.

To further investigate the diagnostic power of *TARBP2*, *DICER*, and *DROSHA* in adrenocortical tumors and overcome limitations of pathological assessment, we applied disease-free survival as the end point for diagnosing ACC, which is similar to the approach applied by de Reynies *et al.* (2009). Among the 72 tumors analyzed,

TARBP2 mRNA expression was a good predictor of ACC (log-rank, $P < 0.001$) and allowed an accurate classification of all seven cases with metastasis at the time of diagnosis. *DROSHA* could also significantly predict ACC cases (log-rank, $P = 0.038$), although two carcinoma cases presenting metastasis at diagnosis were misclassified. However, *DICER* did not show any statistical significance (Supplementary Figure S8). We also tested the predictive potential for *TARBP2* and *DROSHA* to detect ACCs among the 45 potentially malignant adrenocortical tumors after exclusion of tumors with sizes ≤ 3 cm and the seven patients presenting metastases at diagnosis. As shown by Kaplan–Meier analysis, *TARBP2* was still found highly predictive of ACC among those without metastases at diagnosis (log-rank, $P = 0.02$; Fig. 3) while *DROSHA* did not show any statistical significance (data not shown). The classification obtained according to *TARBP2* expression was at least as accurate as the prediction based on *IGF2* and *H19* expressions or the combination of the three genes (Fig. 3).

Validation in independent cohorts of adrenocortical tumors

To strengthen our findings, we studied independent cohorts of 147 adrenocortical tumors and ten normal adrenal cortices from two previously published microarray data sets (Giordano *et al.* 2009, de Reynies *et al.* 2009). Concordantly with our results, *TARBP2*, *DICER*, and *DROSHA* were significantly over-expressed in carcinomas when compared with adenomas or normal adrenal cortices in both studies (Supplementary Figure S9, see section on supplementary data given at the end of this article). Moreover, in both cohorts, we confirmed the higher predictive values for *TARBP2* when compared with *DICER* or *DROSHA* in the classification of ACC among adrenocortical tumors (Giordano *et al.*'s cohort: *TARBP2*, SEN=84%, SPE=100%; de Reynies *et al.*'s cohort: *TARBP2*, SEN=80%, SPE=82%). As observed in our cohort, the combination of *TARBP2*, *IGF2*, and *H19* could improve the overall prediction for ACC classification in both data sets (Supplementary Tables S4 and S5, see section on supplementary data given at the end of this article).

TARBP2 affects cell viability and apoptosis in NCI-H295R cells

The over-expression of *TARBP2* observed in ACC tumors prompted us to investigate the functional consequences of *TARBP2* alteration in human NCI-H295R ACC cells. Upon inhibition of *TARBP2* expression, ACC cells showed a

significant reduction of cell viability over time (Fig. 4A) and a markedly increase ($\sim 30\%$) of cell death (Fig. 4B) in comparison with negative control cells.

TARBP2 sequencing and copy number variations

We investigated possible mechanisms that could induce over-expression of *TARBP2* in ACCs. First, genomic DNA from 23 ACCs was screened for *TARBP2* gene mutations by sequencing. However, no mutations were identified in any of the coding exons analyzed (data not shown).

As the *TARBP2* gene is located in the chromosomal region 12q13.13 that is frequently gained/amplified in ACC (Stephan *et al.* 2008), we investigated *TARBP2* gene copy number alterations in a subset of 28 carcinomas, 18 adenomas, and 3 normal adrenal cortices using TaqMan copy number assay. We observed copy number gain in 57% of the carcinoma cases (16/28), but no changes among the adenomas and normal tissue samples (Supplementary Figure S10, see section on supplementary data given at the end of this article). However, the genomic copy number alteration did not always correspond with the *TARBP2* mRNA levels. For examples, four tumors (Ca3, Ca20, Ca22, and Ca25), showing *TARBP2* copy number gain, were among the cases with relatively low *TARBP2* expression (Supplementary Figure S10). This discrepancy implies that additional mechanism(s), e.g. posttranscriptional regulation by miRNAs, may involve in *TARBP2* regulation.

miR-195 and miR-497 regulate TARBP2 and DICER expression in adrenocortical tumors

To explore a possible involvement of miRNAs in *TARBP2* regulation, we first searched for the published under-expressed miRNAs in ACCs, thus showing an inverse expression pattern compared with *TARBP2*. We found 12 downregulated miRNAs that were reported in more than one study (Soon *et al.* 2009, Tombol *et al.* 2009, Özata *et al.* 2011, Patterson *et al.* 2011, Schmitz *et al.* 2011; Supplementary Table S6, see section on supplementary data given at the end of this article).

Next, we performed bioinformatic analysis using TargetScan (<http://www.targetscan.org/>) to identify whether any of the downregulated miRNAs have potential binding sites in the 3'-UTR of *TARBP2*. The analysis identified *let-7* family members, *miR-195* and *miR-497*, as possible regulators of *TARBP2*. In addition, potential target sites for these miRNAs were also present in the 3'-UTR of *DICER* (Fig. 5A). As *let-7* family members were not significantly deregulated in ACCs compared with

Figure 3

Prediction of carcinoma based on the mRNA expression levels of *TARBP2*, *IGF2*, and/or *H19*. The Kaplan–Meier curves separate the patients according to the expression levels of the molecular predictors *TARBP2*, *IGF2*, and/or *H19*. The 45 clinical cases were divided in two groups with high or low expression levels of *TARBP2*, *IGF2*, or *H19* according to their median

expression levels measured by RT-qPCR. A case was predicted as carcinoma if the expression level was >0.00012 for *TARBP2*, >0.00025 for *IGF2*, <0.0019 for *H19*, or >-0.0013 for combination of the three genes. The curves of each group were compared using log-rank test and $P<0.05$ was considered significant.

adenomas in our cohort (Özata et al. 2011), we focused on *miR-195* and *miR-497*. These miRNAs belong to the same miRNA cluster and show an inverse expression pattern in comparison with *TARBP2* and *DICER* expression levels (Fig. 5B). Noteworthy, the four carcinomas with *TARBP2* copy number gain and low *TARBP2* expression exhibited high expression of *miR-195* and *miR-497*. The same cases also presented low levels of *DICER* (Fig. 5C). These observations prompted us to investigate the role of *miR-195* and *miR-497* as potential regulators of *TARBP2* and/or *DICER*.

To confirm whether *TARBP2* and *DICER* were biological targets of *miR-195* and *miR-497* in adrenocortical tumors, we transfected NCI-H295R cells with miRNA mimics (pre-miR-195 and/or pre-miR-497). We observed

a significant reduction of *TARBP2* and *DICER* mRNA and protein expression levels in the cells over-expressing *miR-195* or *miR-497* compared with the cells transfected with pre-miR-negative control. *miR-497* had a stronger repressive effect on both *TARBP2* and *DICER*, and the combined over-expression of the two miRNAs showed a synergic effect on posttranscriptional regulation (Fig. 5D).

Next, we adopted the Ago2-immunoprecipitation approach to determine whether *TARBP2* and *DICER* are direct targets of *miR-195* and/or *miR-497*. Previous studies have shown that Ago complexes can stably associate with miRNA targets (Beitzinger et al. 2007, Karginov et al. 2007). Thus, we immunoprecipitated Ago2 complexes in NCI-H295R cells after induced over-expression of *miR-195* or *miR-497* and measured the abundance of the

Figure 4

Silencing of *TARBP2* in NCI-H295R cell line affects cell viability and apoptosis. (A) Histogram showing fractions of viable cells measured by WST-1 colorimetric assay in NCI-H295R cells transfected with siTARBP2 or siCTR at different time points (24, 48, and 72 h). (B) Histogram illustrating the increase of cell death evaluated by caspase-3/CPP32 colorimetric assay in NCI-H295R cells transfected with siTARBP2 when compared with siCTR-transfected cells. Error bars represent s.d. of the mean of three independent experiments. Paired t-test was used to determine the differences between groups and $P < 0.05$ were considered significant. (C) A representative western blot showing TARBP2 levels at 24, 48, and 72 h after transfection with siTARBP2 or siCTR. Incubation with a GAPDH antibody was used as a loading control.

co-immunoprecipitated *TARBP2* and *DICER* mRNAs by RT-qPCR. We observed a significant enrichment of endogenous mRNAs for both genes in cells over-expressing *miR-497* ($P < 0.05$). *TARBP2* mRNA was also strongly enriched in the immunoprecipitates of *miR-195* over-expressing cells ($P < 0.05$) while for *DICER* mRNA the enrichment did not show any statistical significance (Fig. 5E).

Discussion

We report dysregulation of miRNA machinery components in adrenocortical tumors and the potential role of *TARBP2* as molecular biomarker for ACC classification. In addition, we show the mechanisms of *TARBP2* regulation and the functional consequences of *TARBP2* deregulation in ACC cells.

Over-expression of *TARBP2*, *DICER*, and *DROSHA* in ACCs

We show that *TARBP2*, *DICER*, and *DROSHA* are significantly over-expressed in ACC when compared with adenomas and adrenal cortices. The expression pattern was validated in publicly available data sets. Disruption of miRNA machinery has been previously associated with tumorigenesis in several cancer types. In line with our findings, over-expression of *TARBP2* and *DICER* was observed in prostate cancer (Fu *et al.* 2010). *DICER* over-expression was also reported in cutaneous melanomas (Ma *et al.* 2011, Sand *et al.* 2011) and was associated with aggressiveness in lung adenocarcinoma (Chiose *et al.* 2007) and poor survival in colorectal cancer (Faber *et al.* 2011). Higher expression of *DROSHA* was found in cervical squamous cell carcinomas (Muralidhar *et al.* 2007) and epithelial skin cancers (Sand *et al.* 2010). Its over-expression was associated with poor prognosis in esophageal cancer (Sugito *et al.* 2006). On the contrary, reduced expression of miRNA-processing factors has also been observed in several tumor types. Decreased expression of *TARBP2* was reported in colorectal and gastric cancers presenting microsatellite instability (Melo *et al.* 2009). Low expression of *DROSHA* and *DICER* was associated with decreased survival in ovarian cancer (Merritt *et al.* 2008). Decreased mRNA level of *DICER* was observed in basal cell carcinomas (Sand *et al.* 2010) and was also associated with poor prognosis in non-small-cell lung cancer (Karube *et al.* 2005). Together, variation of *TARBP2*, *DICER*, and *DROSHA* expression levels among different tumor types suggests that deregulation of miRNA-processing factors can be dependent on cellular context and imply their possible dual role as tumor

Figure 5

miR-195 and *miR-497* target *TARBP2* and *DICER*. (A) Schematic presentation of *miR-195* and *miR-497* predicted target sites identified in the *TARBP2* and *DICER* 3'-UTRs by TargetScan 5.2. (B) *miR-195/miR-497* and *TARBP2/DICER* mRNA relative expression levels in adrenocortical carcinomas ($n = 30$) in comparison to adenomas ($n = 43$) assessed by RT-qPCR. The two miRNAs show an inverse pattern of expression compared with *TARBP2* and *DICER*. (C) Comparison between *miR-195/miR-497* and *TARBP2/DICER* mRNA expression levels in adrenocortical carcinoma cases measured by RT-qPCR. (D) Transfection of pre-miR-195 and/or pre-miR-497 in NCI-H295R cell line represses mRNA and protein expression of *TARBP2* and *DICER* when

compared with pre-miR-CTR. (E) NCI-H295R cells were transfected with pre-miR-CTR, pre-miR-195, or pre-miR-497. The levels of Ago2-associated *TARBP2* and *DICER* mRNAs were measured by RT-qPCR relative to *miR-483-3p*. The mRNA enrichment of *TARBP2* or *DICER* in each anti-Ago2 IP sample was normalized to the corresponding input sample. The results are shown as fold change in comparison to the negative control and the error bars represent s.d. of the mean of three independent experiments. Differences in expression levels between groups were evaluated using paired t-test and $P < 0.05$ was considered significant. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; NS, not significant.

suppressors or oncogenes in human cancers. Despite that we have not investigated whether deregulation of miRNA machinery genes affects the miRNA-processing efficiency in ACC cells, these effects have been demonstrated for TARBP2 and DICER in colorectal and breast cancer cells respectively (Melo *et al.* 2009, 2011, Martello *et al.* 2010).

Interestingly, we observed that TARBP2 was localized both in the cytoplasm and nucleus of tumor cells, whereas its expression was mainly found in the cytoplasm of non-tumor cells. In line with our findings, RNAi pathway components (including DICER, TARBP2, AGO1, and AGO2) have been observed in the nuclear compartment of human cells (Haussecker & Proudfoot 2005, Kim *et al.* 2006, Ahlenstiel *et al.* 2012). These observations suggest that TARBP2 may involve in nuclear RNAi on transcriptional silencing or other unknown function(s). Further analyses are warranted to investigate the potential nuclear role of TARBP2 and its implication in the pathogenesis of adrenocortical tumors.

TARBP2 as molecular predictor of carcinoma and functional role in ACC cells

Discrimination between localized ACCs and adenomas is challenging. To overcome these clinical limitations, the identification of novel molecular biomarkers is certainly needed. In our study, we show that *TARBP2* is a strong predictor of carcinoma and can reliably classify ACC in a cohort of non-metastatic adrenocortical tumors.

Consistent with the over-expression observed in ACC, we also demonstrated that inhibition of TARBP2 expression could affect cell growth and apoptosis in the NCI-H295R cells. In line with these results, previous studies demonstrated that TARBP2 promotes cell growth and transformation *in vitro* and can induce tumor formation in mice (Benkirane *et al.* 1997, Lee *et al.* 2004). These findings suggest a possible oncogenic function of TARBP2 in ACC carcinogenesis and imply the potential use of TARBP2 as a novel therapeutic target in ACC.

Regulatory mechanisms of TARBP2 over-expression in ACC

Despite several studies showing alteration of miRNA machinery factors in different tumor types, the molecular mechanisms that regulate their expression are not fully understood. Given the over-expression and the potential oncogenic role of TARBP2 in ACC, we sought to investigate possible molecular mechanisms involved in

the regulation of TARBP2 expression. *TARBP2* gene mutations causing a loss of TARBP protein expression have been previously shown in colorectal and gastric cancers (Melo *et al.* 2009). However, in our study, the mutational analysis revealed the presence of only wild-type sequences for the *TARBP2* gene in all the ACC cases analyzed, implying that genomic DNA mutations are probably not involved in the deregulation of TARBP2 in ACC. On the other hand, we found *TARBP2* gene copy number gain in 57% (16/28) of the carcinomas, suggesting that copy number gain of *TARBP2* gene may be, at least partially, responsible for its over-expression in ACC.

In addition, we propose a miRNA-mediated regulation of *TARBP2* expression in ACC. While there is no evidence in the literature of miRNAs affecting TARBP2 expression, *let-7* and *miR-103/-107* family are known to regulate expression of DICER (Forman *et al.* 2008, Martello *et al.* 2010). Here, we demonstrate that mature *miR-195* and *miR-497* can directly regulate *TARBP2* and *DICER* expression in ACC. Downregulation of *miR-195* and/or *miR-497* has been observed in several tumor types, including ACC (Soon *et al.* 2009, Özata *et al.* 2011, Patterson *et al.* 2011, Schmitz *et al.* 2011), liver (Xu *et al.* 2009), bladder (Han *et al.* 2011), breast (Li *et al.* 2011), and peritoneal carcinoma (Flavin *et al.* 2009). Reduced expression of *miR-195* is also correlated with lymph node metastasis and poor prognosis in colorectal cancer (Wang *et al.* 2011). In line with the expression pattern, we previously showed that over-expression of *miR-195* and *miR-497* can inhibit cell growth with concomitant increase of apoptosis in NCI-H295R ACC cells (Özata *et al.* 2011). Together, it is intriguing to speculate that the phenotypic effect observed in ACC cell line upon alteration of *miR-195* and *miR-497* expression may be mediated through TARBP2 and DICER downregulation.

In summary, we report frequent over-expression of TARBP2 in ACC and we found that *TARBP2* mRNA expression level is a useful predictor of ACC and able to discriminate adenomas from carcinomas. In addition, we revealed the potential oncogenic role of TARBP2 and mechanisms of its over-expression in ACC.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/ERC-13-0098>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by the Swedish Research Council (523-2009-3517 and 521-2010-3518), the Åke Olsson's Foundation for Haematological Research, the Cancer Research Funds of Radiumhemmet, the Axel and Signe Lagerman's Donation Foundation, the Swedish Cancer Society, Karolinska Institutet, and Stockholm County Council.

Author contribution statement

S Caramuta and W-O Lui conceived and designed the experiments; S Caramuta and L Lee developed the methodology; S Caramuta, L Lee, D M Özata, P Akçakaya, and H Xie performed the experiments; S Caramuta, L Lee, D M Özata, P Akçakaya, H Xie, A Höög, and W-O Lui analyzed and interpreted the data; S Caramuta, L Lee, D M Özata, P Akçakaya, H Xie, A Höög, J Zedenius, M Bäckdahl, C Larsson, and W-O Lui contributed to write and/or revise the manuscript; S Caramuta, L Lee, J Zedenius, M Bäckdahl, C Larsson, and W-O Lui contributed to administrative, technical, or material support; S Caramuta and W-O Lui supervised the study.

Acknowledgements

The authors thank Ms Lisa Ånfalk for her admirable help in collecting the tumor samples and the members of the Medical Genetics group for their help and valuable suggestions. We also thank Dr Daniela Witten (Department of Biostatistics, University of Washington) for statistical advice.

References

- Ahlenstiel CL, Lim HG, Cooper DA, Ishida T, Kelleher AD & Suzuki K 2012 Direct evidence of nuclear Argonaute distribution during transcriptional silencing links the actin cytoskeleton to nuclear RNAi machinery in human cells. *Nucleic Acids Research* **40** 1579–1595. (doi:10.1093/nar/gkr891)
- Allolio B & Fassnacht M 2006 Clinical review: adrenocortical carcinoma: clinical update. *Journal of Clinical Endocrinology and Metabolism* **91** 2027–2037. (doi:10.1210/jc.2005-2639)
- Beitzinger M, Peters L, Zhu JY, Kremmer E & Meister G 2007 Identification of human microRNA targets from isolated argonaute protein complexes. *RNA Biology* **4** 76–84. (doi:10.4161/rna.4.2.4640)
- Benkirane M, Neuveut C, Chun RF, Smith SM, Samuel CE, Gatignol A & Jeang KT 1997 Oncogenic potential of TAR RNA binding protein TRBP and its regulatory interaction with RNA-dependent protein kinase PKR. *EMBO Journal* **16** 611–624. (doi:10.1093/emboj/16.3.611)
- Bertherat J, Groussin L & Bertagna X 2006 Mechanisms of disease: adrenocortical tumors – molecular advances and clinical perspectives. *Nature Clinical Practice. Endocrinology & Metabolism* **2** 632–641. (doi:10.1038/ncpendmet0321)
- Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K & Shiekhattar R 2005 TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature* **436** 740–744. (doi:10.1038/nature03868)
- Chiosea S, Jeletzova E, Chandran U, Luo J, Mantha G, Sobol RW & Dacic S 2007 Overexpression of Dicer in precursor lesions of lung adenocarcinoma. *Cancer Research* **67** 2345–2350. (doi:10.1158/0008-5472.CAN-06-3533)
- DeLellis RA, Lloyd RV, Heitz PU & Eng C 2004 Tumors of the adrenal gland. In *World Health Organization Classification of Tumors. Pathology and Genetics of Endocrine Organs*, 135–146. Lyon, France: IARC Press.
- Faber C, Horst D, Hlubek F & Kirchner T 2011 Overexpression of Dicer predicts poor survival in colorectal cancer. *European Journal of Cancer* **47** 1414–1419. (doi:10.1016/j.ejca.2011.01.006)
- Flavin RJ, Smyth PC, Laios A, O'Toole SA, Barrett C, Finn SP, Russell S, Ring M, Denning KM, Li J et al. 2009 Potentially important microRNA cluster on chromosome 17p13.1 in primary peritoneal carcinoma. *Modern Pathology* **22** 197–205. (doi:10.1038/modpathol.2008.135)
- Forman JJ, Legesse-Miller A & Collier HA 2008 A search for conserved sequences in coding regions reveals that the let-7 microRNA targets Dicer within its coding sequence. *PNAS* **105** 14879–14884. (doi:10.1073/pnas.0803230105)
- de Fraipont F, El Atifi M, Cherradi N, Le Moigne G, Defaye G, Houlgatte R, Bertherat J, Bertagna X, Plouin PF, Baudin E et al. 2005 Gene expression profiling of human adrenocortical tumors using complementary deoxyribonucleic acid microarrays identifies several candidate genes as markers of malignancy. *Journal of Clinical Endocrinology and Metabolism* **90** 1819–1829. (doi:10.1210/jc.2004-1075)
- Fu X, Xue C, Huang Y, Xie Y & Li Y 2010 The activity and expression of microRNAs in prostate cancers. *Molecular BioSystems* **6** 2561–2572. (doi:10.1039/c0mb00100g)
- Gao ZH, Suppola S, Liu J, Heikkilä P, Janne J & Voutilainen R 2002 Association of H19 promoter methylation with the expression of H19 and IGF-II genes in adrenocortical tumors. *Journal of Clinical Endocrinology and Metabolism* **87** 1170–1176. (doi:10.1210/jc.87.3.1170)
- Gicquel C, Bertagna X, Schneid H, Francillard-Leblond M, Luton JP, Girard F & Le Bouc Y 1994 Rearrangements at the 11p15 locus and overexpression of insulin-like growth factor-II gene in sporadic adrenocortical tumors. *Journal of Clinical Endocrinology and Metabolism* **78** 1444–1453. (doi:10.1210/jc.78.6.1444)
- Gicquel C, Bertagna X, Gaston V, Coste J, Louvel A, Baudin E, Bertherat J, Chapuis Y, Duclos JM, Schlumberger M et al. 2001 Molecular markers and long-term recurrences in a large cohort of patients with sporadic adrenocortical tumors. *Cancer Research* **61** 6762–6767.
- Giordano TJ, Kuick R, Else T, Gauger PG, Vinco M, Bauersfeld J, Sanders D, Thomas DG, Doherty G & Hammer G 2009 Molecular classification and prognostication of adrenocortical tumors by transcriptome profiling. *Clinical Cancer Research* **15** 668–676. (doi:10.1158/1078-0432.CCR-08-1067)
- Hammarstedt L, Muth A, Wangberg B, Bjorneld L, Sigurjonsdottir HA, Gotherstrom G, Almqvist E, Widell H, Carlsson S, Ander S et al. 2010 Adrenal lesion frequency: a prospective, cross-sectional CT study in a defined region, including systematic re-evaluation. *Acta Radiologica* **51** 1149–1156. (doi:10.3109/02841851.2010.516016)
- Han Y, Chen J, Zhao X, Liang C, Wang Y, Sun L, Jiang Z, Zhang Z, Yang R, Li Z et al. 2011 MicroRNA expression signatures of bladder cancer revealed by deep sequencing. *PLoS ONE* **6** e18286. (doi:10.1371/journal.pone.0018286)
- Haussecker D & Proudfoot NJ 2005 Dicer-dependent turnover of intergenic transcripts from the human beta-globin gene cluster. *Molecular and Cellular Biology* **25** 9724–9733. (doi:10.1128/MCB.25.21.9724-9733.2005)
- Iivasmäki V, Kahri AI, Miettinen PJ & Voutilainen R 1993 Insulin-like growth factors (IGFs) and their receptors in adrenal tumors: high IGF-II expression in functional adrenocortical carcinomas. *Journal of Clinical Endocrinology and Metabolism* **77** 852–858. (doi:10.1210/jc.77.3.852)
- Karginov FV, Conaco C, Xuan Z, Schmidt BH, Parker JS, Mandel G & Hannon GJ 2007 A biochemical approach to identifying microRNA targets. *PNAS* **104** 19291–19296. (doi:10.1073/pnas.0709971104)
- Karube Y, Tanaka H, Osada H, Tomida S, Tatematsu Y, Yanagisawa K, Yatabe Y, Takamizawa J, Miyoshi S, Mitsudomi T et al. 2005 Reduced expression of Dicer associated with poor prognosis in lung cancer patients. *Cancer Science* **96** 111–115. (doi:10.1111/j.1349-7006.2005.00015.x)
- Kim DH, Villeneuve LM, Morris KV & Rossi JJ 2006 Argonaute-1 directs siRNA-mediated transcriptional gene silencing in human cells. *Nature Structural & Molecular Biology* **13** 793–797. (doi:10.1038/nsmb1142)
- Laurell C, Velazquez-Fernandez D, Lindsten K, Juhlin C, Enberg U, Geli J, Hoog A, Kjellman M, Lundberg J, Hamberger B et al. 2009 Transcriptional profiling enables molecular classification of

- adrenocortical tumours. *European Journal of Endocrinology* **161** 141–152. (doi:10.1530/EJE-09-0068)
- Lee JY, Kim H, Ryu CH, Kim JY, Choi BH, Lim Y, Huh PW, Kim YH, Lee KH, Jun TY *et al.* 2004 Merlin, a tumor suppressor, interacts with transactivation-responsive RNA-binding protein and inhibits its oncogenic activity. *Journal of Biological Chemistry* **279** 30265–30273. (doi:10.1074/jbc.M312083200)
- Li D, Zhao Y, Liu C, Chen X, Qi Y, Jiang Y, Zou C, Zhang X, Liu S, Wang X *et al.* 2011 Analysis of MiR-195 and MiR-497 expression, regulation and role in breast cancer. *Clinical Cancer Research* **17** 1722–1730. (doi:10.1158/1078-0432.CCR-10-1800)
- Ma Z, Swede H, Cassarino D, Fleming E, Fire A & Dadras SS 2011 Up-regulated Dicer expression in patients with cutaneous melanoma. *PLoS ONE* **6** e20494. (doi:10.1371/journal.pone.0020494)
- Martello G, Rosato A, Ferrari F, Manfrin A, Cordenonsi M, Dupont S, Enzo E, Guzzardo V, Rondina M, Spruce T *et al.* 2010 A microRNA targeting dicer for metastasis control. *Cell* **141** 1195–1207. (doi:10.1016/j.cell.2010.05.017)
- Melo SA, Ropero S, Moutinho C, Aaltonen LA, Yamamoto H, Calin GA, Rossi S, Fernandez AF, Carneiro F, Oliveira C *et al.* 2009 A TARBP2 mutation in human cancer impairs microRNA processing and DICER1 function. *Nature Genetics* **41** 365–370. (doi:10.1038/ng.317)
- Melo S, Villanueva A, Moutinho C, Davalos V, Spizzo R, Ivan C, Rossi S, Setien F, Casanovas O, Simo-Riudalbas L *et al.* 2011 Small molecule enoxacin is a cancer-specific growth inhibitor that acts by enhancing TAR RNA-binding protein 2-mediated microRNA processing. *PNAS* **108** 4394–4399. (doi:10.1073/pnas.1014720108)
- Merritt WM, Lin YG, Han LY, Kamat AA, Spannuth WA, Schmandt R, Urbauer D, Pennacchio LA, Cheng JF, Nick AM *et al.* 2008 Dicer, Drosha, and outcomes in patients with ovarian cancer. *New England Journal of Medicine* **359** 2641–2650. (doi:10.1056/NEJMoa0803785)
- Muralidhar B, Goldstein LD, Ng G, Winder DM, Palmer RD, Gooding EL, Barbosa-Morais NL, Mukherjee G, Thorne NP, Roberts I *et al.* 2007 Global microRNA profiles in cervical squamous cell carcinoma depend on Drosha expression levels. *Journal of Pathology* **212** 368–377. (doi:10.1002/path.2179)
- Özata DM, Caramuta S, Velazquez-Fernandez D, Akcakaya P, Xie H, Höög A, Zedenius J, Backdahl M, Larsson C & Lui WO 2011 The role of microRNA deregulation in the pathogenesis of adrenocortical carcinoma. *Endocrine-Related Cancer* **18** 643–655. (doi:10.1530/ERC-11-0082)
- Patterson EE, Holloway AK, Weng J, Fojo T & Kebebew E 2011 MicroRNA profiling of adrenocortical tumors reveals miR-483 as a marker of malignancy. *Cancer* **117** 1630–1639. (doi:10.1002/cncr.25724)
- de Reynies A, Assie G, Rickman DS, Tissier F, Groussin L, Rene-Corail F, Dousset B, Bertagna X, Clauser E & Bertherat J 2009 Gene expression profiling reveals a new classification of adrenocortical tumors and identifies molecular predictors of malignancy and survival. *Journal of Clinical Oncology* **27** 1108–1115. (doi:10.1200/JCO.2008.18.5678)
- Roh MS, Kwak JY, Kim SJ, Lee HW, Kwon HC, Hwang TH, Choi PJ & Hong YS 2005 Expression of double-stranded RNA-activated protein kinase in small-size peripheral adenocarcinoma of the lung. *Pathology International* **55** 688–693. (doi:10.1111/j.1440-1827.2005.01892.x)
- Sand M, Gambichler T, Skrygan M, Sand D, Scola N, Altmeyer P & Bechara FG 2010 Expression levels of the microRNA processing enzymes Drosha and dicer in epithelial skin cancer. *Cancer Investigation* **28** 649–653. (doi:10.3109/07357901003630918)
- Sand M, Gambichler T, Sand D, Altmeyer P, Stuecker M & Bechara FG 2011 Immunohistochemical expression patterns of the microRNA-processing enzyme Dicer in cutaneous malignant melanomas, benign melanocytic nevi and dysplastic melanocytic nevi. *European Journal of Dermatology* **21** 18–21.
- Schmitz KJ, Helwig J, Bertram S, Sheu SY, Suttrop AC, Seggewiss J, Willscher E, Walz MK, Worm K & Schmid KW 2011 Differential expression of microRNA-675, microRNA-139-3p and microRNA-335 in benign and malignant adrenocortical tumours. *Journal of Clinical Pathology* **64** 529–535. (doi:10.1136/jcp.2010.085621)
- Schteingart DE 2007 Adjuvant mitotane therapy of adrenal cancer – use and controversy. *New England Journal of Medicine* **356** 2415–2418. (doi:10.1056/NEJMe078087)
- Soon PS, Tacon LJ, Gill AJ, Bambach CP, Sywak MS, Campbell PR, Yeh MW, Wong SG, Clifton-Bligh RJ, Robinson BG *et al.* 2009 miR-195 and miR-483-5p identified as predictors of poor prognosis in adrenocortical cancer. *Clinical Cancer Research* **15** 7684–7692. (doi:10.1158/1078-0432.CCR-09-1587)
- Stephan EA, Chung TH, Grant CS, Kim S, Von Hoff DD, Trent JM & Demeure MJ 2008 Adrenocortical carcinoma survival rates correlated to genomic copy number variants. *Molecular Cancer Therapeutics* **7** 425–431. (doi:10.1158/1535-7163.MCT-07-0267)
- Stojadinovic A, Brennan MF, Hoos A, Omeroglu A, Leung DH, Dudas ME, Nissan A, Cordon-Cardo C & Ghossein RA 2003 Adrenocortical adenoma and carcinoma: histopathological and molecular comparative analysis. *Modern Pathology* **16** 742–751. (doi:10.1097/01.MP.0000081730.72305.81)
- Sugito N, Ishiguro H, Kuwabara Y, Kimura M, Mitsui A, Kurehara H, Ando T, Mori R, Takashima N, Ogawa R *et al.* 2006 RNAseN regulates cell proliferation and affects survival in esophageal cancer patients. *Clinical Cancer Research* **12** 7322–7328. (doi:10.1158/1078-0432.CCR-06-0515)
- Tombol Z, Szabo PM, Molnar V, Wiener Z, Tolgyesi G, Horanyi J, Riesz P, Reismann P, Patocs A, Liko I *et al.* 2009 Integrative molecular bioinformatics study of human adrenocortical tumors: microRNA, tissue-specific target prediction, and pathway analysis. *Endocrine-Related Cancer* **16** 895–906. (doi:10.1677/ERC-09-0096)
- Wang X, Wang J, Ma H, Zhang J & Zhou X 2011 Downregulation of miR-195 correlates with lymph node metastasis and poor prognosis in colorectal cancer. *Medical Oncology* **29** 919–927. (doi:10.1007/s12032-011-9880-5)
- Weiss LM, Medeiros LJ & Vickery AL, Jr 1989 Pathologic features of prognostic significance in adrenocortical carcinoma. *American Journal of Surgical Pathology* **13** 202–206. (doi:10.1097/00000478-198903000-00004)
- Xie H, Zhao Y, Caramuta S, Larsson C & Lui WO 2012 miR-205 expression promotes cell proliferation and migration of human cervical cancer cells. *PLoS ONE* **7** e46990. (doi:10.1371/journal.pone.0046990)
- Xu T, Zhu Y, Xiong Y, Ge YY, Yun JP & Zhuang SM 2009 MicroRNA-195 suppresses tumorigenicity and regulates G1/S transition of human hepatocellular carcinoma cells. *Hepatology* **50** 113–121. (doi:10.1002/hep.22919)
- Yano T, Linehan M, Anglard P, Lerman MI, Daniel LN, Stein CA, Robertson CN, LaRocca R & Zbar B 1989 Genetic changes in human adrenocortical carcinomas. *Journal of the National Cancer Institute* **81** 518–523. (doi:10.1093/jnci/81.7.518)

Received in final form 2 May 2013

Accepted 10 May 2013

Made available online as an Accepted Preprint

13 May 2013