Is *IGSF1* involved in human pituitary tumor formation?

Fabio R Faucz^{1,2,*}, Anelia D Horvath^{1,*}, Monalisa F Azevedo^{1,*}, Isaac Levy¹, Beata Bak³, Ying Wang³, Paraskevi Xekouki¹, Eva Szarek¹, Evgenia Gourgari^{1,4}, Allison D Manning¹, Rodrigo Bertollo de Alexandre^{1,2}, Emmanouil Saloustros¹, Giampaolo Trivellin¹, Maya Lodish¹, Paul Hofman⁵, Yvonne C Anderson^{5,6}, Ian Holdaway⁷, Edward Oldfield⁸, Prashant Chittiboina⁹, Maria Nesterova¹, Nienke R Biermasz¹⁰, Jan M Wit¹¹, Daniel J Bernard³ and Constantine A Stratakis^{1,4}

¹Section on Endocrinology and Genetics, Program on Developmental Endocrinology and Genetics (PDEGEN) Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), National Institutes of Health, NIH-Clinical Research Center, 10 Center Drive, Building 10, Room 1-3330, MSC1103, Bethesda, Maryland 20892, USA

²Group for Advanced Molecular Investigation, Graduate Program in Health Science, Medical School, Pontificia Universidade Católica do Paraná, Curitiba 80215, Brazil

³Department of Pharmacology and Therapeutics, McGill University, Montréal, Québec, Canada H3G 1Y6

⁴Pediatric Endocrinology Inter-institute Training Program, Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), National Institutes of Health (NIH), Bethesda, Maryland 20892, USA

⁵The Liggins Institute, University of Auckland, Auckland 1023, New Zealand

⁶Taranaki Base Hospital, New Plymouth 4310, New Zealand

⁷Auckland City Hospital & Greenlane Clinical Centre, Auckland 1142, New Zealand

⁸Department of Neurosurgery, University of Virginia Health Sciences Center, University of Virginia, Charlottesville, Virginia 22904, USA

⁹Surgical Neurology Branch, National Institute for Neurological Diseases and Stroke (NINDS), National Institutes of Health, Bethesda, Maryland 20892, USA

Departments of ¹⁰Endocrinology and Metabolic Disorders, and ¹¹Pediatrics, Leiden University Medical Center, Leiden 2333, The Netherlands

*F R Faucz, A D Horvath and M F Azevedo contributed equally to this work

Correspondence
should be addressed
to C A Stratakis
Email
stratakc@mail.nih.gov

Abstract

IGSF1 is a membrane glycoprotein highly expressed in the anterior pituitary. Pathogenic mutations in the *IGSF1* gene (on Xq26.2) are associated with X-linked central hypothyroidism and testicular enlargement in males. In this study, we tested the hypothesis that IGSF1 is involved in the development of pituitary tumors, especially those that produce growth hormone (GH). *IGSF1* was sequenced in 21 patients with gigantism or acromegaly and 92 healthy individuals. Expression studies with a candidate pathogenic *IGSF1* variant were carried out in transfected cells and immunohistochemistry for IGSF1 was performed in the sections of GH-producing adenomas, familial somatotroph hyperplasia, and in normal pituitary. We identified the sequence variant p.N604T, which *in silico* analysis suggested could affect IGSF1 function, in two male patients and one female with somatotroph hyperplasia from the same family. Of 60 female controls, two carried the same variant and seven were heterozygous for other variants. Immunohistochemistry showed increased IGSF1 staining in the GH-producing tumor from the patient with the *IGSF1* p.N604T variant compared with a GH-producing adenoma from a patient negative for any *IGSF1* variants and with normal control pituitary tissue. The *IGSF1* gene appears polymorphic in the general population. A potentially pathogenic variant identified in the germline of three patients with gigantism from the same family (segregating with the disease) was also detected in two healthy female controls.

Key Words

- ▶ pituitary tumor
- ▶ IGSF1
- ▶ growth hormone
- ▶ overgrowth
- ▶ acromegaly
- ▶ gigantism

Variations in IGSF1 expression in pituitary tissue in patients with or without *IGSF1* germline mutations point to the need for further studies of IGSF1 action in pituitary adenoma formation.

Endocrine-Related Cancer
(2015) 22, 47–54

Introduction

Gigantism, a rare condition that causes abnormal growth in children, often has a genetic etiology. Indeed, cases running in a family were first reported in the late nineteenth century (de Herder 2012). Growth hormone (GH)-secreting pituitary adenoma and/or hyperplasia are the main causes of gigantism in childhood. These lesions are much more common in adults and have an annual incidence of ~3/1 000 000 and a prevalence of about 60/1 000 000 (Ezzat *et al.* 2004).

IGSF1 is a plasma membrane glycoprotein encoded by the 'Ig superfamily, member 1' (*IGSF1*) gene, located on Xq26.2. This gene is conserved in mammals and is expressed with transcripts of different lengths in many tissues, including muscle, heart, brain, testis, and pancreas (Frattoni *et al.* 1998). A previous study by Sun *et al.* (2012) demonstrated that *IGSF1* is highly expressed in the Rathke's pouch and the adult anterior pituitary in humans. IGSF1 deficiency has been linked to congenital central hypothyroidism (CCH), hypoprolactinemia, delayed puberty, testicular enlargement, increased body weight, and GH deficiency (Sun *et al.* 2012, Joustra *et al.* 2013, Nakamura *et al.* 2013, Tajima *et al.* 2013), which is mainly observed in males, as expected from an X-linked genetic defect. In *Igsf1*-knockout mice, a decrease in pituitary and circulating thyroid-stimulating hormone (TSH) was observed, most probably secondary to impaired thyrotropin-releasing hormone (TRH) receptor expression and signaling (Sun *et al.* 2012).

Based on the recent work from Sun *et al.* (2012), we have investigated *IGSF1* germline variations in patients with gigantism and/or familial acromegaly from the NIH data registry and in healthy controls. We also tested the expression of IGSF1 in GH-producing adenomas. Although our data do not prove a definitive link between pituitary tumor formation and *IGSF1*, the variation in its pituitary expression and the high number of polymorphisms suggests that *IGSF1* should be studied further as a possible modifier of somatotropinoma formation and/or their clinical expression.

Materials and methods

Subjects & protocol

The *IGSF1* gene was screened for germline mutations in 21 patients (seven females and 14 males; one female and two males from the same family) with gigantism or acromegaly and in 92 previously described controls (100% white Americans, 60 females, and 32 males) with a negative family history of endocrine disorders (Horvath *et al.* 2009). All patients were previously reported (Stratakis *et al.* 2010, Glasker *et al.* 2011). Gigantism or acromegaly were diagnosed based on established criteria (Cook *et al.* 2004): high IGF1 levels according to age and sex and serum GH concentration >1 ng/ml after a 2-h 75 g oral glucose tolerance test in an appropriate clinical context, and pituitary macro- (>10 mm) or micro- (<10 mm) adenomas or pituitary hyperplasia in magnetic resonance imaging (MRI) imaging. Leukocyte DNA was isolated from each patient. Written informed consent was obtained from all participants and the study was approved by the Institutional Review Boards of the participating institutions.

IGSF1 sequencing analysis

DNA was extracted from peripheral blood leucocytes according to manufacturer's protocols (Qiagen). For all patients and controls, the complete *IGSF1*-coding and flanking intronic sequence was analyzed, as described previously (Faucz *et al.* 2011) using the primers and conditions described in Supplementary Table 1 (see section on supplementary data given at the end of this article).

In silico analyses

The Sorting Intolerant from Tolerant (SIFT) (<http://sift.bii.a-star.edu.sg/>), the Align-GVGD (Grantham Variation Grantham Deviation) (<http://agvgd.iarc.fr/>), and PolyPhen-2 (Polymorphism Phenotyping v2) (<http://genetics.bwh.harvard.edu/pph2/>) software packages were used to predict the pathogenic potential of the identified missense variants in *IGSF1*.

Cell cultures

GH3 cells (rat somatomammotroph pituitary cell line) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with high glucose (4500 mg/l), 10% fetal bovine serum (FBS), penicillin (100 IU/ml), and streptomycin (100 mg/ml) in a humidified atmosphere at 37 °C with 5% CO₂. HEK293 cells were cultured as described previously (Sun *et al.* 2012).

Transfection and ELISA

The p.N604T (c.1811A>C, rs146462069, NM_001170961.1) *IGSF1* mutation was introduced by overlapping PCR in a pCMV6 *IGSF1* gene open-reading frame plasmid (ORIGENE, Rockville, MD, USA; cat#209621) or by site-directed mutagenesis as described in Sun *et al.* (2012). Approximately, 1×10^5 GH3 cells were plated per well in 12-well cluster dishes overnight (37 °C), washed, and replenished with Opti-MEM. GH3 cell lines were transiently transfected by electroporation using the Basic Nucleofector Kit for Primary Mammalian Endothelial Cells (Lanza, Basel, Switzerland; cat#VCA-1001) following the manufacturer's protocol. The cells were transfected with plasmid DNA (6 µg), expressing either the WT or the variant form of *IGSF1* and harvested 48 h after the transfection. The supernatant from GH3 transfected cells was analyzed for GH expression levels using the rat/mouse GH ELISA Kit (cat# EZRMGH-45K; Millipore, St Charles, MO, USA) following the manufacturer's instructions.

Pulse-chase analysis (metabolic labeling)

Metabolic labeling studies were performed as described by Rejon *et al.* (2013). Briefly, 1×10^6 HEK293 cells were seeded in 35 mm dishes and transiently transfected the following day with 2 µg HA-tagged WT or p.N604T variant *IGSF1* expression vector using Lipofectamine 2000 (Invitrogen), following the manufacturer's instructions. The cells were cultured in methionine and cysteine-free DMEM (supplemented with 4 mM glutamine) for 3 h at 37 °C. The culture medium was then additionally supplemented with 198 µCi [³⁵S] methionine/[³⁵S] cysteine (per ml) for 15 min. The cells were washed twice with warm PBS and then incubated in DMEM/10% FBS supplemented with 2 mM methionine, 2 mM cysteine, and 4 mM glutamine. Following 2, 4, 8, or 24 h at 37 °C cells were lysed for 15 min, on ice, in 200 µl lysis buffer (PBS with 0.5% deoxycholate, 1% Triton X-100, 10 mM EDTA (pH 8.0), 1 mM phenylmethylsulfonyl fluoride, and Complete Mini

Protease Inhibitor Cocktail tablets, Roche). At the various time points cells were lysed after 15 min incubation with [³⁵S] methionine/[³⁵S] cysteine. Protein extracts were centrifuged for 15 min at 16 000 g to remove insoluble material. Ten microliter of supernatant was saved to assess the effectiveness of the labeling ('total') and the remaining immunoprecipitate (IP) with EZ-view Red anti-HA affinity gel, according to the instructions of the manufacturer (Sigma). The gel-bound proteins were eluted by adding 16 µl of 2× loading buffer containing β-mercaptoethanol and boiling for 5 min at 95 °C minimum. The proteins were resolved by SDS-PAGE (8% Tris-Glycine) and visualized by autoradiography.

Membrane trafficking analyses

Membrane transport of WT and p.N604T variant forms of *IGSF1* was assessed by immunofluorescence or cell-surface biotinylation of transiently transfected HEK293 cells, as described previously (Nakamura *et al.* 2013, Sun *et al.* 2012).

Immunohistochemistry

Human pituitary tissue used for immunostaining was obtained during surgery, formalin-fixed, and paraffin-embedded, and sections (5 µm) were mounted onto 3-aminopropyl-triethoxylasine coated slides (Sigma Chemical, Co.). Routine staining with hematoxylin and eosin (H&E, Histoserv, Inc, Germantown, MD, USA) was performed on several sections across each sample. Unstained slides were used for immunostaining. The procedure is outlined as follows: the sections were deparaffinized with Histoclear (National Diagnostics, Atlanta, GA, USA) and rehydrated with a graded series of ethanol (absolute, 95, 70, and 50% ethanol and distilled water), followed by antigen retrieval, which was performed by boiling the tissue sections in the Antigen Unmasking Solution (pH 6; Vector Laboratories, Burlingame, CA, USA) for 20 min in a pressure cooker; the sections were then allowed to cool to room temperature (20 min). The sections were immunostained using the antibody against IGSF1 (rabbit anti-human diluted 1:1000, Abcam ab66509, Cambridge, MA, USA). Immunostaining was identified by colorimetric staining using the ImmPRESS Polymer Detection Kit (Rabbit; Vector Laboratories) and counterstained with hematoxylin. Briefly, once sections had cooled to room temperature following antigen retrieval, sections were blocked in 0.3% (v/v) H₂O₂ (Sigma) made in methanol for 30 min. After a wash with Tris-buffered Saline (TBS, pH 7.4; diluted

to 1× from a 10× stock, Quality Biologicals, Inc., Gaithersburg, MD) containing 0.01% Tween 20 (Life Technologies) (TBS-T; 1×15 min) sections were blocked for 1 h at room temperature using 2.5% normal horse serum (provided with the ImmPRESS Polymer Detection Kit). The tissue sections were then incubated with primary antibody made in blocking serum, in a humidified chamber overnight at 4 °C. They were subsequently washed with TBS-T (3×5 min) and incubated for 30 min at room temperature with the ImmPRESS reagent, followed by another washing step in TBS-T (3×5 min). They were incubated with counterstain ImmPACT DAB peroxidase substrate (Vector Laboratories), as per manufacturer's instructions. The sections were then counterstained with Mayer's hematoxylin, mounted, and coverslips applied with VectaMount AQ Aqueous Mounting Medium (Vector Laboratories), or mounted without counterstaining.

Homology model

Homology models of the 6th Ig loop in IGSF1 with Asn or Thr at position 604 were generated using Phyre2 (<http://www.sbg.bio.ic.ac.uk/phyre2/>) and Quark (<http://zhanglab.ccmb.med.umich.edu/QUARK/>). Figures of the resulting models were generated using PyMol (<http://pymol.org/>).

Statistical analyses

Data are described as frequencies and percentages, mean ± standard deviation or median (inter-quartile range), as appropriate, and were analyzed using SAS v9.1 (SAS, Inc, Cary, NC, USA). All experiments were repeated at least

three times. Categorical data were compared using the Fisher's exact test. A *P* value ≤0.05 was considered to be statistically significant.

Results

We identified a nonsynonymous genetic *IGSF1* variant (p.N604T, c.1811A>C, rs146462069) in two males and one female (from the same family – a mother and two sons) (Glasker *et al.* 2011). All three patients showed evidence of central hypothyroidism; however, they had either pituitary somatomammotroph tumors and/or hyperplasia or were seen in our hospital after they had their first operation. Eighteen additional unrelated patients were negative for the variant. All patients have been described previously (Stratakis *et al.* 2010, Glasker *et al.* 2011). The minor allele frequency of this variant is 0.01 (<http://www.1000genomes.org/>). *In silico* analysis predicted this variant as possibly damaging. Of the 60 healthy female controls, two carried this variant and seven carried various other nonsynonymous variants (one nonsense, one frameshift, and six missense, Table 1). The variant was not seen in any of the 32 healthy males studied.

To investigate whether the p.N604T variant could be related to gigantism in our patients, we examined its impact on IGSF1 protein function. Transfection of GH3 cells with the p.N604T *IGSF1* variant did not significantly affect GH production compared with cells transfected with WT *IGSF1* (*P*=0.1561, data not shown). Metabolic studies showed that both the WT and mutant proteins exhibit the same pattern of maturation and stability when expressed in heterologous cells (see Supplementary Figure 1, see section

Table 1 Variants found in 92 healthy controls

Chrom X position (in hg19)	DNA change ^a	Protein change	Exon	SNP id	Domains	In silico modeling prediction		
						SIFT	Polyphen	Align-GVGD
130,420,579	c.70C>T	p.R24W	2	COSM1465661	Signal peptide	Tolerated	Benign	Probably damaging
130,419,882	c.238A>G	p.I80V	4	Novel	NTD	Tolerated	Benign	Benign
130,419,793	c.327G>T	p.W109C	4	Novel	NTD	Tolerated	Probably damaging	Probably damaging
130,417,192	c.714G>A	p.M238I	6	Novel	NTD	Tolerated	Benign	Benign
130,412,680	c.1811A>C	p.N604T	12	rs146462069	CTD	Tolerated	Probably damaging	Possibly damaging
130,412,680	c.1811A>C	p.N604T	12	rs146462069	CTD	Tolerated	Probably damaging	Possibly damaging
130,412,078	c.2091dupC	p.Thr698Hisfs*24	13	Novel	CTD			
130,412,018	c.2147G>A	p.G716E	13	Novel	CTD	Tolerated	Benign	Probably damaging
130,408,107	c.3844delC	p.V1282X	19	Novel	CTD			

CTD, c-terminal domain; GVGD, grantham variation/grantham deviation; NTD, n-terminal domain.

^aAll DNA changes were found in females in heterozygosity.

Figure 1

IGSF1 p.N604T traffics to the plasma membrane. (A) HEK293 cells were transfected with pcDNA3, *IGSF1* WT, or *IGSF1* p.N604T constructs and subjected to immunofluorescence under permeabilizing (bottom panels) and non-permeabilizing conditions (top panels). The proteins were detected with an antibody that recognizes the N-terminus of the *IGSF1* CTD. (B) HEK293 cells were transfected with the indicated constructs and subjected to cell surface biotinylation followed by immunoprecipitation.

Streptavidin–HRP signals of equal intensity were detected in cell lysates from WT and p.N604T transfected cells (top panel). The middle panel shows equivalent IP of the two proteins, whereas the bottom panel shows equivalent expression of the two proteins. Note that *IGSF1* migrates as a double band on SDS-PAGE. The lower band corresponds to the core (immature) glycoform, whereas the upper band is the mature glycoform.

on supplementary data given at the end of this article). Both proteins were detected in the plasma membrane by immunofluorescence (Figure 1A) or cell surface biotinylation (Fig. 1B). By western blotting analysis, the WT and p.N604T proteins were indistinguishable (Fig. 1B, lower panels, lanes 2 and 3).

Asn604 maps to Ig loop 6 (of a total of 12) in the *IGSF1* protein. *IGSF1* is co-translationally cleaved into N-terminal domains (NTD) and C-terminal domains (CTD) respectively). Ig loop 6 is at the N-terminus of the CTD. We generated a homology model of Ig loop 6 and mapped Asn604 to a solvent-exposed surface (Fig. 2). Although no *IGSF1* interacting partners have yet been identified, it is possible that the Asn 604 Thr substitution could affect a protein–protein interaction surface.

Finally, immunohistochemical analysis of our patient's pituitary tumors carrying the *IGSF1* variant showed increased *IGSF1* staining compared with normal pituitary control or a GH-producing adenoma from a patient without the *IGSF1* variant (Fig. 3). Three additional GH-producing adenomas from patients with acromegaly and no *IGSF1* variants showed a similar pattern to that of the tumor shown in Figure 3 (data not shown).

Discussion

We identified a hemizygous/heterozygous *IGSF1* variant (p.N604T) segregating with gigantism in three patients in a family. Mutations in this gene were recently linked with CCH both in humans and mice (Sun *et al.* 2012, Nakamura *et al.* 2013, Tajima *et al.* 2013), perhaps secondary to loss of

Figure 2

3D homology model of the *IGSF1* Ig loop 6 in which the variant p.N604T resides. The arrows show the position of Asn (N) 604 and Thr (T) 604.

Figure 3

Immunohistochemistry of IGSF1 in normal pituitary and in GH-producing pituitary adenomas with and without p.N604T IGSF1 sequencing variant. (A/B) Normal Pituitary; (C/D) Patient with a GH-producing adenoma

without IGSF1 sequencing variants; (E/F) Patient with pituitary hyperplasia with the p.N604T IGSF1 variant; (G/H) negative control.

TRH receptor expression and signaling in the pituitary (Sun *et al.* 2012). Considering its involvement in pituitary function and particularly in abnormal TSH secretion, as well as its demonstrated expression in murine somatotrope cells, we examined whether IGSF1 plays a causative role in our cohort of patients with gigantism or early-onset acromegaly.

Sequencing analysis of our 21 patients revealed an *IGSF1*-sequencing variant that we also observed in two female heterozygous controls. To date, 14 pathogenic mutations have been described in the *IGSF1* gene (Sun *et al.* 2012, Nakamura *et al.* 2013, Tajima *et al.* 2013), all impair either protein maturation or membrane trafficking. Among these mutations, two are complete gene deletions and the remaining 12 are located in the extracellular portion of the CTD, an indication that this may represent an important functional domain of the protein.

Eight different variants (one nonsense, one frameshift, and six missense) were identified in our control group; all in females. We would expect that if the nonsense and frameshift mutations were carried by males, they might lead to some IGSF1 deficiency phenotypes, as CCH and several of the other clinical features of IGSF1 deficiency syndrome. Among the missense variants, two (p.W109C and p.N604T) were predicted as damaging by two programs and the remaining four (p.R24W, p.I80V, p.M238I and p.G716E) were predicted as benign by at least two programs. At a minimum, these observations reveal considerable polymorphism in this gene, including variants that could impair protein function.

A difference in the expression of IGSF1 in human pituitary was also noted; IGSF1 was intensely expressed

in the tumor from one of the hemizygous (male) patients carrying the *IGSF1* variant compared with patients with GH-producing adenomas without an *IGSF1* variant or with a normal pituitary tissue. However, we did not observe differences in GH secretion from GH3 cells engineered to express the p.N604T variant versus WT *IGSF1*, indicating that the variant is not sufficient to directly affect GH production and secretion.

Previously described pathogenic *IGSF1* mutations affect the maturation and plasma membrane trafficking of the protein's CTD. However, we failed to detect any effect of the p.N604T IGSF1 variant on protein expression, maturation, stability, or membrane trafficking in heterologous cells. Therefore, if this variant alters IGSF1 function, it is likely through its extracellular activities. Given the prediction that the modified residue is solvent exposed and modifies the surface charge in the 6th Ig loop (N-terminus of the CTD), it is tempting to speculate that the variant might alter IGSF1's interaction with an extracellular partner. However, this possibility must await the identification of extracellular ligands for the protein.

Although the sequence variant detected in our patients does not appear to affect GH levels, it may be acting as a modifier in GH and prolactin production and secretion. This is supported by the observation that the same variant was also found in a Finnish family with apparent X-linked delayed puberty, the index male case showing tall stature (197 cm) and hyperprolactinemia with normal brain MRI (Joustra *et al.* 2014).

In conclusion, the *IGSF1* gene is polymorphic in the general population. The variant identified in three of our

patients with gigantism was also detected in apparently healthy female controls. It is also possible that the variant-bearing allele is in linkage disequilibrium with the actual genetic cause of the phenotype in our family (Trivellin *et al.* 2014). Indeed, the apparent increase in pituitary IGSF1 expression in patients with an *IGSF1* germline variant may indicate that *IGSF1*, while not playing a causative role in pituitary tumor development, could act synergistically in tumor development. Note that following the acceptance of this paper, our laboratory identified the genetic defect cosegregating with one of the described *IGSF1* variants (c.1811A>C) (Trivellin *et al.* 2014). As this paper indicates, we believe that the phenotype of the family carrying the defect identified by Trivellin *et al.* (2014) and first described by Glasker *et al.* (2011) is indeed modified by this potentially functional *IGSF1* variant.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/ERC-14-0465>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This study was supported by the Intramural Research Program, Eunice Kennedy Shriver National Institute of Child Health & Human Development (NICHD), National Institutes of Health (NIH) (to C A Stratakis). Additional support was provided from a post-doctoral fellowship grant from CNPq and from the University of Brasilia, Brazil (to M F Azevedo; mentor: Dr C A Stratakis), in part, by a grant from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Process: 311166/2011-3 – PQ-2 (to Fabio Rueda Faucz), and in part by a grant from Government of Canada-Canadian Institutes of Health Research, grant number CIHR MOP-133557 (to D J Bernard).

Author contribution statement

The contribution of each author is as follows: F R Faucz performed the *in silico* analyses, some of the molecular experiments described in the manuscript, and prepared the manuscript. A D Horvath and M F Azevedo, A D Manning, R B de Alexandre and G Trivellin performed most of the molecular experiments. I Levy and E Saloustros were involved in immunohistochemical studies. B Bak and Y Wang performed *in vitro* analyses of the IGSF1 variant. P Xekouki and E Szarek participated in manuscript preparation and editing; E Gourgari, M Lodish, P Hofman, Y C Anderson, I Holdaway, and N R Biermasz were the clinicians involved in clinical analysis, patients' caring, and samples collection. E Oldfield was the physician who took care of the described kindred at the NIH. P Chittiboia was involved in pathology evaluation. M Nesterova prepared the clinical specimens for genetic analysis. J M Wit was involved in the interpretation of the data and in revising the various draft versions of the manuscript. D J Bernard directed the *in vitro* investigations of the IGSF1 variant, generated the homology model, and contributed to the writing of the

manuscript. C A Stratakis was the senior investigator at NICHD, which provided most of the funding for this project under the NIH Intramural Research Program, and overall supervised the experiments, presentation of results, design of figures, and writing the manuscript.

Acknowledgements

The authors thank Dr Sjoerd Joustra for sharing unpublished data on a family with the same gene variant, Dr Dmitry Rodionov for his assistance with the homology modeling, Dr Tanya Silander for her help with preparing some of the figures, and Dr Stephen Butler for assistance in specimen collection.

References

- Cook DM, Ezzat S, Katznelson L, Kleinberg DL, Laws ER Jr, Nippoldt TB, Swearingen B, Vance ML & Force AAGT 2004 AACE Medical Guidelines for Clinical Practice for the diagnosis and treatment of acromegaly. *Endocrine Practice* **10** 213–225. (doi:10.4158/EP.10.3.213)
- Ezzat S, Asa SL, Couldwell WT, Barr CE, Dodge WE, Vance ML & McCutcheon IE 2004 The prevalence of pituitary adenomas: a systematic review. *Cancer* **101** 613–619. (doi:10.1002/cncr.20412)
- Fauz FR, Horvath A, Rothenbuhler A, Almeida MQ, Libe R, Raffin-Sanson M-L, Bertherat J, Carraro DM, Soares FA, Molina Gde *et al.* 2011 Phosphodiesterase 11A (PDE11A) genetic variants may increase susceptibility to prostatic cancer. *Journal of Clinical Endocrinology and Metabolism* **96** E135–E140. (doi:10.1210/jc.2010-1655)
- Frattini A, Faranda S, Redolfi E, Allavena P & Vezzoni P 1998 Identification and genomic organization of a gene coding for a new member of the cell adhesion molecule family mapping to Xq25. *Gene* **214** 1–6. (doi:10.1016/S0378-1119(98)00253-4)
- Glasker S, Vortmeyer AO, Lafferty AR, Hofman PL, Li J, Weil RJ, Zhuang Z & Oldfield EH 2011 Hereditary pituitary hyperplasia with infantile gigantism. *Journal of Clinical Endocrinology and Metabolism* **96** E2078–E2087. (doi:10.1210/jc.2011-1401)
- de Herder WW 2012 Familial gigantism. *Clinics* **67** (Suppl 1) 29–32. (doi:10.6061/clinics/2012(Sup01)06)
- Horvath A, Korde L, Greene MH, Libe R, Osorio P, Fauz FR, Raffin-Sanson ML, Tsang KM, Drori-Herishanu L, Patronas Y *et al.* 2009 Functional phosphodiesterase 11A mutations may modify the risk of familial and bilateral testicular germ cell tumors. *Cancer Research* **69** 5301–5306. (doi:10.1158/0008-5472.CAN-09-0884)
- Joustra SD, Schoenmakers N, Persani L, Campi I, Bonomi M, Radetti G, Beck-Peccoz P, Zhu H, Davis TM, Sun Y *et al.* 2013 The IGSF1 deficiency syndrome: characteristics of male and female patients. *Journal of Clinical Endocrinology and Metabolism* **98** 4942–4952. (doi:10.1210/jc.2013-2743)
- Joustra SD, Wehkalampi K, Oostdijk W, Biermasz NR, Howard S, Silander T, Bernard DJ, Wit JM, Dunkel L & Losekoot M 2014 IGSF1 variants in boys with familial delayed puberty. *European Journal of Pediatrics* [in press]. (doi:10.1007/s00431-014-2445-9)
- Nakamura A, Bak B, Silander TLR, Lam J, Hotsubo T, Yorifuji T, Ishizu K, Bernard DJ & Tajima T 2013 Three novel IGSF1 mutations in four Japanese patients With X-linked congenital central hypothyroidism. *Journal of Clinical Endocrinology and Metabolism* **98** E1682–E1691. (doi:10.1210/jc.2013-1224)
- Rejon CA, Ho CC, Wang Y, Zhou X, Bernard DJ & Hebert TE 2013 Cycloheximide inhibits follicle-stimulating hormone beta subunit transcription by blocking de novo synthesis of the labile activin type II receptor in gonadotrope cells. *Cellular Signalling* **25** 1403–1412. (doi:10.1016/j.cellsig.2013.03.002)
- Stratakis CA, Tichomirowa MA, Boikos S, Azevedo MF, Lodish M, Martari M, Verma S, Daly AF, Raygada M, Keil MF *et al.* 2010 The role of germline AIP, MEN1, PRKAR1A, CDKN1B and CDKN2C mutations in causing

- pituitary adenomas in a large cohort of children, adolescents, and patients with genetic syndromes. *Clinical Genetics* **78** 457–463. (doi:10.1111/j.1399-0004.2010.01406.x)
- Sun Y, Bak B, Schoenmakers N, van Trotsenburg ASP, Oostdijk W, Voshol P, Cambridge E, White JK, le Tissier P, Gharavy SN *et al.* 2012 Loss-of-function mutations in IGSF1 cause an X-linked syndrome of central hypothyroidism and testicular enlargement. *Nature Genetics* **44** 1375–1381. (doi:10.1038/ng.2453)
- Tajima T, Nakamura A & Ishizu K 2013 A novel mutation of IGSF1 in a Japanese patient of congenital central hypothyroidism without macroorchidism. *Endocrine Journal* **60** 245–249. (doi:10.1507/endocrj.EJ13-0009)
- Trivellin G, Daly AF, Faucz FR, Yuan B, Rostomyan L, Larco DO, Scherthaner-Reiter MH, Szarek E, Leal LF, Caberg JH *et al.* 2014 Gigantism and acromegaly due to Xq26 microduplications and GPR101 mutation. *New England Journal of Medicine* [in press]. (doi:10.1056/NEJMoa1408028)

Received in final form 17 November 2014

Accepted 19 November 2014