mTOR inhibitors response and mTOR pathway in pancreatic neuroendocrine tumors

Simona Falletta¹, Stefano Partelli², Corrado Rubini³, Dominik Nann⁴, Andrea Doria³, Ilaria Marinoni⁴, Vanessa Polenta², Carmelina Di Pasquale¹, Ettore degli Uberti¹, Aurel Perren⁴, Massimo Falconi² and Maria Chiara Zatelli¹

¹Department of Medical Science, Section of Endocrinology and Internal Medicine, University of Ferrara, Ferrara, Italy

²Pancreatic Surgery Unit, Pancreas Translational and Research Institute, San Raffaele Hospital, Vita-Salute San Raffaele University, Milan, Italy

³Department of Biomedical Sciences and Public Health, Polytechnic University of Marche, Ancona, Italy

⁴Institut für Pathologie, University of Bern, Bern, Switzerland

Correspondence
should be addressed
to M C Zatelli
Email
ztlmch@unife.it

Abstract

Medical therapy of pancreatic neuroendocrine tumors (P-NET) may take advantage of Everolimus treatment. However, the extent of therapeutic response cannot be predicted. This study was aimed to identify the possible predictive markers of response to Everolimus in P-NET. We found that Everolimus reduced the cell viability and induced apoptosis in primary cultures of 6 P-NET (P-NET-R), where the proliferative and antiapoptotic effects of IGF1 were blocked by Everolimus. On the contrary, 14 P-NET primary cultures (P-NET-NR) were resistant to Everolimus and IGF1, suggesting an involvement of PI3K/AKT/mTOR pathway in the mechanism of resistance. The response to Everolimus *in vitro* was associated with an active AKT/mTOR pathway and seemed to be associated with a greater clinical aggressiveness. In addition, a patient sensitive to Everolimus *in vitro* was sensitive to this drug *in vivo* also and showed a positive p-AKT immunohistochemistry (IHC) at tissue level. Similarly, a patient resistant to Everolimus treatment after surgery was not sensitive to the drug *in vitro* and had a negative p-AKT IHC staining. Therefore, present data confirm that P-NET primary cultures may be considered a model for testing medical treatment efficacy and that IHC characterization of p-AKT might help in identifying human P-NET who can benefit from Everolimus treatment. These data encourage conducting a prospective multicenter study involving different groups of P-NET patients treated with Everolimus.

Key Words

- ▶ pancreatic neuroendocrine tumors
- ▶ Everolimus
- ▶ IGF1
- ▶ primary cultures
- ▶ predictive markers

Endocrine-Related Cancer
(2016) 23, 883–891

Introduction

Pancreatic neuroendocrine tumors (P-NETs) are heterogeneous neoplasms originating from pancreatic neuroendocrine cells (Singh *et al.* 2015, Berardi *et al.* 2016), representing 1–4% of all pancreatic malignancies,

with an increasing estimated annual incidence of 1 case/100,000 inhabitants (Al-Kurd *et al.* 2014). Based on the hormonal secretion, P-NETs are divided into functioning and nonfunctioning tumors, depending

on the clinical manifestation of a syndrome related to specific hormonal hypersecretion (Halfdanarson *et al.* 2008, Missiaglia *et al.* 2010, Ro *et al.* 2013). Radical surgery represents the first-line treatment for localized disease, whereas it is often not feasible for locally advanced or metastatic tumors (Vallea *et al.* 2014). The most effective pharmaceutical option in functioning tumors is represented by somatostatin analogues (SSAs) that are useful to achieve symptomatic control. These drugs have also been recommended for the control of P-NET growth by recent guidelines (Pavel *et al.* 2016).

The mammalian target of rapamycin (mTOR) is a serine threonine kinase located downstream the PI3K/AKT signaling pathway that plays a crucial role in mediating basic cellular functions (Jiang & Liu 2008). The PI3K/AKT/mTOR pathway is constitutively activated in different NETs, including those of pancreatic origin (Capdevila *et al.* 2011). The expression of mTOR pathway components has been suggested to correlate with clinical behavior and responsiveness to mTOR inhibitors (Righi *et al.* 2010, Qian *et al.* 2013, Zatelli *et al.* 2016). The mTOR inhibitor has recently shown antitumor activity in advanced, well-differentiated and moderately differentiated P-NETs (Yao *et al.* 2011), but resistance to this treatment may develop. However, the mechanisms regulating the sensitivity/resistance to mTOR inhibitors are still scanty elucidated, and predictive biomarkers of Everolimus efficacy in P-NET are still lacking. We recently demonstrated that responsiveness to Everolimus in

human bronchial carcinoid primary cultures correlates with an active AKT/mTOR signaling pathway (Gagliano *et al.* 2013). However, PI3K/AKT/mTOR signaling does not always predict the sensitivity to mTOR inhibitors and is difficult to assess (Zatelli *et al.* 2016). Therefore, the aim of this study was to investigate the possible predictors of sensitivity to Everolimus in P-NET, to identify patients who may benefit from treatment avoiding useless side effects in resistant patients.

Materials and methods

Human pancreatic neuroendocrine tumors

Twenty samples, including 18 primary P-NETs and 2 lymph node metastases, were derived from 16 patients (10 males and 6 females; mean age 58.6 ± 4.2 years) diagnosed with P-NET, who were operated on at the University of Ancona (Pancreatic Surgery Unit, Department of Surgery) and at the University of Ferrara (Section of Endocrinology and Clinical Surgery). Patients and tumor characteristics are displayed in Table 1. Among the investigated patients, 13 did not receive medical therapy before surgery; 3 had been previously treated with somatostatin analogues (SSA) alone (1 patient), in association with chemotherapy (1 patient) or with Everolimus (1 patient, who showed disease progression after being treated with capecitabine).

Most P-NETs were diagnosed as G1 or G2 (14/16), displaying a Ki67 $\leq 10\%$, with pauci cellular focal necrosis

Table 1 Clinical characteristics of P-NET patients.

Patient no.	Sex	Age	Hormonal secretion	Diagnosis	TNM	Stage	Diameter (cm)	Grade	Mitosis no.	Ki67 (%)
1	F	67	NF	NET G2	T3 N1 M1	IV	4	2	2/50	8
2	M	74	NF	NEC G3 ^a	T2 N0 M1	IV	2.8	3	42/10	65
3	M	53	NF	NET G2	T2 N1 M0	IIB	3.4	2	2/10	9
4	M	76	NF	NET G1	T1 N0 M0	I	1.5	1	2/10	2
5 ^b	F	38	NF	NET G2	T4 N0 M0	III	9.5	2	2/10	10
6 ^c	M	56	NF	NEC G3	T3 N1 M1	IIB	8	3	5/10	25
7 ^{b,c}	F	65	NF	NET G2	T3 N1 M1	IV	2.5	2	13/10	20
8	F	32	NF	NET G1	T2 N0 M0	IB	3	1	1/50	1
9	M	77	NF	NET G1	T1 N0 M0	I	1.1	1	2/10	2
10	F	74	NF	NET G2	T2 N0 MX	IB	1.8	2	2/10	5
11	M	23	Insulinoma	NET G2	T3 N1 M0	IIB	2.5	2	2/10	7
12 ^d	M	65	NF	NET G1	T1 N0 M0	I	1.3	1	2/10	1
13	M	75	Insulinoma	NET G1	T1 N0 M0	I	1.9	1	1/10	2
14	F	65	NF	NET G1	T1 N0 M0	I	2	1	2/10	2
15 ^d	M	45	Insulinoma	NET G1	T1 N0 M0	I	0.5	1	0	<2
16	M	53	NF	NET G2	T2 N0 M0	I	2.5	2	NA	3

NA, not available; NEC, neuroendocrine carcinoma; NET, neuroendocrine tumor; NF, non-functioning.

^aPoorly differentiated neuroendocrine carcinoma; ^bNET patients for whom 1 sample from the primary tissue and 1 sample from the lymph node metastasis were available, each generating a primary culture; ^cNET patients treated with Everolimus *in vivo*; ^dNET patients for whom 2 primary tissue samples were available, each generating a primary culture.

in some cases, without lymph node (11/16) or distant metastases (13/16).

Tissue collection and primary cultures

Tissue samples were collected following the guidelines of the local committee on human research. Informed consent of the patients was obtained for disclosing clinical investigation and performing the *in vitro* study. Tissue samples were stored in RNA-later solution (Sigma) for expression studies; upon arrival in the lab, they were immediately frozen in liquid nitrogen under ribonuclease (RNase)-free conditions and stored at -80°C until protein isolation was performed. A portion of the fresh tissue was collected in culture medium for primary culture studies; upon arrival in the lab, the tissue was immediately processed for primary culture experiments as described previously, with minor modifications (Molè *et al.* 2011). Tumor cells were resuspended in F-12 with 10% fetal bovine serum (FBS) and antibiotics (Euroclone Ltd, Wetherby, UK), seeded at 2×10^4 cells/well in 96-well black plates and incubated at 37°C in a humidified atmosphere of 5% CO_2 -95% air, as described previously (Zatelli *et al.* 2010a). After approximately 18 h, cells were treated with test compounds, with further evaluation of cell viability and/or caspase activity.

Compounds

Everolimus was provided by Novartis; IGF1 was purchased from PeproTek (Rocky Hill, NJ, USA). All other reagents were purchased from Sigma, if not otherwise indicated.

Viable cell number and caspase activation assessment

Cell viability and caspase activation were measured as described previously (Zatelli *et al.* 2006, Zatelli *et al.* 2010a,b) using the ATPlite kit (PerkinElmer Life Sciences) and the caspase-Glo 3/7 assay (Promega), respectively. Briefly, 20 P-NETs primary cultures, derived from 16 patients, were treated with or without 100 nM Everolimus for 48 h in the absence or in the presence of 100 nM IGF1. Luminescent output (relative luminescence units, RLU) was recorded after 48 h for each assessment by the Envision Multilable Reader (PerkinElmer). Results are expressed as mean value $\pm$ S.E.M. percent RLU vs untreated control cells.

According to the response to Everolimus, samples were divided into primary cultures displaying a significant reduction ($P < 0.05$ vs untreated cells) in

cell viability under Everolimus treatment, indicated as responders (P-NET-R), and into primary cultures in which Everolimus did not reduce cell viability, indicated as non-responders (P-NET-NR).

Kinase activity assay

Protein isolation from frozen human P-NET tissues was performed as described previously (Gagliano *et al.* 2013). RIPA buffer (Pierce) was used to dissolve tissues that were kept in ice for 30 min, and then centrifuged for 10 min. Protein concentration in the supernatant was measured by the BCA Protein Assay Reagent Kit (Pierce), as described previously (Tagliati *et al.* 2006). Each sample was evaluated in 3–6 replicates by assessing the same total protein amount. Phosphorylated levels of IGF1R (Tyr1135/1136), AKT (Ser473), mTOR (Ser2448) and 4EBP1 (Thr37/46) were measured using the AlphaScreen SureFire assays (PerkinElmer Life Sciences). Normalization against glyceraldehyde-3-phosphate dehydrogenase (GAPDH) using the appropriate AlphaScreen SureFire assay kit was performed. Results are expressed as mean value $\pm$ S.E.M. AlphaScreen signal (counts).

Tissue microarray construction

Paraffin blocks were available for 15 P-NET patients. Tissue microarrays (TMA) were manually assembled from formalin-fixed, paraffin-embedded tissues. Three representative 1 mm diameter normal and tumoral tissue cores were taken from each block, for a total of 45 specimens. Multiple 3- μm sections were cut with a Leica microtome (Leica Microsystems), dried at 60°C and then transferred to adhesive-coated slides for immunohistochemical staining.

Immunohistochemistry

Immunohistochemistry (IHC) was performed by an anti-p-AKT (Ser473) antibody obtained by Abcam. Slides were de-paraffinized and re-hydrated in dewax solution (Leica Biosystems), and endogenous peroxidase activity was blocked with H_2O_2 solution for 4 min. After antigen retrieval using citrate buffer, pH 6.0 (15 min at 100°C), slides were incubated with the anti-p-AKT antibody (1:300) for 30 min, at room temperature. Immunoreactions were revealed by using the Bond Polymer Refine Kit on the automated system BOND RX (both from Leica Biosystems). Antigen-antibody complexes were detected using the cobalt-3,3'-diaminobenzidine (Co-DAB)

as the chromogen (Leica Biosystems). Samples were counterstained with hematoxylin and mounted in Aquatex (Merck). The specificity of all reactions was validated in parallel control sections omitting the primary antibodies for each IHC run. Endometrial cancers that are known to have, in the majority of cases, a PTEN mutation leading to PI3K/AKT pathway constitutive activation were used as positive controls.

Evaluation of staining of tissue slides

Tumor staining characteristics were reviewed and interpreted independently by two pathologists (A P and D N). IHC data were visually scored. Intensity and location (nuclear and membranous) were taken into consideration. The immunostaining intensity was evaluated subjectively as being negative (0), weak (1), moderate (2) and strong (3) and then the staining was classified into two categories: negative (no immunostaining) and positive (weak-moderate and strong immunostaining), in more than 10% of tumor cells.

Statistical analysis

Results are expressed as the mean $\pm$ S.E.M. Concerning the results of cell viability and caspase 3/7 activation experiments, a preliminary analysis was carried out to determine whether the datasets conformed to a normal distribution. The results were compared within each group and between the groups using ANOVA. If the *F* values

were significant ($P < 0.05$), Student's paired or unpaired *t*-test was used to evaluate individual differences between the means. *P* values < 0.05 were considered significant. For the AlphaScreen assays, after proof of homogeneity of variance and normality test, Student's unpaired *t*-test was used. In case of non-normal distribution, Mann-Whitney test was used.

To compare the clinicopathological features, a preliminary analysis was carried out to determine whether the datasets conformed to a normal distribution and then Mann-Whitney test was used, due to not normal distribution for Ki67, tumor size and number of mitoses. For sex, primary vs. metastatic tissue, pre-treatment with SSA, Octreoscan/68GA-PET uptake, site, TNM, stage, necrosis and tumor grade, chi-square test corrected for linearity was applied. Differences in age were assessed by the Student's *t*-test. Data were analyzed using GraphPad (Prism v-5.0).

Results

Effects of Everolimus and IGF1 on P-NET primary cultures

Six P-NET primary cultures were defined as P-NET-R and 14 P-NETs were defined as P-NET-NR. As shown in Fig. 1A (upper panel), in P-NET-R, Everolimus significantly reduced the cell viability (-29.9% ; $P < 0.05$ vs untreated cells), an effect completely counteracted by co-incubation with IGF1. In P-NET-NR (Fig. 1B, upper panel), cell viability was significantly induced by Everolimus

Figure 1

Effects of Everolimus and IGF1 on P-NET primary cultures. P-NET primary cultures were incubated in 96-well plates for 48h in a culture medium supplemented with 100nM Everolimus with (black bars) or without 100nM IGF1 (white bars); control cells were treated with a vehicle solution. Cell viability and caspase activation of each primary culture were measured as a luminescent signal. According to the response to Everolimus in terms of cell viability inhibition, the samples were divided into 'responder' (6 samples) (A) and 'non-responder' (14 samples) (B). Data from P-NET primary cultures were evaluated independently with six replicates each, and expressed as the mean value $\pm$ S.E.M. percent cell viability inhibition vs untreated control cells (upper panels) or as the mean value $\pm$ S.E.M. percent caspase activation vs untreated control cells (lower panels). * $P < 0.05$ and *** $P < 0.001$ vs untreated control cells. # $P < 0.05$ and ### $P < 0.001$ vs cells treated with Everolimus alone.

(+30.2%; $P < 0.001$ vs control cells) and was not affected by IGF1, alone or in combination with Everolimus.

To verify whether Everolimus modifies cell viability by influencing the apoptotic process, caspase 3/7 activation was evaluated. As shown in Fig. 1A (lower panel), in P-NET-R Everolimus significantly induced caspase activation (+73%; $P < 0.001$ vs untreated cells), an effect completely counteracted by co-incubation with IGF1. In P-NET-NR (Fig. 1B, lower panel), both Everolimus and IGF1 did not significantly affect caspase activation. These results indicate that P-NET may respond differently to Everolimus *in vitro* and that P-NET-NRs are not sensitive to IGF1 either, suggesting that IGF1 transduction pathway is not active in these tissues. Our data suggest that IGF1 pathway may be involved in the mechanism of resistance to Everolimus *in vitro*.

IGF1/mTOR signaling pathway expression in P-NET tissues

As IGF1 was capable of reversing the inhibitory effects of Everolimus on cell viability in P-NET-R, we investigated the levels of IGF1 downstream signaling proteins involved in mTOR pathway. Proteins isolated from frozen P-NET tissues, classified as P-NET-R and P-NET-NR on the basis of the results of primary cultures, were used to evaluate the expression levels of phosphorylated IGF1 R (p-IGF1 R), AKT (p-AKT), mTOR (p-mTOR) and 4EBP1 (p-4EBP1) by using AlphaScreen SureFire assay. As shown in Fig. 2A, we found that p-IGF1 R protein levels were significantly ($P < 0.001$) >2-fold lower in P-NET-NR compared with those

in P-NET-R. Similarly, p-AKT levels were >2-fold lower in P-NET-NR compared with those in P-NET-R (Fig. 2B); however, statistical significance was not reached, possibly due to the small number of samples. Moreover, p-mTOR and p-4EBP1 protein levels were significantly ($P < 0.02$ and $P < 0.001$) >2-fold lower in P-NET-NR compared with P-NET-R (Fig. 2C and D). These data indicate that response to Everolimus *in vitro* is associated with an active AKT/mTOR pathway.

Clinical and pathological correlations

When the clinical characteristics of the patients were evaluated according to Everolimus responsiveness *in vitro*, there was no significant difference between patients with P-NET-R compared with those with P-NET-NR in terms of sex, age, primary vs metastatic tissue, pre-treatment with SSA, Octreoscan/ ^{68}Ga -PET uptake, site, TNM, stage, size, number of mitoses and necrosis. However, ki67 was significantly ($P < 0.05$) higher in P-NET-R (median 10%) than that in P-NET-NR (median 2%) (Fig. 3A). In addition, a trend for a linear correlation between tumor grade and response to Everolimus was found. All G3 P-NET responded to the treatment *in vitro*, whereas only 14% of G1 P-NET were P-NET-R (Fig. 3B).

Expression of mTOR pathway components in P-NET and association with clinical outcomes

To validate the identified putative markers of response to Everolimus in P-NET, the expression of p-AKT protein was

Figure 2

IGF1/mTOR signaling pathway expression in P-NET tissues. Total proteins were isolated from 20 P-NET tissues and AlphaScreen analysis for p-IGF1 R (A), p-AKT (B), p-mTOR (C) and p-4EBP1 (D) expression in the pooled P-NET R and in the pooled P-NET NR tissues was performed, as described in the 'Materials and methods' section. Values are expressed after normalization against glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as AlphaScreen signal (counts). ** $P < 0.02$ and *** $P < 0.001$ vs P-NET R.

Figure 3

Patients clinical characteristics and Everolimus responsiveness *in vitro*. (A) Median values and range of Ki67 labelling index in P-NET-R and in P-NET-NR. * $P < 0.05$ vs P-NET-R.

(B) % P-NET-R tissues according to tumor grade. G1 P-NET-R = 1/7; G2 P-NET-R = 2/7; G3 P-NET-R = 2/2.

evaluated using IHC. Four tissue samples did not provide reliable results due to technical problems. Therefore, we obtained reliable results for 11 patients, 3 P-NET-R and 8 P-NET-NR. IHC for p-AKT was positive in all the paraffin-embedded tissues that correspond *in vitro* to 3 P-NET-R. IHC for p-AKT was positive in 5 out of 8 paraffin-embedded tissues that corresponded *in vitro* to P-NET-NR. There was no statistically significant correlation between p-AKT positivity by IHC and responsiveness to Everolimus *in vitro*, possibly due to the low number of available samples. The response to Everolimus *in vitro*, as well as p-AKT protein levels in frozen tissues were also compared with the response to Everolimus *in vivo*, in terms of tumor stabilization/progression and survival. Two patients had been treated with Everolimus after surgery due to the presence of non-resectable liver metastases. The first patient was a 56-year-old male operated on for a well-differentiated non-functioning G3 P-NET (T3N1M1, Ki67=25%), which turned out to be a P-NET-R *in vitro*. The patient displayed stabilization of the

disease during Everolimus treatment, with a progression-free survival of 21 months (still alive at the end of the study). Furthermore, he showed positive p-AKT IHC at tissue level (Fig. 4A). The second patient was a 65-year-old female operated on for a non-functioning G2 P-NET (T3N1M1, Ki67=20%), that turned out to be a P-NET-NR *in vitro*. The patient displayed disease progression after 12 months of treatment with Everolimus, with a survival of 21 months (still alive at the end of the study). Data from IHC showed negative p-AKT staining (Fig. 4B).

Discussion

Our study shows that P-NET primary cultures are a useful model for testing medical therapy *in vitro*. We also identified putative markers that may predict the response to Everolimus treatment *in vivo*. In our settings, 30% of human P-NET primary cultures responded to Everolimus treatment with a significant reduction in cell viability and induction of apoptosis, whereas the remaining 70% did not. These data may seem in contrast with the demonstrated efficacy of Everolimus in prolonging PFS in advanced P-NET (Yao et al. 2011, Wolin 2014). It should be underlined that we defined as responders those primary cultures displaying a significant cell viability reduction *in vitro* after 48 h. This parameter may not completely reflect the results obtained on clinical grounds in terms of PFS, which is influenced by many clinical variables and not by tumor growth only. On the other hand, our results may help in identifying those patients who may better benefit from treatment not only in terms of longer PFS but also in terms of tumor bulk reduction. Moreover, our series shows intra-patient and intra-tumor heterogeneity in primary cultures, further underlining P-NET heterogeneity. In addition, we show that the effect of Everolimus was influenced by IGF1 only in human P-NET tissues classified as responders to the drug *in vitro*, but not in human P-NET tissues resistant to the drug. The importance of IGF1 in regulating NET proliferation is underlined by the evidence of an enhanced and constitutive expression of IGF1 and

Figure 4

Association between p-AKT expression and clinical outcomes.

(A) Immunohistochemical expression of p-AKT in a non-functioning G3 P-NET (T3N1M1), sensitive to Everolimus after surgery, defined as P-NET-R *in vitro*. (B) Immunohistochemical expression of p-AKT in a non-functioning G2 P-NET (T3N1M1), resistant to Everolimus after surgery and defined as P-NET-NR *in vitro*.

IGF1R in gastroenteropancreatic NET, which leads to autocrine and paracrine growth stimulation (von Wichert *et al.* 2000, Höpfner *et al.* 2008). Moreover, we previously demonstrated that IGF1 exerts protective effects toward the anti-proliferative action of Everolimus in a group of medullary thyroid carcinoma primary cultures (Gentilin *et al.* 2015). Therefore, these data suggest that IGF1 pathway may play an important role in the development of Everolimus resistance. The latter represents a relevant issue because previous evidence demonstrated that some patients do not benefit from Everolimus treatment, due to the development of primary or secondary acquired resistance to this drug (Fazio 2015), despite a significant prolongation of PFS shown by the RADIANT-3 trial in a significant number of patients with P-NET (Yao *et al.* 2011). However, the precise mechanism underlying Everolimus resistance remains unknown (Zatelli *et al.* 2016).

Everolimus inhibits mTOR activity by interacting with the mTORC1 complex but not with mTORC2 complex. Selective blockade of mTORC1 by Everolimus leads to an increase in mTORC2 activity and results in a positive feedback activation of AKT, by phosphorylation on Ser473, and of IGF1/IGFR signaling due to inhibition of the S6K negative feedback (O'Reilly *et al.* 2006). As an evident association between the expression levels of PI3K/AKT/mTOR components and response to Everolimus *in vitro* was observed in bronchial carcinoids and in nonfunctioning pituitary adenomas (Zatelli *et al.* 2010a, Gagliano *et al.* 2013), we hypothesized that the investigation of mTOR pathway components could be useful to predict the responsiveness to Everolimus treatment in P-NET. In this setting, AlphaScreen data show that P-NET-R tissues display higher levels of p-IGF1R, p-AKT, p-mTOR and p-4EBP1 compared with those of P-NET-NR tissues, supporting the hypothesis that the expression of these proteins could be used to separate 'responder' tissues from 'resistant' tissues, in agreement with our previous results (Zatelli *et al.* 2010a, Gagliano *et al.* 2013). On this basis, our data suggest that the lack of response to Everolimus *in vitro* may be due to the low expression levels of PI3K/AKT/mTOR pathway components. Our results are in contrast with a recent study showing that high p-p70S6K protein levels are associated with a worse outcome under Everolimus treatment in NET (Benslama *et al.* 2016). However, the latter study was performed including a very heterogeneous group of NET of different origin. Therefore, the results obtained by Benslama and coworkers may not reflect P-NET behavior (Benslama *et al.* 2016).

Besides putative molecular markers, we also investigated whether P-NET clinical characteristics may

differ between responders and non-responders. We found a higher ki67 index in human P-NET-R compared with P-NET-NR and a trend for a linear correlation between tumor grade and response to Everolimus, suggesting that Everolimus responsiveness may associate with a greater clinical aggressiveness of P-NET.

Even though AlphaScreen represents a very sensitive method, validated by Western blot in bronchial carcinoids (Gagliano *et al.* 2013), it is not widely available. Therefore, we assessed IHC for p-AKT protein to validate the identified putative markers of response to Everolimus by means of a method that may be used also by other centers. The low number of available samples limited the significance of our findings as the correlation between p-AKT positivity by IHC and responsiveness to Everolimus *in vitro* did not reach statistical significance. The lack of correlation between AlphaScreen results obtained in fresh frozen tissues and IHC data for p-AKT expression is of importance. Although the fresh tissues were frozen after being stored several hours in RNAlater, which is a stabilizing RNA but does not influence protein phosphorylation, the paraffin-embedded tissues were processed routinely in the pathology lab with a variation in fixation times. The IHC protocols were performed on tissues that were fixed as large blocks and formalin is known to penetrate at 1 mm/h. During this time, phosphorylation levels of AKT may profoundly change (Baker *et al.* 2005), possibly explaining the different results obtained using these two different approaches. However, it is interesting to notice that p-AKT positivity by IHC was found in a P-NET-R patient who showed tumor stabilization under Everolimus treatment *in vivo*, suggesting that even P-NET-G3 may take advantage of this drug. On the contrary, IHC failed to demonstrate a strong p-AKT staining in a P-NET-NR who showed disease progression under Everolimus treatment *in vivo*. These findings suggest the presence of a good correlation between the responsiveness to Everolimus *in vitro* and that *in vivo* and provide the basis for further studies aimed at validating the identified putative markers of Everolimus responsiveness.

In conclusion, our results show that P-NET primary cultures may represent a suitable model for testing medical treatment efficacy. In addition, our data indicate that IGF1 pathway is involved in the development of resistance to Everolimus and that that IHC characterization of p-AKT may help in identifying human P-NET who can benefit from Everolimus treatment. However, these preliminary data need to be confirmed and encourage conducting a prospective multicenter study involving different groups of P-NET patients treated with Everolimus.

Declaration of interest

M C Z has received consultant fees from Novartis and Genzyme. E d U has received consultant fees from Novartis and grant support from Sanofi. The other authors have nothing to disclose concerning a possible conflict of interest.

Funding

This work was supported by grants from the Italian Ministry of Education, Research and University (FIRB RBAP11884M, RBAP1153LS), Fondazione Dott. Carlo Fornasini and Associazione Italiana per la Ricerca sul Cancro (AIRC) in collaboration with Laboratorio in rete del Tecnopolo 'Tecnologie delle terapie avanzate' (LTTA) of the University of Ferrara. The funding sources had no involvement in study design, in the collection, analysis and interpretation of data, in the writing of the report and in the decision to submit the article for publication.

Author contribution statement

S Falletta wrote the manuscript, performed primary cultures and AlphaScreen experiments and helped in setting up the immunohistochemistry experiments; S Partelli helped in providing the surgical specimens; C Rubini provided the paraffin-embedded tissues; D Nann performed the immunohistochemistry experiments; A Doria helped in providing the paraffin-embedded tissues; I Marinoni provided support for the immunohistochemistry experiments and reviewed the manuscript; V Polenta helped in providing the clinical data; C Di Pasquale helped in performing the primary culture experiments; E degli Uberti provided support for the clinical correlations; A Perren supervised the immunohistochemistry experiments and reviewed the manuscript; M Falconi supervised the surgical specimen provision and reviewed the manuscript and M C Zatelli wrote the manuscript, supervised the *in vitro* experiments and the project development.

Acknowledgements

The authors thank Dr Miguel Sampedro Nuñez for his help in the statistical evaluation.

References

- Al-Kurd A, Chapchay K, Grozinsky-Glasberg S & Mazeh H 2014 Laparoscopic resection of pancreatic neuroendocrine tumors. *World Journal of Gastroenterology* **20** 4908–4916. (doi:10.3748/wjg.v20.i17.4908)
- Baker AF, Dragovich T, Ihle NT, Williams R, Fenoglio-Preiser C & Powis G 2005 Stability of phosphoprotein as a biological marker of tumor signaling. *Clinical Cancer Research* **11** 4338–4340. (doi:10.1158/1078-0432.CCR-05-0422)
- Benslama N, Bollard J, Vercherat C, Massoma P, Roche C, Hervieu V, Peron J, Lombard-Bohas C, Scoazec JY & Walter T 2016 Prediction of response to everolimus in neuroendocrine tumors: evaluation of clinical, biological and histological factors. *Investigational New Drugs* **34** 654–662. (doi:10.1007/s10637-016-0363-6)
- Berardi R, Morgese F, Torniai M, Savini A, Partelli S, Rinaldi S, Caramanti M, Ferrini C, Falconi M & Cascinu S 2016 Medical treatment for gastro-entero-pancreatic neuroendocrine tumours. *World Journal of Gastrointestinal Oncology* **8** 389–401. (doi:10.4240/wjgs.v8.i5.389)
- Capdevila J, Salazar R, Halperin I, Abad A & Yao JC 2011 Innovations therapy: mammalian target of rapamycin (mTOR) inhibitors for the treatment of neuroendocrine tumors. *Cancer and Metastasis Review* **30** 27–34. (doi:10.1007/s10555-011-9290-3)
- Fazio N 2015 Neuroendocrine tumors resistant to mammalian target of rapamycin inhibitors: A difficult conversion from biology to the clinic. *World Journal of Clinical Oncology* **6** 194–197. (doi:10.5306/wjco.v6.i6.194)
- Gagliano T, Bellio M, Gentilin E, Molè D, Tagliati F, Schiavon M, Cavallese NG, Andriolo LG, Ambrosio MR, Rea F, et al. 2013 mTOR, p70S6K, AKT, and ERK1/2 levels predict sensitivity to mTOR and PI3K/mTOR inhibitors in human bronchial carcinoids. *Endocrine-Related Cancer* **20** 463–475. (doi:10.1530/ERC-13-0042)
- Gentilin E, Di Pasquale C, Rossi M, Tagliati F, Gagliano T, Rossi R, Pelizzo M, Merante Boschini I, degli Uberti EC & Zatelli MC 2015 Igf-I influences everolimus activity in medullary thyroid carcinoma. *Frontiers in Endocrinology* **6** 63. (doi:10.3389/fendo.2015.00063)
- Halfdanarson TR, Rabe KG, Rubin J & Petersen GM 2008 Pancreatic neuroendocrine tumors (PNETs): incidence, prognosis and recent trend toward improved survival. *Annals of Oncology* **19** 1727–1733. (doi:10.1093/annonc/mdn351)
- Höpfner M, Schuppan D & Scherübl H 2008 Treatment of gastrointestinal neuroendocrine tumors with inhibitors of growth factor receptors and their signaling pathways: recent advances and future perspectives. *World Journal of Gastroenterology* **14** 2461–2473. (doi:10.3748/wjg.14.2461)
- Jiang BH & Liu LZ 2008 Role of mTOR in anticancer drug resistance: perspectives for improved drug treatment. *Drug Resistance Updates* **11** 63–76. (doi:10.1016/j.drug.2008.03.001)
- Missiaglia E, Dalai I, Barbi S, Beghelli S, Falconi M, della Peruta M, Piemonti L, Capurso G, Di Florio A, delle Fave G, et al. 2010 Pancreatic endocrine tumors: expression profiling evidences a role for AKT-mTOR pathway. *Journal of Clinical Oncology* **28** 245–255. (doi:10.1200/JCO.2008.21.5988)
- Molè D, Gagliano T, Gentilin E, Tagliati F, Pasquali C, Ambrosio MR, Pansini G, degli Uberti EC & Zatelli MC 2011 Targeting protein kinase C by Enzastaurin restrains proliferation and secretion in human pancreatic endocrine tumors. *Endocrine-Related Cancer* **18** 439–450. (doi:10.1530/ERC-11-0055)
- O'Reilly KE, Rojo F, She QB, Solit D, Mills GB, Smith D, Lane H, Hofmann F, Hicklin DJ, Ludwig DL, et al. 2006 mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates Akt. *Cancer Research* **66** 1500–1508. (doi:10.1158/0008-5472.CAN-05-2925)
- Pavel M, O'Toole D, Costa F, Capdevila J, Gross D, Kianmanesh R, Krenning E, Knigge U, Salazar R, Pape UF, et al. 2016 ENETS consensus guidelines update for the management of distant metastatic disease of intestinal, pancreatic, bronchial neuroendocrine neoplasms (NEN) and NEN of unknown primary site. *Neuroendocrinology* **103** 172–185. (doi:10.1159/000443167)
- Qian ZR, Ter-Minassian M, Chan JA, Imamura Y, Hooshmand SM, Kuchiba A, Morikawa T, Brais LK, Daskalova A, Heafield R, et al. 2013 Prognostic significance of mTOR pathway component expression in neuroendocrine tumors. *Journal of Clinical Oncology* **31** 3418–3425. (doi:10.1200/JCO.2012.46.6946)
- Righi L, Volante M, Rapa I, Tavaglione V, Inzani F, Pelosi G & Papotti M 2010 Mammalian target of rapamycin signaling activation patterns in neuroendocrine tumors of the lung. *Endocrine-Related Cancer* **17** 977–987. (doi:10.1677/ERC-10-0157)
- Ro C, Chai W, Yu VE & Yu R 2013 Pancreatic neuroendocrine tumors: biology, diagnosis, and treatment. *Chinese Journal of Cancer* **32** 312–324. (doi:10.5732/cjc.012.10295)
- Singh S, Dey C, Kennecke H, Kocha W, Maroun J, Metrakos P, Mukhtar T, Pasiaka J, Rayson D, Rowsell C, et al. 2015 Consensus recommendations for the diagnosis and management of pancreatic neuroendocrine tumors: guidelines from a Canadian National Expert Group. *Annals of Surgical Oncology* **22** 2685–2699. (doi:10.1245/s10434-014-4145-0)

- Tagliati F, Zatelli MC, Bottoni A, Piccin D, Luchin A, Culler MD & degli Uberti EC 2006 Role of complex cyclin D1/Cdk4 in somatostatin subtype 2 receptor-mediated inhibition of cell proliferation of a medullary thyroid carcinoma cell line in vitro. *Endocrinology* **147** 3530–3538. (doi:10.1210/en.2005-1479)
- Vallea J, Eatock M, Clueit B, Zahava G, Roxanne F & Mitchell S 2014 A systematic review of non-surgical treatments for pancreatic neuroendocrine tumours. *Cancer Treatment Reviews* **40** 376–389. (doi:10.1016/j.ctrv.2013.08.007)
- von Wichert G, Jehle PM, Hoeflich A, Koschnick S, Dralle H, Wolf E, Wiedenmann B, Boehm BO, Adler G & Seufferlein T 2000 Insulin-like growth factor-I is an autocrine regulator of chromogranin A secretion and growth in human neuroendocrine tumor cells. *Cancer Research* **60** 4573–4581.
- Wolin EM 2014 Long-term Everolimus treatment of patients with pancreatic neuroendocrine tumors. *Chemotherapy* **60** 143–150. (doi:10.1159/000369780)
- Yao JC, Shah MH, Ito T, Bohas CL, Wolin EM, Van Cutsem E, Hobday TJ, Okusaka T, Capdevila J, de Vries EG, et al. 2011 Everolimus for advanced pancreatic neuroendocrine tumors. *New England Journal of Medicine* **364** 514–523. (doi:10.1056/NEJMoa1009290)
- Zatelli MC, Piccin D, Tagliati F, Bottoni A, Luchin A, Vignali C, Margutti A, Bondanelli M, Pansini GC, Pelizzo MR, et al. 2006 Selective activation of somatostatin receptor subtypes differentially modulates secretion and viability in human medullary thyroid carcinoma primary cultures: potential clinical perspectives. *Journal of Clinical Endocrinology and Metabolism* **91** 2218–2224. (doi:10.1210/jc.2006-0334)
- Zatelli MC, Minoia M, Martini C, Tagliati F, Ambrosio MR, Schiavon M, Buratto M, Calabrese F, Gentilin E, Cavallero G, et al. 2010a Everolimus as a new potential antiproliferative agent in aggressive human bronchial carcinoids. *Endocrine-Related Cancer* **17** 719–729. (doi:10.1677/ERC-10-0097)
- Zatelli MC, Gentilin E, Daffara F, Tagliati F, Reimondo G, Carandina G, Ambrosio MR, Terzolo M & degli Uberti EC 2010b Therapeutic concentrations of mitotane (o,p'-DDD) inhibit thyrotroph cell viability and TSH expression and secretion in a mouse cell line model. *Endocrinology* **151** 2453–2461. (doi:10.1210/en.2009-1404)
- Zatelli MC, Fanciulli G, Malandrino P, Ramundo V, Faggiano A, Colao A & NIKE Group 2016 Predictive factors of response to mTOR inhibitors in neuroendocrine tumours. *Endocrine-Related Cancer* **23** R173–R183. (doi:10.1530/ERC-15-0413)

Received in final form 21 September 2016

Accepted 3 October 2016

Accepted Preprint published online 3 October 2016