RESEARCH LETTER

8th edition of the AJCC/TNM staging system of thyroid cancer: what to expect (ITCO#2)

Dear Editor,

Differentiated thyroid cancer (DTC) has become one of the most frequently diagnosed malignancies, especially among women and young adults (Davies & Welch 2014). The outcomes are generally very good: disease recurrence rates are low (Durante *et al.* 2013), and survival rates are excellent (Tuttle *et al.* 2017a). Evidence-based management is crucial to avoid overtreatment of these low-risk tumors, which can reduce quality of life and yet identify accurately those requiring more aggressive therapy. Several staging systems have been generated to inform DTC management. One of the most widely used is the tumor-node-metastasis (TNM) classification elaborated by the American Joint Committee on Cancer (AJCC), which allows to predict the risk of cancer-related death. The 8th edition of the AJCC staging system for thyroid cancer (AJCC-8) was recently published (Tuttle *et al.* 2017b) and is scheduled to be implemented on 1 January 2018. Revision of the system was undertaken to address several specific limitations identified in the 7th edition (AJCC-7), which has been in use since 2009 (Tuttle *et al.* 2017a,b). The main changes (described in detail below and summarized in Table 1) are as follows: (1) an increase in the age threshold for defining high risk of thyroid cancer-related death and (2) a decrease in the unfavorable prognostic significance attributed to certain findings (i.e., cervical lymph node metastases and microscopic extrathyroidal extension (ETE), which has been re-defined to include only invasion of the perithyroidal muscle).

To assess the impact of transitioning to the new AJCC-8 in terms of stage distribution and prevalence of each stage class, we analyzed data extracted from the web-based database of the Italian Thyroid Cancer Observatory (ITCO) (www.itcofoundation.org), a network of thyroid cancer centers (including primary and tertiary centers) located throughout Italy. The database includes prospectively updated, observational data provided by ITCO member centers on patients consecutively diagnosed with thyroid

cancer since 2013 (Lamartina *et al.* 2017). Cases included in our study met all the following criteria: (1) histological diagnosis of thyroid cancer of follicular origin; (2) date of diagnosis between 1 January 2013 and 1 March 2017; (3) complete data on primary tumor pathology, including minimal ETE, and initial treatment.

The selected cohort analyzed included 1765 patients, 76% of whom were females. The median age at diagnosis was 48 years (range: 10–87). Total thyroidectomy (or lobectomy+completion thyroidectomy) was performed in 1727 (98%) cases and followed by radioiodine remnant ablation in 954 (55%). Neck dissection was performed in 711 (40%) of the 1765 patients. Most of the tumors (n=1657, 94%) were papillary thyroid cancers; the remaining 108 (6%) were follicular or Hürthle cell carcinomas. Estimated risks of recurrence calculated according to the criteria recommended in 2015 by the American Thyroid Association were low in 1046 (59%), intermediate in 612 (35%) and high in 107 (6%) of the cases. Microscopic ETE was found in 410 (23%), but only 40 (2%) of these patients had gross invasion of the strap muscles (sternohyoid, sternothyroid, thyroidhyoid and/or omohyoid muscles). Lymph node status for the 711 patients who underwent lymph node dissection was as follows: pN0 (no metastasis) in 338 (19%); pN1a (central compartment metastases) in 221 (12%) and pN1b (lateral compartment metastases) 152 (9%). Distant metastases were found in 32 (1.8%) patients.

As noted above, in the AJCC-8, the age threshold for high risk of disease-specific mortality was raised from 45 years – the median age at diagnosis in several published series – to 55 years (Nixon *et al.* 2016). This change increases the proportion of relatively young patients whose mortality risk can be defined solely on the basis of the absence or presence of distant metastases (stages I and II, respectively) (Table 1). As shown in Fig. 1A, the percentage of patients classified as ‘younger’ in our

Table 1 AJCC TNM staging: 7th and 8th editions.

7th edition	Age <45 years			8th edition	Age <55 years		
I	Any tumor size	Any lymph node status	Absence of distant metastases (M0)	I	Any tumor size	Any lymph node status	Absence of distant metastases (M0)
II	Any tumor size	Any lymph node status	Presence of distant metastases (M1)	II	Any tumor size	Any lymph node status	Presence of distant metastases (M1)
7th edition	Age ≥45 years			8th edition	Age ≥55 years		
I	Tumor of ≤2 cm limited to the thyroid (T1)	Absence of lymph node metastases (Nx/N0)	Absence of distant metastases (M0)	I	Tumor of ≤4 cm limited to the thyroid (T2)	Absence of lymph node metastases (Nx/N0)	Absence of distant metastases (M0)
II	Tumor of ≤4 cm limited to the thyroid (T2)	Absence of lymph node metastases (Nx/N0)	Absence of distant metastases (M0)	II	Tumor of any size with lymph node metastases (N1) or with gross extrathyroidal extension invading only strap muscles (sternohyoid, sternothyroid, thyroidhyoid, omohyoid) with/without lymph node metastases (T3b)		Absence of distant metastases (M0)
III	Tumor of any size with lymph node metastases of the central compartment (N1a) or with minimal extrathyroid extension (T3) with/without lymph node metastases of the central compartment		Absence of distant metastases (M0)	III	Gross extrathyroidal extension invading subcutaneous soft tissues, larynx, trachea, esophagus, or recurrent laryngeal nerve (T4a)	Any lymph node status	Absence of distant metastases (M0)
IVa	Tumor of any size with lymph node metastases of the lateral compartment (N1b) or with gross extrathyroidal extension invading subcutaneous soft tissues, larynx, trachea, esophagus, or recurrent laryngeal nerve (T4a) with/without lymph node metastases		Absence of distant metastases (M0)	IVa	Gross extrathyroidal extension invading prevertebral fascia or encasing the carotid artery or mediastinal vessels (T4b)	Any lymph node status	Absence of distant metastases (M0)
IVb	Gross extrathyroidal extension invading prevertebral fascia or encasing the carotid artery or mediastinal vessels (T4b)	Any lymph node status	Absence of distant metastases (M0)	IVb	Any tumor size	Any lymph node status	Presence of distant metastases (M1)
IVc	Any tumor size	Any lymph node status	Presence of distant metastases (M1)	–	–	–	–

cohort rose from 40% (698/1765) with the AJCC-7 to 65% (1150/1765) with the AJCC-8, but in both cases, the proportion of patients with distant metastases (i.e., those classified as stage II) was identically low (2%). Given the overall increase in the age of the 'younger' stage II patients, their estimated disease-specific survival at 10 years (DSS-10) drops from 95–99% to 85–95% (Kim *et al.* 2017b, Tuttle *et al.* 2017a,b).

Among the 615 patients classified by both the AJCC-7 and the AJCC-8 as 'older' (Fig. 1B), 193 (31%) originally

classified as stage III or IVa were re-classified as stage II, raising the proportion of patients with stages I–II DTC from 64% to 94%. Over half of the downstagings (112/193, 58%) involved patients whose previous T3 classification had been based solely on the presence of microscopic ETE, which has no effect on the T category or overall disease stage in the AJCC-8 (Tuttle *et al.* 2017b). In the remaining patients, the AJCC-7 classification as stage III or IV had been based on the presence of regional metastases alone (45/193, 23%), which no longer necessitates assignment

Figure 1

DTC stage distributions in the ITCO cohort based on the 7th and 8th editions of the AJCC system. (A) The younger subcohorts defined by the AJCC-7 and AJCC-8 comprised 698 patients aged <45 years and 1150 patients <55 years, respectively. In both subcohorts, 98% of the patients were classified as stage I. (B) The older subcohorts defined by the AJCC-7 and AJCC-8 comprised 1067 patients aged >45 years and 615 patients >55 years, respectively. Restaging with the AJCC-8 increased the percentages of patients with stage I (from 56 to 64%) or II (from 8 to 31%) disease.

to stage III (Tuttle *et al.* 2017b), or regional metastases plus microscopic ETE (18/193, 9%). It should be noted that, differently from the AJCC-7, the ITCO database has always classified level VII lymph node metastases as central neck node lesions. This reflects the well-known difficulties in distinguishing levels VI and VII and is consistent with the revised definitions adopted in the AJCC-8 (Tuttle *et al.* 2017b). Since the same classification was also used for our AJCC-7 staging, some cases that met the criteria for AJCC-7 stage IVa may have been erroneously reported herein as stage III. As for the entire cohort, application of the AJCC-8 criteria downstaged 477 (27%) of the 1765 DTC patients. As a result, the estimated risk for 10-year disease-specific mortality was <15% for almost all the patients. Higher estimated risks (>40%) were restricted to 2% (31/1765) of the patients who were 55 or older and had gross ETE (T4), with or without distant metastases (stages III and IV). Importantly, the mortality risk was not always paralleled by the likelihood of recurrence. The risk of recurrence, as defined by the American Thyroid Association, was rated intermediate in 25% of AJCC-8 stage I patients and 90% of AJCC-8 stage II patients aged ≥ 55 , and high risks of recurrence were found in 3% and 4% of these groups, respectively, owing to the presence of unfavorable histologic findings (i.e., widely invasive follicular and Hürthle cell cancer with foci of vascular invasion).

The AJCC-8 stage distribution for our DTC cohort resembles those reported for retrospectively analyzed cohorts (Kim *et al.* 2017a,b), where the downstaging affected an even larger subset of patients (38%). Compared with the ITCO cohort, the Korean cohorts included higher percentages of patients with regional metastasis or microscopic ETE (56% and 32–40%, respectively), which might reflect a referral-center-selection bias. Microscopic ETE reporting can be unreliable as it is often variable even between skilled pathologists (Su *et al.* 2016). However,

a far better agreement was found for the presence of gross ETE with invasion of perithyroidal muscles compared with invasion of perithyroidal fat only. The extensive downstaging effect of the AJCC-8 was intentional: the TNM staging system assesses the risk of DTC-related death, which has proved to be very low for most patients. Indeed, the results of retrospective cohort analyses confirm that the AJCC-8 provides more accurate estimates of DTC patients' DSS (Kim *et al.* 2017a,b, Pontius *et al.* 2017, Tuttle *et al.* 2017a). From a practical standpoint, application of the AJCC-8 criteria can be expected to simplify the staging process for most DTC patients, who will now be classified as 'younger'. It will also markedly reduce the number of patients with a substantial mortality risk (those with stages III–IV disease, where the DSS-10 is <50–70%). This new tool, together with the risk of recurrence stratification tools, should facilitate clinicians' attempts to provide more cost-effective management of DTCs, and this improvement should also have benefits in terms of their patients' quality of life.

Livia Lamartina^{1,*}

Giorgio Grani^{1,*}

Emanuela Arvat²

Alice Nervo²

Maria Chiara Zatelli³

Roberta Rossi³

Ef시오 Puxeddu⁴

Silvia Morelli⁴

Massimo Torlontano⁵

Michela Massa⁵

Rocco Bellantone⁶

Alfredo Pontecorvi⁷

Teresa Montesano⁸

Loredana Pagano⁹

Lorenzo Daniele¹⁰

Laura Fugazzola^{11,12}

Graziano Ceresini¹³Rocco Bruno¹⁴Ruth Rossetto¹⁵Salvatore Tumino¹⁶Marco Centanni¹⁷Domenico Meringolo¹⁸Maria Grazia Castagna¹⁹Domenico Salvatore²⁰Antonio Nicolucci²¹Giuseppe Lucisano²¹Sebastiano Filetti¹Cosimo Durante¹¹⁶Dipartimento di Medicina Clinica e Sperimentale,
Università di Catania, Catania, Italy¹⁷Dipartimento di Scienze e Biotechnologie Medico-Chirurgiche,
Università di Roma Sapienza, Latina, Italia¹⁸Unità Operativa Semplice Dipartimentale, Ospedale
Bentivoglio, Ausl Bologna, Bologna, Italy¹⁹Dipartimento di Scienze Mediche, Chirurgiche
e Neurologiche, Università di Siena, Siena, Italy²⁰Dipartimento di Medicina Clinica e Chirurgia, Università
degli Studi di Napoli 'Federico II', Napoli, Italy²¹Center for Outcomes Research and Clinical Epidemiology,
Pescara, Italy¹Dipartimento di Medicina Interna e Specialità Mediche,
Università di Roma Sapienza, Roma, Italy²Dipartimento di Scienze Mediche, Università di Torino,
Torino, Italy³Unità Operativa di Endocrinologia e Malattie del Ricambio,
Azienda Ospedaliera Universitaria di Ferrara, Ferrara, Italy⁴Dipartimento di Medicina, Università di Perugia,
Perugia, Italy⁵Dipartimento di Scienze Mediche, Ospedale Casa Sollievo
della Sofferenza-IRCCS, San Giovanni Rotondo, Italy⁶Cattedra di Chirurgia Endocrina, Area di Endocrinologia
e Malattie Metaboliche, Fondazione Policlinico Universitario
Agostino Gemelli, Università Cattolica del Sacro Cuore,
Roma, Italy⁷Cattedra di Endocrinologia, Area di Endocrinologia
e Malattie Metaboliche, Fondazione Policlinico Universitario
Agostino Gemelli, Università Cattolica del Sacro Cuore,
Roma, Italy⁸Unità di Medicina Nucleare, Università di Roma Sapienza,
Roma, Italy⁹Unità di Endocrinologia, Dipartimento di Medicina
Traslationale, Università del Piemonte Orientale,
Novara, Italy¹⁰Unità di Patologia, Ospedale Mauriziano Umberto I di
Torino, Torino, Italy¹¹Divisione di Malattie Endocrine e Metaboliche, Istituto
Auxologico Italiano IRCCS, Milano, Italy¹²Dipartimento di Fisiopatologia Medico-Chirurgica e dei
Trapianti, Università degli Studi di Milano, Milano, Italy¹³Dipartimento di Medicina e Chirurgia, Università di Parma,
Parma, Italy¹⁴Unità di Endocrinologia, Ospedale di Tinchi-Pisticci,
Matera, Italy¹⁵Divisione di Endocrinologia, Diabetologia e Metabolismo,
Dipartimento di Scienze Mediche, Ospedale Molinette, A.O.U.
Città della Salute e della Scienza di Torino, Università di
Torino, Torino, Italy(Correspondence should be addressed to S Filetti:
sebastiano.filetti@uniroma1.it)*(L Lamartina and G Grani contributed equally to this
work)**Declaration of interest**The authors declare that there is no conflict of interest that could be
perceived as prejudicing the impartiality of this article.**Funding**

This paper was funded by the Umberto Di Mario Foundation.

Author contribution statement Livia Lamartina and Giorgio Grani: data management and interpretation,
analysis and writing. Antonio Nicolucci and Giuseppe Lucisano: statistical
analysis and data interpretation. Emanuela Arvat, Alice Nervo, Maria
Chiara Zatelli, Roberta Rossi, Efisio Puxeddu, Silvia Morelli, Massimo
Torlontano, Michela Massa, Rocco Bellantone, Alfredo Pontecorvi,
Teresa Montesano, Loredana Pagano, Lorenzo Daniele, Laura Fugazzola,
Graziano Ceresini, Rocco Bruno, Ruth Rossetto, Salvatore Tumino, Marco
Centanni, Domenico Meringolo, Maria Grazia Castagna and Domenico
Salvatore: data collection and analysis, and manuscript editing. Cosimo
Durante and Sebastiano Filetti: concept, data interpretation, analysis and
manuscript editing.**Acknowledgements**G G and L L contributed to this paper as recipients of the PhD program of
Biotechnologies and Clinical Medicine of the University of Rome, Sapienza.
Medical editing services were provided by M E Kent and funded by the
Umberto Di Mario Foundation. The authors thank all the members of the
ITCO foundation: Ancona: Alessia Smerilli, Augusto Taccaliti. Bari: Vincenzo
Triggiani, Giuseppina Renzulli. Brescia: Maria Beatrice Panarotto. Catania:
Simona Quartararo, Dario Tumino. Ferrara: Ettore Degli Uberti. La Spezia:
Laura Camerieri, Mario Cappagli. Latina: Maria Giulia Santaguida.
Livorno: Daniele Barbaro, Paola Lapi. Matera: Antonella Carbone. Milano:
Università degli Studi di Milano: Carla Colombo, Simone De Leo, Luca
Persani; Univesrità Humanitas di Milano: Andrea Lania. Napoli: Università
degli Studi di Napoli 'Federico II': Tommaso Porcelli. Istituto Nazionale
Tumori ITCCS 'Fondazione G. Pascale': Maria Grazia Chiofalo, Luciano
Pezzullo. Novara: Gianluca Aimaretti, Chiara Mele. Padova: Caterina Mian,

Simona Censi. Palermo: Andrea Attard, Marco Attard, Adele Maniglia. Parma: Michela Marina. Pisa: Nadia Caraccio, Poupak Fallahi, Filippo Niccolai, Giuseppe Pasqualetti, Armando Patrizio. Roma: Policlinico Universitario Campus Biomedico: Anna Crescenzi, Angelo Lauria, Rossella Del Toro; Università Cattolica del Sacro Cuore: Francesca Ianni, Celestino Pio Lombardi, Valeria Loiacono, Manuela Nestola, Emanuela Traini; Università di Roma Sapienza: Laura Ciotti, Cristiano Lomonaco, Salvatore Monti, Sara Nazzarena Morgante, Giuseppe Ronga, Maria Segni, Serena Elisa Tempera, Vincenzo Toscano. San Giovanni Rotondo: Umberto Crocetti, Leonardo D'Aloiso. Siena: Raffaella Forleo, Fabio Maino, Furio Pacini. Torino: Ospedale Mauriziano Umberto I di Torino: Paolo Limone. Università di Torino: Stefania Corvisieri, Francesco Felicetti, Marco Gallo, Ezio Ghigo, Alessandro Piovesan. Verona: Chiara Conforti, Amalia Robustelli, Giovanna Spiazzi.

References

- Davies L & Welch HG 2014 Current thyroid cancer trends in the United States. *JAMA Otolaryngology: Head and Neck Surgery* **140** 317–322. (<https://doi.org/10.1001/jamaoto.2014.1>)
- Durante C, Montesano T, Torlontano M, Attard M, Monzani F, Tumino S, Costante G, Meringolo D, Bruno R, Trulli F, *et al.* 2013 Papillary thyroid cancer: time course of recurrences during postsurgery surveillance. *Journal of Clinical Endocrinology and Metabolism* **98** 636–642. (<https://doi.org/10.1210/jc.2012-3401>)
- Kim M, Kim WG, Oh HS, Park S, Kwon H, Song DE, Kim TY, Shong YK, Kim WB, Sung TY, *et al.* 2017a Comparison of the seventh and eighth editions of the American Joint Committee on cancer/union for international cancer control tumor-node-metastasis staging system for differentiated thyroid cancer. *Thyroid* **27** 1149–1155. (<https://doi.org/10.1089/thy.2017.0050>)
- Kim TH, Kim YN, Kim HI, Park SY, Choe JH, Kim JH, Kim JS, Oh YL, Hahn SY, Shin JH, *et al.* 2017b Prognostic value of the eighth edition AJCC TNM classification for differentiated thyroid carcinoma. *Oral Oncology* **71** 81–86. (<https://doi.org/10.1016/j.oraloncology.2017.06.004>)
- Lamartina L, Durante C, Lucisano G, Grani G, Bellantone R, Lombardi CP, Pontecorvi A, Arvat E, Felicetti F, Zatelli MC, *et al.* 2017 Are evidence-based guidelines reflected in clinical practice? An analysis of prospectively collected data of the Italian Thyroid Cancer Observatory (ITCO#1). *Thyroid* **27** 1490–1497. (<https://doi.org/10.1089/thy.2017.0299>)
- Nixon IJ, Wang LY, Migliacci JC, Eskander A, Campbell MJ, Aniss A, Morris L, Vaisman F, Corbo R, Momesso D, *et al.* 2016 An international multi-institutional validation of age 55 years as a cutoff for risk stratification in the AJCC/UICC staging system for well-differentiated thyroid cancer. *Thyroid* **26** 373–380. (<https://doi.org/10.1089/thy.2015.0315>)
- Pontius LN, Oyekunle TO, Thomas SM, Stang MT, Scheri RP, Roman SA & Sosa JA 2017 Projecting survival in papillary thyroid cancer: a comparison of the seventh and eighth editions of the American Joint Commission on cancer/union for international cancer control staging systems in two contemporary national patient cohorts. *Thyroid* **27** 1408–1416. (<https://doi.org/10.1089/thy.2017.0306>)
- Su HK, Wenig BM, Haser GC, Rowe ME, Asa SL, Baloch Z, Du E, Faquin WC, Fellegara G, Giordano T, *et al.* 2016 Inter-observer variation in the pathologic identification of minimal extrathyroidal extension in papillary thyroid carcinoma. *Thyroid* **26** 512–517. (<https://doi.org/10.1089/thy.2015.0508>)
- Tuttle RM, Haugen B & Perrier ND 2017a Updated American Joint Committee on cancer/tumor-node-metastasis staging system for differentiated and anaplastic thyroid cancer (eighth edition): what changed and why? *Thyroid* **27** 751–756. (<https://doi.org/10.1089/thy.2017.0102>)
- Tuttle M, Morris LF, Haugen B, Shah J, Sosa JA, Rohren E, Subramaniam RM, Hunt JL & Perrier ND 2017b Thyroid-differentiated and anaplastic carcinoma (Chapter 73). In *AJCC Cancer Staging Manual*, 8th edition. Eds MB Amin, SB Edge, F Greene, D Byrd, RK Brookland, MK Washington, CC Compton, KR Hess, DC Sullivan, JM Jessup, *et al.* New York, NY, USA: Springer International Publishing.

Received in final form 24 November 2017

Accepted 29 November 2017

Accepted Preprint published online 30 November 2017