THEMATIC REVIEW

Na⁺/I[−] symporter expression, function, and regulation in non-thyroidal tissues and impact on thyroid cancer therapy

Sissy M Jhiang^{1,2} and Jennifer A Sipos²¹Department of Physiology and Cell Biology, The Ohio State University Comprehensive Cancer Center, The Ohio State University, Columbus, Ohio, USA²Department of Internal Medicine, Division of Endocrinology and Metabolism, The Ohio State University Comprehensive Cancer Center, Columbus, Ohio, USACorrespondence should be addressed to S Jhiang: jhiang.1@osu.edu

This paper is part of a thematic review section celebrating 80 Years of the Use of Radioiodine. The guest editor for this section was Christopher John McCabe.

Abstract

For the past 80 years, radioiodine (¹³¹I) has been used to ablate thyroid tissue not removed by surgery or to treat differentiated thyroid cancer that has metastasized to other parts of the body. However, the Na⁺/I[−] symporter (NIS), which mediates active iodide uptake into thyroid follicular cells, is also expressed in several non-thyroidal tissues. This NIS expression permits ¹³¹I accumulation and radiation damage in these non-target tissues, which accounts for the adverse effects of radioiodine therapy. We will review the data regarding the expression, function, and regulation of NIS in non-thyroidal tissues and explain the seemingly paradoxical adverse effects induced by ¹³¹I, the self-limited gastrointestinal adverse effects in contrast to the permanent salivary dysfunction that is seen after ¹³¹I therapy. We propose that prospective studies are needed to uncover the time-course of pathological processes underlying development and progression or ultimate resolution of ¹³¹I-induced salivary ductal obstruction and nasolacrimal duct obstruction. Finally, preventive measures and early therapeutic interventions that can be applied potentially to eliminate or alleviate long-term radioiodine adverse effects will be discussed.

Key Words

- ▶ Na⁺/I[−] symporter
- ▶ non-thyroid tissues
- ▶ radioiodine adverse effect
- ▶ salivary ductal obstruction
- ▶ nasolacrimal duct obstruction

Endocrine-Related Cancer
(2021) **28**, T167–T177

Introduction

The thyroid gland is the only tissue in the body that utilizes iodine (Berman *et al.* 1968). Radioiodine (¹³¹I) ablation is, therefore, a logical and highly specific targeted therapy for remnant ablation after total thyroidectomy and/or for treatment of metastatic lesions derived from well-differentiated, follicular cell-derived thyroid cancer (Mazzaferri & Jhiang 1994). After administration, radioiodine taken into the thyroid is retained via iodine organification (Berman *et al.* 1968). The majority of the

remaining radioiodine not taken up by thyroid tissues is eliminated in the urine within 2 days (Berman *et al.* 1968). For the first 2 days after radioiodine administration, notable radioiodine accumulation is also found in the salivary glands (Myant 1960) and stomach (Berman *et al.* 1968), which leads to the early complications of sialadenitis (Goolden *et al.* 1957, Van Nostrand *et al.* 1986) and gastrointestinal symptoms (Van Nostrand *et al.* 1986). Gastrointestinal symptoms typically resolve quickly,

while salivary gland dysfunction may recur and can persist in some patients (Van Nostrand *et al.* 1986, Clement *et al.* 2015). Additional adverse effects of radioiodine therapy include xerostomia, bone marrow suppression, gonadal damage, dry eye(s), nasolacrimal duct obstruction, and secondary cancer (Fard-Esfahani *et al.* 2014, Clement *et al.* 2015). While the incidence and severity of the aforementioned adverse effects are generally correlated with the cumulative radioactivity administered (Clement *et al.* 2015, Hollingsworth *et al.* 2016), factors contributing to variations in incidence and severity of radioiodine adverse effects among individual patients are yet to be delineated. Molecular cloning of the Na⁺/I⁻ symporter (NIS) that mediates active iodide uptake into thyroid follicular cells (Dai *et al.* 1996, Smanik *et al.* 1996), subsequently developing NIS antibodies (Levy *et al.* 1997, Jhiang *et al.* 1998), has made it possible to investigate NIS expression, function, and regulation in non-thyroidal tissues. A better understanding of NIS expression and modulation in non-thyroid tissues may uncover the underlying pathological process of ¹³¹I-induced adverse effects and allow us to devise better strategies to prevent ¹³¹I-induced adverse effects.

NIS expression in non-thyroidal tissues

NIS expression in thyroidal and non-thyroidal tissues may be detected by RT-quantitative PCR (RT-qPCR) or Northern blot analysis at the mRNA level and by immunohistochemical (IHC) staining or Western blot analysis at the protein level. RT-qPCR is the most sensitive method that can detect minute quantities of NIS mRNA. However, the physiological or functional significance of such an infinitesimal detection is unclear. IHC staining is superior in terms of detecting NIS expression restricted to a specific cell type and in revealing plasma membrane localization of NIS, where NIS mediates radioiodine uptake (Jhiang *et al.* 1998, Wapnir *et al.* 2003). However, IHC staining results can be compromised by cross-reactivity to non-NIS proteins due to variations in NIS antibody specificity and stringency of antibody incubation and washing conditions. Both Northern and Western blot analyses suffer from limited detection sensitivity but may reveal alternative spliced transcripts and post-translational modifications, respectively. For example, with those tissues composed of multiple cell types, NIS expression may not be detected by Northern or Western blot analyses, particularly if its expression is restricted to a specific cell type that comprises only a minor component of the overall tissue composition (Spitzweg *et al.* 1998).

Taken together, RT-qPCR accompanied by IHC staining using a NIS antibody of high specificity can be informative for investigating NIS expression in non-thyroidal tissues at different developmental stages and in various pathological conditions.

According to the Human Protein Atlas website (<https://www.proteinatlas.org/ENSG00000105641-SLC5A5/tissue>), thyroid, salivary gland, and stomach are tissues with significant NIS detection at both mRNA and protein levels. The GeneCards website (<https://www.genecards.org/cgi-bin/carddisp.pl?gene=SLC5A5&keywords=SLC5A5>) indicates that RNAseq analysis has identified numerous tissues expressing NIS mRNA including blood cells, lymph nodes, brain, cortex, cerebellum, spinal cord, tibial nerve, heart, artery, skeletal muscle, small intestine, colon, adipocyte, kidney, liver, lung, spleen, stomach, esophagus, bladder, pancreas, salivary gland, adrenal gland, pituitary, breast, skin, ovary, uterus, prostate and testis. Among these tissues, NIS mRNA level was the highest in the stomach, followed by the thyroid, salivary glands, and cerebellum.

NIS expression in salivary gland tissues

It was recognized that NIS is expressed in the salivary glands before NIS molecule cloning. In patients with autosomal recessive, congenital dysmaturational goiters, the abnormal saliva/serum iodide ratio, as well as the lack of ^{99m}TcO₄⁻ uptake in both thyroid and salivary glands were used to reveal the underlying iodide transport defect (Wolff 1983). However, the localization of NIS expression in salivary ductal cells rather than acinar cells was not discovered until the development of NIS antibodies. IHC staining demonstrates that NIS is highly expressed in the basolateral membranes of the striated ducts (Jhiang *et al.* 1998) (Fig. 1A) whereas weakly expressed in intercalated and excretory ductal cells (La Perle *et al.* 2013). Accordingly, ¹³¹I is concentrated from blood circulation into saliva in the ductal lumen mainly from salivary striated ductal cells. In sialoadenitis, NIS expression in striated ducts has been demonstrated to be heterogeneously decreased or absent (La Perle *et al.* 2013) (Fig. 1B), suggesting that inflammation decreases NIS expression in striated ductal cells. This pathologic process translates well with clinical observations of decreased uptake and impaired clearance of ^{99m}TcO₄⁻ that is seen after RAI therapy (Mandel & Mandel 2003) (Fig. 1C). The ¹³¹I-induced inflammatory infiltrate causes increased periductal pressure with duct constriction and sialadenitis, which usually manifests as bilateral parotid swelling and pain with decreased salivary flow (Goolden *et al.* 1957, Mandel & Mandel 2003). For some patients, resolution of the inflammatory process

Figure 1

NIS expression and function in normal and inflamed salivary glands. (A) NIS immunostaining (brown color) is detected in the basolateral membranes of the normal striated ducts; (B) NIS immunostaining (brown color) is heterogeneously decreased or absent in inflamed striated ductal cells during sialoadenitis; and (C) Technetium-99m pertechnetate (TPT) time activity graph shows normal uptake and secretory clearance of the TPT radioisotope in the functioning left parotid (curve of green color). The inflamed right parotid (curve of blue color) had minimal uptake and no secretory clearance, indicating reduced salivary NIS expression and salivary ductal obstruction, respectively. Lemon candy was given at frame 20 to stimulate secretory clearance. (C) Modified, with permission, from Mandel SJ & Mandel L, Radioactive iodine and the salivary glands, *Thyroid*, 2003, volume 13, pages 265–271.

and improvement in symptoms occurs within days, while other patients experience subclinical inflammation with delayed symptoms (Mandel & Mandel 2003). The latter group is more likely to suffer from recurrent/persistent sialadenitis that eventually leads to permanent xerostomia with progressive susceptibility to dental caries and periodontal disease (Mandel & Mandel 2003). It is important to note that both acute periductal inflammation and chronic sclerosis in the salivary glands may be seen in thyroid cancer patients who have been treated with ^{131}I therapy.

Our study of ^{123}I single-photon-emission CT/anatomic imaging of CT (SPECT/CT) images from 50 thyroid cancer patients prior to radioiodine therapy showed that the ratio of ^{123}I cumulative radioactivity at 24 h after administration between parotid and submandibular glands is 2.38 ± 0.19 (La Perle *et al.* 2013). However, the corresponding ratio of ^{123}I accumulation normalized by volume of interest is 1.19 ± 0.06 . The percentage of NIS-positive striated ductal cells in submandibular salivary glands was statistically greater than in parotid salivary glands, suggesting a higher clearance rate of ^{131}I in submandibular than parotid salivary glands. Indeed, approximately 65–70% of saliva in the oral cavity is produced by the submandibular glands, even though they are much smaller than the parotid glands. These findings are consistent with the recent study showing that damage to the parotid glands is more common than damage to submandibular salivary tissue among radioiodine-treated patients when examined by neck ultrasound (Horvath *et al.* 2020).

We also demonstrated that most ductal salivary gland tumors do not express NIS. The exception is Warthin's tumors of striated duct origin which exhibit consistent and intense NIS staining and avid radioactive iodine uptake (La Perle *et al.* 2013). Taken together, NIS expression in salivary ductal cells appears to be mainly modulated by ductal cell differentiation, and NIS expression is retained in salivary tumors derived from striated ducts.

NIS expression in gastrointestinal tissues

In the gastrointestinal tract, NIS is expressed in the basolateral membrane of gastric mucin-producing cells (Wapnir *et al.* 2003). However, in the small intestine, NIS is functionally expressed on the apical surface of enterocytes (Nicola *et al.* 2009), suggesting that iodine from food sources as well as iodine concentrated in gastric juice is likely to be absorbed and reabsorbed, respectively, via NIS at the apical membrane of the small intestine into the blood circulation. The gastrointestinal epithelium is a constitutively developing tissue, constantly differentiating from stem cells in a progenitor pool throughout the life of the organism (de Santa Barbara *et al.* 2003). The high turnover rate of the gastric mucosa and enterocytes may explain why gastrointestinal symptoms generally resolve quickly after radioiodine therapy.

The pathogenesis of gastric cancer has been associated with abnormal patterns of gastric differentiation and with chronic tissue injury (Mills & Shivdasani 2011).

Altorjay *et al.* (2007) found that NIS staining was observed in gastric mucin-producing cells in gastritis, both in the presence and absence of *Helicobacter pylori*. However, NIS expression was absent in gastric cancer, independent of its histological type. They also observed that focal NIS staining in the direct vicinity of gastric tumors increased gradually and became linear with escalating distance from the tumor, suggesting that loss of NIS expression precedes microscopically identifiable morphological changes. Recently, Shiozaki *et al.* (2019) examined NIS expression by IHC staining in 145 primary gastric cancer samples and showed that 37% of the samples were NIS positive, yet NIS immunostaining was detected in the cytoplasm and non-polarized cell membrane. They further demonstrated that strong NIS expression was associated with a poor prognosis in gastric cancer. A separate study reported a similar prognostic role of NIS expression in gastric cancers (Mishra & Shrivastava 2020).

In the normal gastrointestinal tract, NIS expression is restricted to terminally differentiated cells with a rapid turnover rate. For NIS-negative gastric cancer, one could interpret that malignant transformation suppresses NIS expression or occurs in NIS-negative gastric precursor cells at a less-differentiated stage. For NIS-positive gastric cancer, it is possible that signaling nodes and/or transcription factors that initiate NIS expression in gastric mucosa are aberrantly activated during malignant transformation or malignant transformation occurs after precursor cells are committed to the differentiated gastric mucosa.

NIS expression in breast tissues

NIS expression and radioiodide uptake in breast tissues are at the highest levels during active lactation (Cho *et al.* 2000, Tazebay *et al.* 2000). NIS proteins are detected at the basolateral membrane of breast ductal epithelium. Iodide is transported from the mother's blood circulation to the ductal lumen where milk is accumulated. This process is critical to ensure sufficient iodide is provided to the newborn for thyroid hormone synthesis. The NIS protein level in breast tissue declines significantly during the post-lactational involution process until it is at an undetectable level in non-lactating normal breast tissues (Cho *et al.* 2000).

Incidental findings of radioiodine uptake in non-breastfeeding women have been reported (Hammami & Bakheet 1996), yet the underlying reason(s) remains to be elucidated. Among pathological breast tissues, our unpublished data and others (Kilbane *et al.* 2000, Berger *et al.* 2006, Ryan *et al.* 2011) detected a high level of NIS

expression at the basolateral membrane of fibroadenoma, a benign breast tumor which may change in size during pregnancy, breastfeeding, or while taking hormone replacement therapy. Fibroadenoma is one of the most common benign tumors of the breast in women under 30 years of age. NIS-positive fibroadenomas may account for some of the incidental findings of radioiodine uptake in non-breastfeeding women (Hammami & Bakheet 1996, Kim *et al.* 2017).

Based on communication with our previous collaborator, Dr Juha Kononen at the National Human Genome Research Institute, our analysis of NIS IHC staining in tissue microarray composing of 650 breast tumors showed that 17% of 650 breast tumors had strong plasma membrane NIS staining that did not correlate with the presence or absence of estrogen receptor, progesterone receptor, nor human EGF receptor 2 (HER2) (unpublished data). In contrast, Wapnir *et al.* (2003) reported a significant number of breast cancers with cytoplasmic NIS immunostaining. It is possible that the detectability of plasma membrane NIS and cytoplasmic NIS is epitope-dependent and can vary among NIS antibodies. However, the identity of cytoplasmic NIS immunostaining in both thyroid and breast cancers has been questioned (Peyrottes *et al.* 2009). If possible, an additional method of NIS detection, such as RT-qPCR with an acceptable Ct value (cycle threshold), needs to be applied to confirm NIS identity in breast cancers with exclusively cytoplasmic NIS staining. Of interest, Smith *et al.* (Smith *et al.* 2009, 2013) reported that inhibition of PTTG binding factor (PBF) increases NIS plasma membrane localization with improved radioiodine uptake in thyroid and other tumors. Recently, Fletcher *et al.* (2020) reported that an inhibitor for valosin-containing protein (VCP) significantly increases NIS cell surface localization and results in a marked increase in radioiodine uptake activity in both mouse and human thyroid models.

NIS expression in lacrimal sac and nasolacrimal duct

Nasolacrimal duct obstruction has been associated with high-dose radioiodine therapy (Kloos *et al.* 2002, Ali 2016, da Fonseca *et al.* 2016). Morgenstern *et al.* (2005) showed that NIS mRNA was detected in normal lacrimal sacs via RT-PCR and NIS immunostaining was detected at the basolateral membrane of the stratified columnar epithelial cells in the lacrimal sac (Fig. 2A) and nasolacrimal duct (Fig. 2B). In contrast, NIS immunostaining was not detected in the lacrimal glands (Fig. 2C), Wolfring or Krause glands, conjunctiva, canaliculus, nor nasal mucosa. In lacrimal

Figure 2

NIS expression in normal lacrimal drainage tissues. NIS immunostaining (brown color) is detected at the basolateral membrane of the stratified columnar epithelial cells in (A) the lacrimal sac, and (B) the nasolacrimal duct. NIS immunostaining is absent in (C) lacrimal glands. NIS expression was not investigated in (D) lacrimal excretory ducts (red arrow).

sacs removed from radioiodine-treated patients, NIS-expressing cells were absent, and marked fibrosis was noted at sites of the stratified columnar epithelial cells (Morgenstern *et al.* 2005). Morgenstern *et al.* (2005) concluded that ^{131}I accumulation at the lacrimal sac and nasolacrimal duct is responsible for the development of nasolacrimal drainage system obstruction, as patients diagnosed with nasolacrimal obstruction typically do not demonstrate signs of proximal canaliculus stenosis. Instead, the association of dry eye with radioiodine therapy could be attributed to an increased risk of eye infection or inflammation from improper tear drainage (Morgenstern *et al.* 2005). As illustrated in Fig. 2D, tears produced by the lacrimal glands are secreted into excretory ducts which empty into the superior conjunctival fornix and then spread over to the cornea due to blinking. However, NIS expression in the excretory ducts has not been examined. If NIS is expressed in the excretory ducts, it would explain the detection of radioactive tears after ^{131}I administration, and ^{131}I -induced excretory duct obstruction may lead to dry eyes. Since dry eyes can be triggered by multiple factors, such as aging, medications, allergy, menopause, etc., the role of ^{131}I in the development of dry eyes in ^{131}I -treated patients is uncertain.

NIS expression in other non-thyroid tissues

Using a very sensitive detection method, RT-PCR followed by Southern hybridization, Spitzweg *et al.* (1998) reported

that NIS expression was detected in the pituitary gland, pancreas, testis, mammary gland, gastric mucosa, prostate and ovary, adrenal gland, heart, thymus, and lung. Wapnir *et al.* (2003) reported apical membrane NIS staining in placenta cytotrophoblasts and renal distal and collecting tubules, as well as NIS intracellular immunostaining in many normal tissues including bladder mucosa, colonic mucosal cells, bile duct, etc. However, the implications of cytoplasmic NIS immunostaining without concomitant detection of equivalent NIS mRNA levels remains speculative. Likewise, the physiological and clinical significance of extremely low tissue expression of NIS mRNA without unequivocal plasma membrane NIS immunostaining within a specific cell type remains uncertain.

NIS function in non-thyroidal tissues

NIS function in the thyroid gland is well defined. NIS mediates active iodide uptake against its concentration gradient from the circulating blood into thyroid follicular cells for thyroid hormone synthesis. Since iodine is a rare but essential element for thyroid hormone synthesis, the reabsorption of iodide is a critical step for maintaining a circulating pool. The iodine that has not yet been retained by the thyroid gland can be reabsorbed into the circulating blood and may be eventually taken up via NIS into thyroid follicular cells. Indeed, iodide is taken from circulating blood by NIS at salivary ducts and gastric mucosa into saliva and gastric juice, respectively. Iodide-containing saliva and gastric fluids that move into the small intestine are then resorbed into the bloodstream via NIS at the apical surface of enterocytes to complete entero-thyroid circulation of iodide (Josefsson *et al.* 2002). Taken together, NIS function in the salivary gland and gastrointestinal tract plays an important role in iodide conservation for thyroid hormone synthesis.

In the placenta, NIS in the apical membrane of cytotrophoblasts mediates iodide transport from maternal blood circulation to fetal blood circulation for thyroid hormone synthesis by the fetus. In lactating breast tissue, NIS in the basolateral membrane transports iodide from maternal blood circulation to maternal milk so that the newborn has sufficient iodine for his/her own thyroid hormone synthesis. Maternal iodine deficiency has been reported to have a negative impact on an infant's thyroid function (Dei-Tutu *et al.* 2020).

Additionally, it has been proposed that the function of NIS in non-thyroidal tissues is to provide antimicrobial protection and healing (Venturi & Venturi 2009).

However, the clinical impact of this possible role is unclear. The clinical phenotype for ITD patients with low or absent radioiodide uptake by the thyroid and other NIS-expressing organs has been limited to the consequence of thyroid hormone deficiency only. To date, there is no evidence to support increased susceptibility to oral or gastrointestinal infections nor pathological dysfunction in NIS-expressing non-thyroid organs for ITD patients. Similarly, the phenotype of NIS-knockout mice is characterized by hypothyroidism, with reduced thyroid hormone and increased thyroid-stimulating hormone levels (Ferrandino *et al.* 2017). Taken together, NIS function appears not to be critical in NIS-expressing non-thyroidal tissues beyond iodide conservation for thyroid hormone synthesis.

NIS regulation in non-thyroidal tissues

Among NIS-expressing tissues, NIS expression is generally elevated and restricted to highly functioning differentiated cells. In the thyroid, NIS is most prominent in the thyrocytes of patients with Graves' disease (Jhiang *et al.* 1998), when thyrocytes are actively synthesizing and releasing thyroid hormones. In breast tissue, NIS is most abundant during lactation (Cho *et al.* 2000, Wapnir *et al.* 2003) when providing iodine to the infant. In the stomach, NIS is restricted to the gastric mucosa (Wapnir *et al.* 2003) where gastric juice is secreted for food digestion. In salivary glands, NIS is most prominent in striated ducts (Jhiang *et al.* 1998, La Perle *et al.* 2013), which plays a major role in the modification of the ion composition of the primary saliva produced by acini. Taken together, it is reasonable to postulate that NIS modulation is tissue-specific and is likely modulated by the factor(s) driving lineage commitment to the highly functioning differentiated cells.

Among NIS-expressing non-thyroid tissues, the breast is the most extensively studied tissue for NIS modulation. Tazebay *et al.* (2000) showed that a s.c. injection of 1 IU oxytocin for 3 consecutive days significantly increased NIS expression and radioiodide uptake in the mammary gland of nubile mice. For ovariectomized nubile mice, the combination of oxytocin, prolactin and 17- β -estradiol stimulated maximal NIS induction. Our previous study demonstrates that radioiodide uptake in the lactating mammary glands of rats is partially decreased by a selective oxytocin antagonist or bromocriptine that inhibits prolactin release (Cho *et al.* 2000). In MCF-7 cells, an estrogen receptor-positive human breast cancer cell line, all-trans retinoic acid (ATRA) induces NIS expression and radioiodide uptake (Kogai *et al.* 2000). ATRA-induced NIS

expression can be further enhanced by co-treatment with dexamethasone (Kogai *et al.* 2005, Unterholzner *et al.* 2006). Additional factors that modulate NIS expression in ATRA+dexamethason-treated MCF-7 cells include miR-339-5p (Lakshmanan *et al.* 2015), a MEK inhibitor (Zhang *et al.* 2013), a staurosporine-related protein kinase inhibitor, KT5823 (Beyer *et al.* 2011), and a potent activator of pregnane-X-receptor, carbamazepine (Willhauck *et al.* 2011). A histone deacetylase inhibitor was reported to increase NIS mRNA in a variety of established cell lines (Liu & Xing 2012, Kelkar *et al.* 2016). Examining NIS expression in mammary gland tumors from various genetically engineered mouse models, cAMP or PI3K activation appears to associate with NIS up-regulation (Knostman *et al.* 2004).

In contrast to the breast, selective modulation of NIS expression in other non-thyroid tissues has not been investigated except that NIS expression is driven by lineage commitment to the highly functioning differentiated cells. La Perle *et al.* (2013) reported that inflammation decreases NIS expression in salivary striated ductal cells. It will be of interest to investigate if NIS expression in other non-thyroid tissues is also reduced by acute or chronic inflammation.

Similar to the thyroid, NIS expression is likely modulated by iodide concentration in non-thyroid tissues that facilitate dietary iodide absorption or iodide conservation for thyroid hormone synthesis. Indeed, excess iodide downregulates NIS protein levels and NIS-mediated iodide absorption in rat's small intestine as well as in small intestinal cell line (Nicola *et al.* 2009, 2012). This finding emphasizes the important role that dietary iodine plays in enterocyte physiology, as it controls its own NIS-mediated absorption and, by extension, manipulating the supply of iodine to the body (Nicola *et al.* 2015). NIS in gastric mucosa and salivary ductal cells mediates iodide uptake from circulating blood into epithelial cells such that iodide not taken by the thyroid can be released into gastric juice and saliva, respectively. Iodide returned to the gastrointestinal lumen can then be reabsorbed via NIS in small intestines. It would be of interest to investigate if NIS expression is modulated by iodide concentration in other non-thyroid tissues.

Impact of NIS expression in non-thyroidal tissues on thyroid cancer radioiodine therapy

NIS expression in non-thyroid tissues makes these tissues susceptible to radiation damage during radioiodine therapy for thyroid cancer patients. Thus, transiently

silencing/reducing NIS expression or inhibiting NIS function in non-thyroid tissues for 24–48 h after ^{131}I administration would prevent or alleviate ^{131}I -induced adverse effects. However, the proper mechanism to achieve transient, targeted, and restricted delivery of siRNA or a NIS inhibitor to target cells without compromising the ^{131}I therapeutic efficacy on thyroid cancer remains to be elucidated. Nevertheless, identifying the specific cell type in which NIS is abundantly expressed in non-thyroidal tissues has provided a better understanding of the development and progression of ^{131}I -induced adverse effects. For example, dry mouth is likely a secondary effect of salivary duct obstruction, as NIS expression is restricted to ductal cells. Indeed, loss of acinar cells and salivary gland atrophy occurs quickly after ligation of salivary excretory ducts, however, acinar cells can be regenerated after ductal ligation is removed in an experimental animal model (Cotroneo *et al.* 2010, Weng *et al.* 2018). Accordingly, if salivary duct obstruction could be recognized and relieved on time, not only could the discomfort of full-blown salivary gland swelling and pain be alleviated, ^{131}I -induced dry mouth could also potentially be prevented. Uncovering the pathological process and intervention at an early, potentially reversible stage is an alternate therapeutic intervention to reduce the severity of ^{131}I -induced adverse effects. Taken together, we are now in a better position to devise more effective strategies to alleviate the adverse effects of radioiodine therapy and thereby improve the quality of life for thyroid cancer survivors.

Among radioiodine adverse effects, salivary and lacrimal complications are relatively common and have a major negative impact on the quality of life for thyroid cancer survivors. As stated in a recent joint multidisciplinary clinical consensus statement of otolaryngology, ophthalmology, nuclear medicine and endocrinology (Singer *et al.* 2020), there is a need to better understand the true incidence, optimal treatments and importantly, prevention of salivary gland and lacrimal dysfunction elicited by radioiodine therapy. Since salivary duct and nasolacrimal drainage ductal systems are relatively accessible for local delivery of reagents, the incidence and severity of these adverse effects can be significantly reduced with collective effort from our community in pursuing personalized preventive measure and therapeutic intervention.

Radioiodine-induced salivary duct obstruction

Many thyroid cancer survivors suffer from life-long morbidity of radioiodine-induced salivary gland

dysfunction, including recurrent sialadenitis, persistent xerostomia, and progressive susceptibility to dental caries and periodontal disease. The prevalence of chronic symptomatic salivary dysfunction (16–54%) and abnormal salivary gland scintigraphic findings (37–72%) varies among studies (Clement *et al.* 2015). Various prevention strategies for ^{131}I -induced salivary gland (SG) dysfunction have been implemented, including salivary stimulation, radioprotection, and anti-inflammatory medications. Currently, the most common practice includes adequate hydration and initiation of salivary stimulation, such as sucking sour candy or lemon juice for the first 48 h after ^{131}I treatment. However, the efficacy and optimal timing of salivary stimulation have not been systematically evaluated and currently remain controversial (Nakada *et al.* 2005, Jentzen *et al.* 2010, Van Nostrand *et al.* 2010). The fact that chronic SG dysfunction is a common adverse effect among patients who have received high doses of ^{131}I indicates that novel approaches with better efficacy need to be developed.

Since the salivary accumulation of ^{131}I is absent in NIS-knockout mice and in patients with a congenital iodide transport defect who carry NIS loss-of-function mutation, one could hypothesize that ^{131}I -induced salivary gland damage can be avoided if salivary NIS expression/activity is temporarily inhibited during the time when blood circulating ^{131}I is high. The salivary ductal system, the primary injury site of ^{131}I , is accessible for local delivery of NIS silencing or NIS inhibition reagent. However, the challenge is to ensure that NIS silencing or NIS inhibition will be retained solely within the salivary glands without entering blood circulation to compromise radioiodine therapeutic efficacy for thyroid cancer. For ^{131}I -treated patients who suffer from radioiodine-induced sialadenitis with ductal stenosis and mucus plugs, sialendoscopy with dilatation, saline irrigation, and steroid injection have resulted in clinical improvement of symptoms for 75–100% of patients (Bhayani *et al.* 2015, Cung *et al.* 2017). Sialendoscopy interventions at an early stage of sialadenitis have a better outcome than patients who already have developed chronic salivary gland dysfunction. Of interest, sialendoscopy also shown to be effective in salivary gland protection by reducing the inflammatory effects of PSMA targeting alpha therapy in prostate cancer patients (Rathke *et al.* 2019). Since not all patients are at risk to develop ^{131}I -induced chronic salivary dysfunction, it is important to identify which patients will benefit from preventive measures prior to radioiodine therapy and which patients will require sialendoscopy intervention at an early stage of sialadenitis.

The incidence and severity of ^{131}I -induced salivary dysfunction in thyroid cancer survivors increase with cumulative ^{131}I radioactivity received (Clement *et al.* 2015). In addition to cumulative ^{131}I radioactivity, we found that female gender, previous incidence of sialadenitis, and personal history of autoimmune disease associated with Sjogren's syndrome are risk factors for the development of sialadenitis after ^{131}I therapy (Hollingsworth *et al.* 2016). Excluding patients with pre-existing salivary gland diseases, Horvath *et al.* (2020) reported ~25% of patients had anatomical damage in salivary gland(s) revealed by neck ultrasonography conducted before and after ^{131}I therapy. They found that the risk of sialadenitis significantly correlated with ^{131}I radioactivity administered and female gender. Salivary gland atrophy was found in 21, 47 and 78% of patients receiving 100, 150, and ≥ 200 mCi radioactivity, respectively. Therefore, the challenge is to identify individual patients who are susceptible to ^{131}I -induced chronic salivary dysfunction for prevention or early intervention.

The underlying pathobiological process associated with transient, recurrent, and chronic sialadenitis induced by ^{131}I has not been studied. This can be accomplished by conducting prospective studies investigating changes in answers to subjective questions related to the health status of the salivary gland (Le Roux *et al.* 2020), along with changes in objective evaluation of saliva composition (Klein Hesselink *et al.* 2016), such as saliva electrolyte and molecule profiles, scintigraphic functional imaging (Jeong *et al.* 2013), and neck ultrasound imaging (Horvath *et al.* 2020), prior to radioiodine therapy and at follow-up and then correlating these findings with the development of sialadenitis and/or dry mouth. By doing so, each patient can serve as their own control to identify the magnitude of changes in saliva composition and imaging characteristics by scintigraphy and/or ultrasonic images that lead to transient, recurrent, or chronic sialadenitis induced by ^{131}I . If we could use quantifiable saliva biomarkers or changes in salivary gland anatomic/functional images to monitor the onset and progression of salivary gland dysfunction induced by ^{131}I , intervention strategies can be applied to patients on time. We can investigate whether salivary biomarkers detected in the early stages of salivary dysfunction induced by ^{131}I are predictive of future progression to chronic disease. We may also be able to distinguish which patients are more susceptible to developing life-long ^{131}I -induced salivary gland dysfunction. The acquired insight on mechanisms underlying the development and progression of salivary gland dysfunction induced by ^{131}I can lead to alternative preventive measures and/or therapeutic interventions.

Radioiodine-induced nasolacrimal duct obstruction

The incidence of ^{131}I -induced nasolacrimal duct obstruction is 2–18% and typically occurs months after radioiodine therapy (Ali 2016). Preventive measures to reduce ^{131}I -induced nasolacrimal duct obstruction have not been proposed. A recent study demonstrated that a patient with pre-existing asymptomatic and incomplete nasolacrimal duct obstruction was able to be identified on the pre-therapy ^{123}I whole-body scan and SPECT/CT and further confirmed in post-therapy ^{131}I whole-body scan and SPECT/CT of head and neck (Kay *et al.* 2020). Thus, it is reasonable to assume that retrospective studies examining the pre-therapy and post-therapy whole-body scans along with SPECT/CT of head and neck from patients with multiple radioiodine treatments may help determine the prevalence of pre-existing nasolacrimal duct obstruction and ^{131}I -induced asymptomatic nasolacrimal duct obstruction among thyroid cancer patients. It would be possible to investigate whether additional ^{131}I therapy further exacerbates the severity of asymptomatic nasolacrimal duct obstruction, leading to symptomatic disease. With a better understanding of the development and progression of ^{131}I -induced nasolacrimal duct obstruction, we will be poised to devise appropriate prevention measures and intervention strategies.

Conclusion

NIS expression in non-thyroidal tissues allows radioiodine accumulation in these tissues after radioiodine administration. Consequently, these tissues are also vulnerable to radiation damage in ^{131}I -treated thyroid cancer patients. NIS expression in non-thyroidal tissues is generally restricted to or enriched in functioning differentiated cells, and tissue damage elicited by radioiodine is caused by direct or indirect radiation-induced inflammation. The severity and duration of adverse effects in these non-thyroidal tissues depend on the extent and duration of radioiodine accumulation, the tissue radiosensitivity, the turnover rate of radioiodine injured cells, and the individual tissue repair capacity. While NIS silencing in non-thyroidal tissues prior to ^{131}I administration can eliminate ^{131}I -induced adverse effects, additional technology is needed to ensure restricted delivery of NIS siRNA within targeted cells. Fully uncovering the pathological processes underlying ^{131}I -induced damage is critical for devising preventive measures and/or early therapeutic interventions that will facilitate complete resolution of radiation-induced inflammation. Finally,

the incidence and severity of ^{131}I -induced adverse effects vary among individual patients. Risk factors in developing long-term salivary dysfunction or nasolacrimal duct obstruction other than cumulative radioactivity administered need to be identified to aid in clinical management and decision-making.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was, in part, supported by National Institute of Health (NIH) Grant P50CA168505 (Project 3 leader: SM Jhiang, PI: MD Ringel) and the Ohio State University Comprehensive Cancer Center Bridge Fund.

Author contribution statement

SJ contributed to conceptualization, referencing, and writing of the draft. JS contributed to editing and providing clinical insight of the manuscript.

References

- Ali MJ 2016 Iodine-131 therapy and nasolacrimal duct obstructions: what we know and what we need to know. *Ophthalmic Plastic and Reconstructive Surgery* **32** 243–248. (<https://doi.org/10.1097/IOP.0000000000000647>)
- Altortay A, Dohán O, Szilágyi A, Paroder M, Wapnir IL & Carrasco N 2007 Expression of the Na⁺/I⁻ symporter (NIS) is markedly decreased or absent in gastric cancer and intestinal metaplastic mucosa of Barrett esophagus. *BMC Cancer* **7** 5. (<https://doi.org/10.1186/1471-2407-7-5>)
- Berger F, Unterholzner S, Diebold J, Knesewitsch P, Hahn K & Spitzweg C 2006 Mammary radioiodine accumulation due to functional sodium iodide symporter expression in a benign fibroadenoma. *Biochemical and Biophysical Research Communications* **349** 1258–1263. (<https://doi.org/10.1016/j.bbrc.2006.08.170>)
- Berman M, Hoff E, Barandes M, Becker DB, Sonenberg M, Benua R & Koutras DA 1968 Iodine kinetics in man—a model. *Journal of Clinical Endocrinology and Metabolism* **28** 1–14. (<https://doi.org/10.1210/jcem-28-1-1>)
- Beyer S, Lakshmanan A, Liu YY, Zhang X, Wapnir I, Smolenski A & Jhiang S 2011 KT5823 differentially modulates sodium iodide symporter expression, activity, and glycosylation between thyroid and breast cancer cells. *Endocrinology* **152** 782–792. (<https://doi.org/10.1210/en.2010-0782>)
- Bhayani MK, Acharya V, Kongkiatkamon S, Farah S, Roberts DB, Sterba J, Chambers MS & Lai SY 2015 Sialendoscopy for patients with radioiodine-induced sialadenitis and xerostomia. *Thyroid* **25** 834–838. (<https://doi.org/10.1089/thy.2014.0572>)
- Cho JY, Léveillé R, Kao R, Rousset B, Parlow AF, Burak WE Jr, Mazzaferri EL & Jhiang SM 2000 Hormonal regulation of radioiodide uptake activity and Na⁺/I⁻ symporter expression in mammary glands. *Journal of Clinical Endocrinology and Metabolism* **85** 2936–2943. (<https://doi.org/10.1210/jcem.85.8.6727>)
- Clement SC, Peeters RP, Ronckers CM, Links TP, van den Heuvel-Eibrink MM, Nieveen van Dijkum EJ, van Rijn RR, van der Pal HJ, Neggers SJ, Kremer LC, *et al.* 2015 Intermediate and long-term adverse effects of radioiodine therapy for differentiated thyroid carcinoma—a systematic review. *Cancer Treatment Reviews* **41** 925–934. (<https://doi.org/10.1016/j.ctrv.2015.09.001>)
- Cotroneo E, Proctor GB & Carpenter GH 2010 Regeneration of acinar cells following ligation of rat submandibular gland retraces the embryonic-perinatal pathway of cytodifferentiation. *Differentiation: Research in Biological Diversity* **79** 120–130. (<https://doi.org/10.1016/j.diff.2009.11.005>)
- Cung TD, Lai W, Svider PF, Hanba C, Samantray J, Folbe AJ, Shkoukani M & Raza SN 2017 Sialendoscopy in the management of radioiodine induced sialadenitis: a systematic review. *Annals of Otolaryngology, Rhinology, and Laryngology* **126** 768–773. (<https://doi.org/10.1177/0003489417732795>)
- da Fonseca FL, Yamanaka PK, Kato JM & Matayoshi S 2016 Lacrimal system obstruction after radioiodine therapy in differentiated thyroid carcinomas: a prospective comparative study. *Thyroid* **26** 1761–1767. (<https://doi.org/10.1089/thy.2015.0657>)
- Dai G, Levy O & Carrasco N 1996 Cloning and characterization of the thyroid iodide transporter. *Nature* **379** 458–460. (<https://doi.org/10.1038/379458a0>)
- de Santa Barbara P, van den Brink GR & Roberts DJ 2003 Development and differentiation of the intestinal epithelium. *Cellular and Molecular Life Sciences* **60** 1322–1332. (<https://doi.org/10.1007/s00018-003-2289-3>)
- Dei-Tutu SA, Manful A, Heimbürger DC, Malechi H, Moore DJ, Oppong SA, Russell WE & Aliyu MH 2020 Correlating maternal iodine status with neonatal thyroid function in two hospital populations in Ghana: a multicenter cross-sectional pilot study. *BMC Pediatrics* **20** 26. (<https://doi.org/10.1186/s12887-020-1932-6>)
- Fard-Esfahani A, Emami-Ardekani A, Fallahi B, Fard-Esfahani P, Beiki D, Hassanzadeh-Rad A & Eftekhari M 2014 Adverse effects of radioactive iodine-131 treatment for differentiated thyroid carcinoma. *Nuclear Medicine Communications* **35** 808–817. (<https://doi.org/10.1097/MNM.0000000000000132>)
- Ferrandino G, Kaspari RR, Reyna-Neyra A, Boutagy NE, Sinusas AJ & Carrasco N 2017 An extremely high dietary iodide supply forestalls severe hypothyroidism in Na⁺/I⁻ symporter (NIS) knockout mice. *Scientific Reports* **7** 5329. (<https://doi.org/10.1038/s41598-017-04326-z>)
- Fletcher A, Read ML, Thornton CEM, Larner DP, Poole VL, Brookes K, Nieto HR, Alshahrani M, Thompson RJ, Lavery GG, *et al.* 2020 Targeting novel sodium iodide symporter interactors ADP-ribosylation factor 4 and valosin-containing protein enhances radioiodine uptake. *Cancer Research* **80** 102–115. (<https://doi.org/10.1158/0008-5472.CAN-19-1957>)
- Goolden AW, Mallard JR & Farran HE 1957 Radiation sialitis following radioiodine therapy. *British Journal of Radiology* **30** 210–212. (<https://doi.org/10.1259/0007-1285-30-352-210>)
- Hammami MM & Bakheet S 1996 Radioiodine breast uptake in nonbreastfeeding women: clinical and scintigraphic characteristics. *Journal of Nuclear Medicine* **37** 26–31.
- Hollingsworth B, Senter L, Zhang X, Brock GN, Jarjour W, Nagy R, Brock P, Coombes KR, Kloos RT, Ringel MD, *et al.* 2016 Risk factors of ^{131}I -induced salivary gland damage in thyroid cancer patients. *Journal of Clinical Endocrinology and Metabolism* **101** 4085–4093. (<https://doi.org/10.1210/jc.2016-1605>)
- Horvath E, Skoknic V, Majlis S, Tala H, Silva C, Castillo E, Whittle C, Niedmann JP & González P 2020 Radioiodine-induced salivary gland damage detected by ultrasonography in patients treated for papillary thyroid cancer: radioactive iodine activity and risk. *Thyroid* **30** 1646–1655. (<https://doi.org/10.1089/thy.2019.0563>)
- Jentzen W, Balschuweit D, Schmitz J, Freudenberg L, Eising E, Hilbel T, Bockisch A & Stahl A 2010 The influence of saliva flow stimulation on the absorbed radiation dose to the salivary glands during radioiodine therapy of thyroid cancer using ^{124}I PET/(CT) imaging. *European Journal of Nuclear Medicine and Molecular Imaging* **37** 2298–2306. (<https://doi.org/10.1007/s00259-010-1532-z>)

- Jeong SY, Kim HW, Lee SW, Ahn BC & Lee J 2013 Salivary gland function 5 years after radioactive iodine ablation in patients with differentiated thyroid cancer: direct comparison of pre- and postablation scintigraphies and their relation to xerostomia symptoms. *Thyroid* **23** 609–616. (<https://doi.org/10.1089/thy.2012.0106>)
- Jhiang SM, Cho JY, Ryu KY, DeYoung BR, Smanik PA, McGaughy VR, Fischer AH & Mazzaferri EL 1998 An immunohistochemical study of Na⁺/I⁻ symporter in human thyroid tissues and salivary gland tissues. *Endocrinology* **139** 4416–4419. (<https://doi.org/10.1210/endo.139.10.6329>)
- Josefsson M, Grunditz T, Ohlsson T & Ekblad E 2002 Sodium/iodide-symporter: distribution in different mammals and role in entero-thyroid circulation of iodide. *Acta Physiologica Scandinavica* **175** 129–137. (<https://doi.org/10.1046/j.1365-201X.2002.00968.x>)
- Kay MD, Morris-Wiseman LE, Beazer A, Winegar BA & Kuo PH 2020 Primary nasolacrimal duct obstruction visualized on ¹²³I preablation scan for papillary thyroid carcinoma. *Journal of Nuclear Medicine Technology* **48** 77–78. (<https://doi.org/10.2967/jnmt.119.235010>)
- Kelkar MG, Senthilkumar K, Jadhav S, Gupta S, Ahn BC & De A 2016 Enhancement of human sodium iodide symporter gene therapy for breast cancer by HDAC inhibitor mediated transcriptional modulation. *Scientific Reports* **6** 19341. (<https://doi.org/10.1038/srep19341>)
- Kilbane MT, Ajjan RA, Weetman AP, Dwyer R, McDermott EW, O'Higgins NJ & Smyth PP 2000 Tissue iodine content and serum-mediated ¹²⁵I uptake-blocking activity in breast cancer. *Journal of Clinical Endocrinology and Metabolism* **85** 1245–1250. (<https://doi.org/10.1210/jcem.85.3.6442>)
- Kim MH, Kim HS & Park SA 2017 Extrathyroidal radioiodine accumulation in a fibroadenoma of the breast. *Clinical Nuclear Medicine* **42** e123–e125. (<https://doi.org/10.1097/RLU.0000000000001421>)
- Klein Hesselink EN, Brouwers AH, de Jong JR, van der Horst-Schrivers AN, Coppes RP, Lefrandt JD, Jager PL, Vissink A & Links TP 2016 Effects of radioiodine treatment on salivary gland function in patients with differentiated thyroid carcinoma: a prospective study. *Journal of Nuclear Medicine* **57** 1685–1691. (<https://doi.org/10.2967/jnumed.115.169888>)
- Kloos RT, Duvuuri V, Jhiang SM, Cahill KV, Foster JA & Burns JA 2002 Nasolacrimal drainage system obstruction from radioactive iodine therapy for thyroid carcinoma. *Journal of Clinical Endocrinology and Metabolism* **87** 5817–5820. (<https://doi.org/10.1210/jc.2002-020210>)
- Knostman KA, Cho JY, Ryu KY, Lin X, McCubrey JA, Hla T, Liu CH, Di Carlo E, Keri R, Zhang M, et al. 2004 Signaling through 3',5'-cyclic adenosine monophosphate and phosphoinositide-3 kinase induces sodium/iodide symporter expression in breast cancer. *Journal of Clinical Endocrinology and Metabolism* **89** 5196–5203. (<https://doi.org/10.1210/jc.2004-0907>)
- Kogai T, Schultz JJ, Johnson LS, Huang M & Brent GA 2000 Retinoic acid induces sodium/iodide symporter gene expression and radioiodide uptake in the MCF-7 breast cancer cell line. *PNAS* **97** 8519–8524. (<https://doi.org/10.1073/pnas.140217197>)
- Kogai T, Kanamoto Y, Li AI, Che LH, Ohashi E, Taki K, Chandraratna RA, Saito T & Brent GA 2005 Differential regulation of sodium/iodide symporter gene expression by nuclear receptor ligands in MCF-7 breast cancer cells. *Endocrinology* **146** 3059–3069. (<https://doi.org/10.1210/en.2004-1334>)
- La Perle KM, Kim DC, Hall NC, Bobbey A, Shen DH, Nagy RS, Wakely Jr PE, Lehman A, Jarjoura D & Jhiang SM 2013 Modulation of sodium/iodide symporter expression in the salivary gland. *Thyroid* **23** 1029–1036. (<https://doi.org/10.1089/thy.2012.0571>)
- Lakshmanan A, Wojcicka A, Kotlarek M, Zhang X, Jazdzewski K & Jhiang SM 2015 microRNA-339-5p modulates Na⁺/I⁻ symporter-mediated radioiodide uptake. *Endocrine-Related Cancer* **22** 11–21. (<https://doi.org/10.1530/ERC-14-0439>)
- Le Roux MK, Graillon N, Guyot L, Taieb D, Galli P, Godio-Raboulet Y, Chossegros C & Foletti JM 2020 Salivary side effects after radioiodine treatment for differentiated papillary thyroid carcinoma: long-term study. *Head and Neck* **42** 3133–3140. (<https://doi.org/10.1002/hed.26359>)
- Levy O, Dai G, Riedel C, Ginter CS, Paul EM, Lebowitz AN & Carrasco N 1997 Characterization of the thyroid Na⁺/I⁻ symporter with an anti-COOH terminus antibody. *PNAS* **94** 5568–5573. (<https://doi.org/10.1073/pnas.94.11.5568>)
- Liu Z & Xing M 2012 Induction of sodium/iodide symporter (NIS) expression and radioiodine uptake in non-thyroid cancer cells. *PLoS ONE* **7** e31729. (<https://doi.org/10.1371/journal.pone.0031729>)
- Mandel SJ & Mandel L 2003 Radioactive iodine and the salivary glands. *Thyroid* **13** 265–271. (<https://doi.org/10.1089/105072503321582060>)
- Mazzaferri EL & Jhiang SM 1994 Long-term impact of initial surgical and medical therapy on papillary and follicular thyroid cancer. *American Journal of Medicine* **97** 418–428. ([https://doi.org/10.1016/0002-9343\(94\)90321-2](https://doi.org/10.1016/0002-9343(94)90321-2))
- Mills JC & Shivdasani RA 2011 Gastric epithelial stem cells. *Gastroenterology* **140** 412–424. (<https://doi.org/10.1053/j.gastro.2010.12.001>)
- Mishra A & Shrivastava A 2020 Prognostic significance of sodium iodide symporter and deiodinase enzymes mRNA expression in gastric cancer. *International Journal of Applied and Basic Medical Research* **10** 43–48. (https://doi.org/10.4103/ijabmr.IJABMR_287_19)
- Morgenstern KE, Vadsyrisack DD, Zhang Z, Cahill KV, Foster JA, Burns JA, Kloos RT & Jhiang SM 2005 Expression of sodium iodide symporter in the lacrimal drainage system: implication for the mechanism underlying nasolacrimal duct obstruction in I(131)-treated patients. *Ophthalmic Plastic and Reconstructive Surgery* **21** 337–344. (<https://doi.org/10.1097/OI.iop.0000179369.75569.a8>)
- Myant NB 1960 Iodine metabolism of salivary glands. *Annals of the New York Academy of Sciences* **85** 208–214. (<https://doi.org/10.1111/j.1749-6632.1960.tb49959.x>)
- Nakada K, Ishibashi T, Takei T, Hirata K, Shinohara K, Katoh S, Zhao S, Tamaki N, Noguchi Y & Noguchi S 2005 Does lemon candy decrease salivary gland damage after radioiodine therapy for thyroid cancer? *Journal of Nuclear Medicine* **46** 261–266.
- Nicola JB, Basquin C, Portulano C, Reyna-Neyra A, Paroder M & Carrasco N 2009 The Na⁺/I⁻ symporter mediates active iodide uptake in the intestine. *American Journal of Physiology: Cell Physiology* **296** C654–C662. (<https://doi.org/10.1152/ajpcell.00509.2008>)
- Nicola JB, Reyna-Neyra A, Carrasco N & Masini-Repiso AM 2012 Dietary iodide controls its own absorption through post-transcriptional regulation of the intestinal Na⁺/I⁻ symporter. *Journal of Physiology* **590** 6013–6026. (<https://doi.org/10.1113/jphysiol.2012.241307>)
- Nicola JB, Carrasco N & Masini-Repiso AM 2015 Dietary I(-) absorption: expression and regulation of the Na(+)/I(-) symporter in the intestine. *Vitamins and Hormones* **98** 1–31. (<https://doi.org/10.1016/bs.vh.2014.12.002>)
- Peyrottes I, Navarro V, Ondo-Mendez A, Marcellin D, Bellanger L, Marsault R, Lindenthal S, Ettore F, Darcourt J & Pourcher T 2009 Immunoanalysis indicates that the sodium iodide symporter is not overexpressed in intracellular compartments in thyroid and breast cancers. *European Journal of Endocrinology* **160** 215–225. (<https://doi.org/10.1530/EJE-08-0505>)
- Rathke H, Kratochwil C, Hohenberger R, Giesel FL, Bruchertseifer F, Flechsig P, Morgenstern A, Hein M, Plinkert P, Haberkorn U, et al. 2019 Initial clinical experience performing sialendoscopy for salivary gland protection in patients undergoing ²²⁵Ac-PSMA-617 RLT. *European Journal of Nuclear Medicine and Molecular Imaging* **46** 139–147. (<https://doi.org/10.1007/s00259-018-4135-8>)
- Ryan J, Curran CE, Hennessy E, Newell J, Morris JC, Kerin MJ & Dwyer RM 2011 The sodium iodide symporter (NIS) and potential regulators in normal, benign and malignant human breast tissue. *PLoS ONE* **6** e16023. (<https://doi.org/10.1371/journal.pone.0016023>)
- Shiozaki A, Ariyoshi Y, Iitaka D, Kosuga T, Shimizu H, Kudou M, Konishi T, Shoda K, Arita T, Konishi H, et al. 2019 Functional analysis and clinical

- significance of sodium iodide symporter expression in gastric cancer. *Gastric Cancer* **22** 473–485. (<https://doi.org/10.1007/s10120-018-0874-2>)
- Singer MC, Marchal F, Angelos P, Bernet V, Boucai L, Buchholzer S, Burkey B, Eisele D, Erkul E, Faure F, *et al.* 2020 Salivary and lacrimal dysfunction after radioactive iodine for differentiated thyroid cancer: American Head and Neck Society Endocrine Surgery Section and Salivary Gland Section joint multidisciplinary clinical consensus statement of otolaryngology, ophthalmology, nuclear medicine and endocrinology. *Head and Neck* **42** 3446–3459. (<https://doi.org/10.1002/hed.26417>)
- Smanik PA, Liu Q, Furminger TL, Ryu K, Xing S, Mazzaferri EL & Jhiang SM 1996 Cloning of the human sodium iodide symporter. *Biochemical and Biophysical Research Communications* **226** 339–345. (<https://doi.org/10.1006/bbrc.1996.1358>)
- Smith VE, Read ML, Turnell AS, Watkins RJ, Watkinson JC, Lewy GD, Fong JC, James SR, Eggo MC, Boelaert K, *et al.* 2009 A novel mechanism of sodium iodide symporter repression in differentiated thyroid cancer. *Journal of Cell Science* **122** 3393–3402. (<https://doi.org/10.1242/jcs.045427>)
- Smith VE, Sharma N, Watkins RJ, Read ML, Ryan GA, Kwan PP, Martin A, Watkinson JC, Boelaert K, Franklyn JA, *et al.* 2013 Manipulation of PBF/PTTG1IP phosphorylation status: a potential new therapeutic strategy for improving radioiodine uptake in thyroid and other tumors. *Journal of Clinical Endocrinology and Metabolism* **98** 2876–2886. (<https://doi.org/10.1210/jc.2012-3640>)
- Spitzweg C, Joba W, Eisenmenger W & Heufelder AE 1998 Analysis of human sodium iodide symporter gene expression in extrathyroidal tissues and cloning of its complementary deoxyribonucleic acids from salivary gland, mammary gland, and gastric mucosa. *Journal of Clinical Endocrinology and Metabolism* **83** 1746–1751. (<https://doi.org/10.1210/jcem.83.5.4839>)
- Tazebay UH, Wapnir IL, Levy O, Dohan O, Zuckier LS, Zhao QH, Deng HF, Amenta PS, Fineberg S, Pestell RG, *et al.* 2000 The mammary gland iodide transporter is expressed during lactation and in breast cancer. *Nature Medicine* **6** 871–878. (<https://doi.org/10.1038/78630>)
- Unterholzner S, Willhauck MJ, Cengic N, Schütz M, Göke B, Morris JC & Spitzweg C 2006 Dexamethasone stimulation of retinoic acid-induced sodium iodide symporter expression and cytotoxicity of ¹³¹I in breast cancer cells. *Journal of Clinical Endocrinology and Metabolism* **91** 69–78. (<https://doi.org/10.1210/jc.2005-0779>)
- Van Nostrand D, Neutze J & Atkins F 1986 Side effects of "rational dose" iodine-131 therapy for metastatic well-differentiated thyroid carcinoma. *Journal of Nuclear Medicine* **27** 1519–1527.
- Van Nostrand D, Bandaru V, Chennupati S, Wexler J, Kulkarni K, Atkins F, Mete M & Gadwale G 2010 Radiopharmacokinetics of radioiodine in the parotid glands after the administration of lemon juice. *Thyroid* **20** 1113–1119. (<https://doi.org/10.1089/thy.2009.0429>)
- Venturi S & Venturi M 2009 Iodine in evolution of salivary glands and in oral health. *Nutrition and Health* **20** 119–134. (<https://doi.org/10.1177/026010600902000204>)
- Wapnir IL, van de Rijn M, Nowels K, Amenta PS, Walton K, Montgomery K, Greco RS, Dohán O & Carrasco N 2003 Immunohistochemical profile of the sodium/iodide symporter in thyroid, breast, and other carcinomas using high density tissue microarrays and conventional sections. *Journal of Clinical Endocrinology and Metabolism* **88** 1880–1888. (<https://doi.org/10.1210/jc.2002-021544>)
- Weng PL, Aure MH, Maruyama T & Ovitt CE 2018 Limited regeneration of adult salivary glands after severe injury involves cellular plasticity. *Cell Reports* **24** 1464–1470.e3. (<https://doi.org/10.1016/j.celrep.2018.07.016>)
- Willhauck MJ, O Kane DJ, Wunderlich N, Göke B & Spitzweg C 2011 Stimulation of retinoic acid-induced functional sodium iodide symporter (NIS) expression and cytotoxicity of ¹³¹I by carbamazepine in breast cancer cells. *Breast Cancer Research and Treatment* **125** 377–386. (<https://doi.org/10.1007/s10549-010-0835-x>)
- Wolff J 1983 Congenital goiter with defective iodide transport. *Endocrine Reviews* **4** 240–254. (<https://doi.org/10.1210/edrv-4-3-240>)
- Zhang Z, Beyer S & Jhiang SM 2013 MEK inhibition leads to lysosome-mediated Na⁺/I⁻ symporter protein degradation in human breast cancer cells. *Endocrine-Related Cancer* **20** 241–250. (<https://doi.org/10.1530/ERC-12-0342>)

Received in final form 1 April 2021

Accepted 20 April 2021

Accepted Manuscript published online 11 May 2021