

Supplemental Table 4. Incidence rates of colon NETs (# of cases per 100,000 population per year)

Reference	Country (region), registry	# of cases	Time period	Overall incidence	Incidence by gender and race	Comments
Godwin 1975	US, TNCS	n=27	1969–1971	N/A	1969–1971 WM: 0.11; WF: 0.07; BM: 0.15; BF: 0.07	N/A
Ballantyne <i>et al.</i> 1992	US (Connecticut), Connecticut Tumour Registry	n=54	1976–1986	0.31	N/A	Cecum: 48% Asc Colon: 16%
Modlin & Sandor 1997	US: SEER, TNCS	N/A	TNCS: 1969–1971 SEER: 1973–1991		1969–1971 WM: 0.11; WF: 0.07; BM: 0.15; BF: 0.07 1973–1991: WM: 0.10; WF: 0.13; BM: 0.21; BF: 0.19	Excluding appendix

Modlin <i>et al.</i> 2003	US, TNCS and SEER	n=1075	1969–1999 TNCS: 1969– 1971 SEER: 1973– 1999	N/A	1969–1971: WM: 0.11; WF: 0.07; BM: 0.15; BF: 0.07 1973–1991: WM: 0.11; WF: 0.13; BM: 0.23; BF: 0.20 1992–1999: WM: 0.21; WF: 0.18; BM: 0.38; BF: 0.28	Excluding appendix
Crocetti & Paci 2003	US, SEER	Cecum: n=238 Sigmoid: n=167 Colon NOS: n=48	1992–1998	Cecum: 0.06; sigmoid: 0.05; colon NOS: 0.01	N/A	N/A
Maggard <i>et</i>	US, SEER	n=661	1973–1997	1973: 0.10;	N/A	

<i>al.</i> 2004				1997: 0.20		
Gustafsson <i>et al.</i> 2008	US, SEER	N/A	1973–2004	1973: 0.08; 2004: 0.42	N/A	
Yao <i>et al.</i> 2008	US, SEER	N/A	2000–2004	Colon 0.20; cecum 0.16	Colon, M: 0.23; F: 0.17; W: 0.18; AA: 0.38; A/PI: 0.12; AI/AN: 0.22 Cecum, M: 0.16; F: 0.17; W: 0.17; AA: 0.21; A/PI: 0.04; AI/AN: 0.09	N/A
Lawrence <i>et al.</i> 2011	US, SEER	n=3530	1973–2007	1973–2007: 0.27 1973–1977: 0.15 1978–1982: 0.14 1983–1987: 0.17 1988–1992:	N/A	Excluding appendix Most common site in colon = cecum

				0.23 1993–1997: 0.28 1998–2002: 0.35 2003–2007: 0.40		
Hauso <i>et al.</i> 2008	US, SEER	N/A	1993–2004	W: 0.33; B: 0.53	WM: 0.35; WF: 0.32; BM: 0.58; BF: 0.50	N/A
			1993–1997	W: 0.27; B: 0.46	N/A	N/A
			2000–2004	W: 0.39; B: 0.64	N/A	N/A
	Norway, Norwegian registry of cancer	N/A	1993–2004	1993–2004: 0.25 1993–1997: 0.19 2000–2004:	1993–2004: M: 0.22; F: 0.28	N/A

				0.33		
Ellis <i>et al.</i> 2010	UK, NCR of England	n=658	1971–1978	N/A	M: 0.13; F:0.22	N/A
		n=123	1979–1984	N/A	M: 0.05; F:0.04	N/A
		n=139	1985–1989	N/A	M: 0.05; F:0.06	N/A
		n=177	1990–1994	N/A	M:0.07; F: 0.08	N/A
		n=389	1995–1999	N/A	M: 0.16; F: 0.16	N/A
		n=560	2000–2006	N/A	M: 0.17; F:0.17	N/A
Hemminki & Li 2001	Sweden, Swedish Family Cancer Database	n=601	1983–1998		M: 0.10; F: 0.20	N/A
Crocetti <i>et al.</i> 1997	Italy (Tuscany), Tuscany Tumour Registry RTT	n=8	1985–1991	0.05	N/A	Excluding appendix
Caldarella <i>et al.</i> 2011	Italy (Tuscany), Tuscany Cancer Registry RTT	N/A	1985–2005	0.30	M: 0.30; F: 0.40	N/A
Niederle <i>et al.</i> 2010	Austria, Pathology depts in Austria	n=14	2004–2005	0.06	N/A	Malignant only, including

						rectosigmoid junction
--	--	--	--	--	--	--------------------------

Abbreviations: TNCS, Third National Cancer Survey; SEER, Surveillance, Epidemiology, and End Results Database; WM, white male; WF, white female; BM, black male; BF, black female; N/A, not available; ERG, End Results Group; M, male; F, female; AA, AA, African American; A/PI, Asian Pacific Islander, AI/AN, American Indian/Alaskan native