

**Supplementary Table S2. Incidence of most common individual AEs considered treatment-related during the core and OLE studies among the OLE study population (i.e., 1 or more events occurring in  $\geq 5\%$  of patients)**

	LAN-LAN group (n=41)			PBO-LAN group (n=47)	
	Core study	OLE study	Core study and OLE studies (pooled)	Core study	OLE study
Diarrhoea	11 (26.8)	2 (4.9)	12 (29.3)	4 (8.5)	10 (21.3)
Abdominal pain	7 (17.1)	0	7 (17.1)	1 (2.1)	1 (2.1)
Cholelithiasis	4 (9.8)	3 (7.3)	6 (14.6)	3 (6.4)	1 (2.1)
Hyperglycemia	3 (7.3)	1 (2.4)	4 (9.8)	0	0
Flatulence	3 (7.3)	1 (2.4)	3 (7.3)	2 (4.3)	0
Injection site pain	2 (4.9)	1 (2.4)	3 (7.3)	1 (2.1)	1 (2.1)
Steatorrhea	2 (4.9)	0	2 (4.9)	0	3 (6.4)

Data are number (%) of patients with an AE while receiving lanreotide Autogel 120 mg. Adverse events were defined according to the Medical Dictionary for Regulatory Activities version 16.0.

LAN-LAN group, patients receiving lanreotide Autogel in core study as well as the OLE study; OLE, open-label extension; PBO-LAN, patients receiving placebo in the core study before crossing over to lanreotide in the OLE study.