

**Table S2:** Frequencies of metastatic thyroid carcinomas included in this study, distributed by body location.

	<b>PTC</b>	<b>FTC</b>	<b>HCC</b>	<b>PDTC</b>	<b>ATC</b>	<b>MTC</b>
<b>Location</b>						
Lymph Node	80 (50%)	0	5 (37%)	18 (32%)	3 (33%)	4 (67%)
Lung	33 (21%)	0	2 (14%)	7 (12%)	0	0
Bone	14 (9%)	2 (67%)	1 (7%)	8 (14%)	0	0
Neck	9 (6%)	0	2 (14%)	2 (4%)	1 (11%)	0
Mediastinum	2 (1%)	0	0	2 (4%)	1 (11%)	0
Brain	2 (1%)	0	0	3 (5%)	1 (11%)	0
Liver	1 (0.5%)	0	0	1 (2%)	1 (11%)	1 (17%)
Pleura	1 (0.5%)	0	0	2 (4%)	0	0
Esophagus	1 (0.5%)	0	0	0	0	0
Soft Tissue	1 (0.5%)	0	0	0	0	0
Rib	1 (0.5%)	0	0	0	0	0
Adrenal gland	1 (0.5%)	0	0	0	0	0
Kidney	1 (0.5%)	0	0	0	0	0
Chest Wall	0	1 (33%)	0	0	1 (11%)	1 (17%)
Trachea	0	0	1 (7%)	0	0	0
Larynx	0	0	1 (7%)	0	0	0
Thigh	0	0	1 (7%)	0	0	0
Skull	0	0	1 (7%)	0	0	0
Skin	0	0	0	1 (2%)	0	0
Paratracheal	0	0	0	0	1 (11%)	0
Pelvis	0	0	0	0	0	0
Retroperitoneum	0	0	0	0	0	0
Unspecified	15 (9%)	0	0	13 (23%)	0	0